
All new STI episodes seen at GUM clinics in the UK: 1999 - 2008



Data is represented by country, gender and sexual orientation.

Notes on using these tables:

Numbers of diagnoses are adjusted for missing clinic data

Data Source:

England: GUM clinic KC60 returns
Wales: GUM clinic KC60 returns
Northern Ireland: GUM clinic KC60 returns
Scotland: GUM clinic ISD(D)5 returns from 31/03/2004 and STISS returns from 01/04/2004

Data Supplied by:

Health Protection Agency, Centre for Infections
Communicable Disease Surveillance Centre Wales
Communicable Disease Surveillance Centre Northern Ireland
Regional Information Branch, Department of Health, Social Services and Public Safety (DHSS&PS Northern Ireland)
ISD Scotland, Information Services, NHS National Services Scotland

Definitions of selected data totals:

Total new STI diagnoses: refer to 'Code Groups' for included KC60 conditions
Total other STI diagnoses: refer to 'Code Groups' for included KC60 conditions
Total other GUM clinic diagnoses: refer to 'Code Groups' for included KC60 conditions
Total services provided: refer to 'Code Groups' for included KC60 conditions

Data prepared by:

Microbiology and Epidemiology of STIs and HIV (MESH) Department
Health Protection Agency
Centre for Infections
61 Colindale Avenue
London
NW9 5EQ

Tel. 020-8200-6868
Email: hiv/sti@hpa.org.uk
Website: www.hpa.org.uk

Data tables

United Kingdom

England

Code Groups

Further Information

England

Data on selected diagnoses, 1971 to 2008, is available upon request

HPA Centre for Infections

Contact: Geraldine Leong

Tel: +44 (0)20 8200 6868

Email: geraldine.leong@hpa.org.uk

Web: <http://www.hpa.org.uk>

Scotland

Data on selected diagnoses, 1956 to 2008, is available upon request

ISD Scotland

Contact: Kenny McIntyre

Tel: +44(0)131 275 6557

Email: k.mcintyre@nhs.net

Web: www.isdscotland.org.uk

Wales

Data on selected diagnoses is available upon request.

National Public Health Service for Wales

Contact: Daniel Thomas

Tel: +44(0)29 2040 2472

email: daniel.thomas@nphs.wales.nhs.uk

Web: <http://www.nphs.wales.nhs.uk>

Northern Ireland

Data on selected diagnoses is available upon request.

Communicable Disease Surveillance Centre Northern Ireland

Contact: Brian Smyth

Tel: +44(0)2890 263 765

email: dbrian.smyth@hpa.org.uk

Web: www.hpa.org.uk



United Kingdom:

Number of new STI episodes seen at genitourinary medicine clinics by sex, 1999 - 2008

Diagnoses and/or treatment of infection or disease			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
A1,A2	Primary and secondary infectious syphilis	Male (all)	166	264	642	1,118	1,461	2,000	2,362	2,393	2,344	2,248
		-of which MSM	53	130	365	653	826	1,074	1,437	1,431	1,419	1,390
		Female	57	78	111	139	191	283	362	290	289	276
		Total	223	342	753	1,257	1,652	2,283	2,724	2,683	2,633	2,524
A3	Early latent syphilis (first 2 years)	Male (all)	127	134	275	363	459	571	717	751	903	900
		-of which MSM	31	51	118	194	211	301	427	467	534	535
		Female	79	93	104	129	156	201	192	199	172	164
		Total	206	227	379	492	615	772	909	950	1,075	1,064
A4,A5,A6	Other acquired syphilis	Male (all)	697	807	882	909	1,051	1,157	1,156	1,156	1,143	1,094
		-of which MSM	97	96	128	140	166	171	209	257	259	299
		Female	421	476	652	777	830	897	877	759	850	783
		Total	1,118	1,283	1,534	1,686	1,881	2,054	2,033	1,915	1,993	1,877
A7	Congenital syphilis, aged under 2	Male (all)	1	6	2	2	3	5	6	3	1	4
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	4	0	5	0	3	8	3	3	0
Total	2	10	2	7	3	8	14	6	4	4		
A8	Congenital syphilis, aged 2 or over	Male (all)	7	9	4	7	1	6	9	4	4	3
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	14	4	6	9	5	6	14	5	5	2
Total	21	13	10	16	6	12	23	9	9	5		
A9	Epidemiological treatment of suspected syphilis	Male (all)	31	27	64	168	270	423	535	699	783	879
		-of which MSM	7	6	25	86	161	255	324	451	503	578
		Female	35	26	30	52	70	95	126	132	164	148
Total	66	53	94	220	340	518	661	831	947	1,027		
B1,B2	Uncomplicated gonorrhoea	Male (all)	11,326	15,234	16,626	17,868	17,409	15,677	13,887	13,509	12,838	11,113
		-of which MSM	1,966	3,149	3,741	3,618	4,077	3,994	4,391	4,573	3,765	3,316
		Female	5,144	6,566	7,079	7,723	7,564	6,649	5,378	5,354	5,811	5,516
		Total	16,470	21,800	23,705	25,591	24,973	22,326	19,265	18,863	18,649	16,629
B3	Gonococcal ophthalmia neonatorum	Male (all)	6	7	7	8	10	10	12	21	13	15
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	7	6	10	13	16	15	81	9	16	24
Total	13	13	17	21	26	25	93	30	29	39		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	1,641	2,282	2,772	3,179	3,514	3,713	3,852	4,064	4,547	4,736
		-of which MSM	454	648	868	931	1,165	1,192	1,367	1,482	1,440	1,461
		Female	1,907	2,319	2,705	3,115	3,304	3,269	3,317	3,221	3,530	3,842
Total	3,548	4,601	5,477	6,294	6,818	6,982	7,169	7,285	8,077	8,578		
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	81	85	101	113	110	138	105	93	99	100
		-of which MSM	13	9	14	26	13	19	31	21	18	20
		Female	247	278	287	283	327	277	234	251	266	248
Total	328	363	388	396	437	415	339	344	365	348		

UK: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	34	32	30	38	65	88	262	220	265	284
		-of which MSM	0	0	0	0	0	0	1	3	6	4
		Female	24	33	36	39	83	85	67	79	135	94
		Total	58	65	66	77	148	173	329	299	400	378
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	24,435	29,818	33,362	38,676	43,188	48,625	51,976	56,231	60,807	61,863
		-of which MSM	690	1,082	1,531	1,646	2,113	2,228	2,713	3,364	3,699	4,327
		Female	32,556	38,514	43,153	48,912	52,971	56,114	57,530	57,482	60,984	61,155
		Total	56,991	68,332	76,515	87,588	96,159	104,739	109,506	113,713	121,791	123,018
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	480	569	610	632	699	724	810	847	875	1,006
		-of which MSM	26	25	40	35	54	49	74	70	74	89
		Female	1,959	2,365	2,411	2,462	2,426	2,617	2,534	2,486	2,803	2,828
		Total	2,439	2,934	3,021	3,094	3,125	3,341	3,344	3,333	3,678	3,834
C4D	Chlamydial ophthalmia neonatorum	Male (all)	18	20	15	18	23	49	22	17	13	19
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	13	29	29	26	26	41	23	23	11	21
		Total	31	49	44	44	49	90	45	40	24	40
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	12,823	15,700	19,471	23,941	25,173	27,420	29,310	28,923	32,284	33,195
		-of which MSM	277	479	726	1,147	1,491	1,613	1,913	2,217	2,480	2,640
		Female	9,221	11,202	13,291	15,673	16,317	18,691	20,720	20,957	22,922	24,231
		Total	22,044	26,902	32,762	39,614	41,490	46,111	50,030	49,880	55,206	57,426
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	57,963	60,933	64,414	67,295	72,585	75,903	76,252	75,424	73,299	69,218
		-of which MSM	3,603	4,089	4,839	4,841	5,288	5,441	5,547	6,081	5,611	5,374
		Female	197	160	129	134	7,041	7,880	7,989	7,500	7,311	7,076
		Total	58,160	61,093	64,543	67,429	79,626	83,783	84,241	82,924	80,610	76,294
C4I	Epidemiological treatment of NSGI	Male (all)	6,413	6,497	6,363	6,287	6,441	6,598	6,134	6,094	6,378	6,470
		-of which MSM	733	646	723	670	800	829	775	870	779	818
		Female	19,880	20,410	21,161	20,122	18,236	18,613	18,324	17,410	17,146	15,945
		Total	26,293	26,907	27,524	26,409	24,677	25,211	24,458	23,504	23,524	22,415
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	3,341	3,865	4,045	4,214	4,280	4,675	5,427	5,099	5,597	6,199
		-of which MSM	221	216	248	239	257	286	368	354	378	498
		Female	10,790	11,431	11,534	11,964	11,386	13,243	12,718	13,140	14,904	15,560
		Total	14,131	15,296	15,579	16,178	15,666	17,918	18,145	18,239	20,501	21,759
C6A	Trichomoniasis	Male (all)	670	747	775	612	318	347	387	368	435	403
		-of which MSM	0	0	0	0	0	0	0	1	0	0
		Female	5,640	6,196	6,479	6,804	6,388	6,093	5,257	5,384	5,522	5,405
		Total	6,310	6,943	7,254	7,416	6,706	6,440	5,644	5,752	5,957	5,808
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	1,512	1,455	1,531	1,488	1,534	1,534	1,601	1,822	1,848	2,122
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	59,202	64,807	69,403	76,132	77,435	75,413	77,529	80,237	92,364	97,264
		Total	60,714	66,262	70,934	77,620	78,969	76,947	79,130	82,059	94,212	99,386
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	13,632	13,605	13,745	13,482	13,629	13,577	13,064	12,699	14,491	15,452
		-of which MSM	58	40	59	67	72	61	64	65	73	101
		Female	7,817	7,314	7,062	5,895	5,657	6,062	5,107	4,492	4,780	5,320
		Total	21,449	20,919	20,807	19,377	19,286	19,639	18,171	17,191	19,271	20,772

UK: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
C7A	Anogenital candidosis	Male (all)	8,334	8,055	8,359	7,885	6,954	6,777	6,608	7,493	8,643	9,203
		-of which MSM	11	13	13	6	15	19	10	26	14	23
		Female	68,546	70,202	74,608	75,275	71,371	67,929	64,829	63,935	70,675	75,512
		Total	76,880	78,257	82,967	83,160	78,325	74,706	71,437	71,428	79,318	84,715
C7B	Epidemiological treatment of C6 & C7	Male (all)	3,688	3,386	3,761	4,276	4,332	4,285	4,439	4,227	4,763	5,146
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	6,468	5,807	6,704	6,741	7,133	8,004	7,385	7,298	8,075	9,169
		Total	10,156	9,193	10,465	11,017	11,465	12,289	11,824	11,525	12,838	14,315
C8,C9	Scabies/pediculosis pubis	Male (all)	5,044	5,225	4,687	4,091	3,524	2,799	2,231	2,179	2,058	1,970
		-of which MSM	830	951	931	834	724	600	557	587	513	535
		Female	1,452	1,296	1,062	825	593	456	342	297	304	262
		Total	6,496	6,521	5,749	4,916	4,117	3,255	2,573	2,476	2,362	2,232
C10A	Anogenital herpes simplex – first attack	Male (all)	6,660	6,812	7,200	7,218	7,209	7,237	7,606	8,481	10,110	11,273
		-of which MSM	372	439	447	525	523	525	626	615	695	846
		Female	10,849	11,011	11,744	12,220	12,024	11,836	12,242	13,396	16,160	17,684
		Total	17,509	17,823	18,944	19,438	19,233	19,073	19,848	21,877	26,270	28,957
C10B	Anogenital herpes simplex – recurrence	Male (all)	6,428	6,590	6,770	6,965	6,940	6,717	6,915	7,270	7,821	8,688
		-of which MSM	422	463	438	440	503	510	568	589	636	720
		Female	7,741	7,891	8,691	8,822	8,702	8,429	8,719	9,080	10,444	11,673
		Total	14,169	14,481	15,461	15,787	15,642	15,146	15,634	16,350	18,265	20,361
C11A	Anogenital warts – first attack	Male (all)	37,482	37,633	38,522	39,887	40,601	42,727	43,021	44,249	47,140	48,744
		-of which MSM	1,760	1,877	1,987	2,231	2,397	2,396	2,593	2,664	2,757	2,760
		Female	34,266	33,684	34,936	35,082	35,998	37,332	38,292	39,367	42,375	43,781
		Total	71,748	71,317	73,458	74,969	76,599	80,059	81,313	83,616	89,515	92,525
C11B	Anogenital warts – recurrence	Male (all)	25,385	25,420	25,620	24,957	24,718	25,189	26,457	27,932	30,540	31,629
		-of which MSM	1,325	1,337	1,484	1,513	1,693	1,801	1,743	1,914	2,137	2,052
		Female	13,861	14,051	13,869	13,742	14,188	14,622	15,635	17,016	19,015	20,053
		Total	39,246	39,471	39,489	38,699	38,906	39,811	42,092	44,948	49,555	51,682
C11C	Anogenital warts – re-registered cases	Male (all)	10,435	9,992	9,864	9,272	9,535	9,261	9,516	10,090	10,121	10,630
		-of which MSM	51	40	21	32	28	19	37	67	69	80
		Female	8,535	8,036	7,869	7,107	7,202	7,126	7,443	7,580	7,700	7,623
		Total	18,970	18,028	17,733	16,379	16,737	16,387	16,959	17,670	17,821	18,253
C12	Molluscum contagiosum	Male (all)	5,186	5,594	6,049	6,704	7,544	8,728	9,705	10,602	11,689	12,298
		-of which MSM	238	221	243	231	273	285	325	351	364	354
		Female	2,897	2,955	3,277	3,512	3,815	4,277	4,684	4,972	5,394	5,433
		Total	8,083	8,549	9,326	10,216	11,359	13,005	14,389	15,574	17,083	17,731
C13	Antigen positive viral hepatitis B	Male (all)	565	669	748	962	23	31	17	25	40	46
		-of which MSM	109	135	165	203	9	15	6	3	15	10
		Female	264	238	364	496	6	7	3	7	13	8
		Total	829	907	1,112	1,458	29	38	20	32	53	54
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	-	-	619	672	655	651	702	745
		-of which MSM	-	-	-	-	114	93	91	62	58	68
		Female	-	-	-	-	276	328	329	345	361	396
		Total	-	-	-	-	895	1,000	984	996	1,063	1,141

UK: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	-	-	47	55	48	43	48	63
		-of which MSM	-	-	-	-	14	19	14	8	6	18
		Female	-	-	-	-	11	12	30	26	22	14
		Total	-	-	-	-	58	67	78	69	70	77
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	-	-	348	462	419	437	494	567
		-of which MSM	-	-	-	-	121	205	207	230	147	167
		Female	-	-	-	-	151	169	165	197	196	244
		Total	-	-	-	-	499	631	584	634	690	811
C14*	Viral hepatitis C: first diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	957	980	955	1,096	1,053	1,116	1,029	963	895	964
		-of which MSM	83	106	105	145	110	126	123	128	86	133
		Female	454	445	396	503	489	472	447	399	332	378
		Total	1,411	1,425	1,351	1,599	1,542	1,588	1,476	1,362	1,227	1,342
E1A	New HIV diagnosis - asymptomatic	Male (all)	1,125	1,244	1,713	2,365	2,737	2,922	3,152	2,938	3,643	3,263
		-of which MSM	551	624	770	772	1,134	1,299	1,500	1,409	1,588	1,394
		Female	589	691	1,137	2,012	2,154	2,116	2,099	1,769	2,293	1,723
		Total	1,714	1,935	2,850	4,377	4,891	5,038	5,251	4,707	5,936	4,986
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	7,502	8,173	8,758	10,387	198	629	832	915	980	1,098
		-of which MSM	557	122	76	47	84	426	656	691	775	803
		Female	2,280	3,006	3,872	5,630	56	222	202	307	312	350
		Total	9,782	11,179	12,630	16,017	254	851	1,034	1,222	1,292	1,448
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	-	-	26,102	31,372	38,437	39,964	40,140	42,380
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	-	-	12,318	16,561	20,823	21,836	23,280	24,249
		Total	-	-	-	-	38,420	47,933	59,260	61,800	63,420	66,629
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	548	513	684	772	695	701	767	662	511	998
		-of which MSM	294	244	330	265	278	263	276	253	170	320
		Female	198	210	335	457	459	402	454	370	313	393
		Total	746	723	1,019	1,229	1,154	1,103	1,221	1,032	824	1,391
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	8,689	9,421	9,508	9,597	58	433	502	467	481	563
		-of which MSM	521	104	7	8	38	328	405	357	350	434
		Female	2,065	2,376	2,699	3,381	8	107	79	76	98	120
		Total	10,754	11,797	12,207	12,978	66	540	581	543	579	683
E3A*	AIDS – first presentation	Male (all)	414	383	448	468	3	4	5	7	7	6
		-of which MSM	167	189	175	177	0	3	1	4	6	1
		Female	112	141	217	237	0	2	0	2	0	3
		Total	526	524	665	705	3	6	5	9	7	9
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	-	-	182	222	232	190	241	224
		-of which MSM	-	-	-	-	73	62	77	64	75	68
		Female	-	-	-	-	143	120	150	89	97	117
		Total	-	-	-	-	325	342	382	279	338	341
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	-	-	240	156	162	160	111	109
		-of which MSM	-	-	-	-	97	76	61	82	48	57
		Female	-	-	-	-	135	65	70	70	50	55
		Total	-	-	-	-	375	221	232	230	161	164

UK: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
E3B	AIDS – subsequent presentation	Male (all)	6,060	6,625	7,046	7,958	9,761	10,582	11,985	11,921	11,469	11,799
		-of which MSM	291	87	21	8	13	81	106	152	150	180
		Female	1,381	1,683	2,125	2,666	4,060	4,418	5,044	5,230	5,430	5,589
		Total	7,441	8,308	9,171	10,624	13,821	15,000	17,029	17,151	16,899	17,388
D2A	Urinary tract infection	Male (all)	2,138	2,010	2,101	2,625	2,626	2,588	2,135	2,107	2,315	2,708
		-of which MSM	6	3	8	6	2	4	7	8	8	19
		Female	9,349	9,642	9,743	9,480	9,042	10,119	9,686	10,079	12,573	14,780
		Total	11,487	11,652	11,844	12,105	11,668	12,707	11,821	12,186	14,888	17,488
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	56,498	58,716	58,976	62,338	69,190	68,597	70,712	72,440	83,082	94,006
		-of which MSM	1,025	797	682	1,021	1,243	509	391	396	410	437
		Female	51,699	56,342	57,158	58,182	56,791	57,674	58,626	60,101	67,326	75,334
		Total	108,197	115,058	116,134	120,520	125,981	126,271	129,338	132,541	150,408	169,340
D3	Other episodes not requiring treatment	Male (all)	112,394	118,230	139,176	158,614	151,051	183,482	207,094	229,362	261,242	291,328
		-of which MSM	1,113	874	1,043	1,399	1,556	3,398	4,902	5,071	5,936	6,716
		Female	120,252	123,430	143,377	161,215	150,515	176,853	204,275	223,967	262,688	292,673
		Total	232,646	241,660	282,553	319,829	301,566	360,335	411,369	453,329	523,930	584,001
P1A	HIV antibody test (no sexual health screen)	Male (all)	79,819	90,280	117,896	156,922	58,621	57,271	42,333	36,257	34,525	34,078
		-of which MSM	12,041	13,340	17,432	20,186	10,010	9,695	8,272	7,834	7,426	7,358
		Female	63,346	71,831	98,559	138,304	44,666	41,445	31,723	27,085	26,339	26,053
		Total	143,165	162,111	216,455	295,226	103,287	98,716	74,056	63,342	60,864	60,131
P1B	HIV antibody test offered and refused	Male (all)	13,825	15,220	20,396	45,408	100,363	127,092	127,315	122,701	126,857	132,235
		-of which MSM	1,492	1,234	1,765	3,335	6,188	6,973	7,196	7,232	6,898	6,975
		Female	13,086	14,236	20,035	50,304	114,577	143,218	145,059	141,854	155,686	169,188
		Total	26,911	29,456	40,431	95,712	214,940	270,310	272,374	264,555	282,543	301,423
P2	Hepatitis B vaccination	Male (all)	10,324	11,277	12,427	13,996	14,967	16,659	16,502	17,054	17,842	19,525
		-of which MSM	4,892	5,387	6,114	6,961	8,095	8,779	8,970	8,924	8,847	9,636
		Female	2,574	2,718	3,243	4,008	4,222	5,252	5,636	5,986	6,626	7,490
		Total	12,898	13,995	15,670	18,004	19,189	21,911	22,138	23,040	24,468	27,015
P3	Contraception (excluding condom provision)	Male (all)	0	0	0	0	0	12	82	156	466	1,601
		-of which MSM	0	0	0	0	0	2	41	35	46	165
		Female	30,705	33,389	34,075	31,923	33,102	34,664	35,861	39,818	49,049	59,730
		Total	30,705	33,389	34,075	31,923	33,102	34,664	35,861	39,818	49,049	59,730
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	8,060	6,509	4,926	3,849	5,358	3,450	1,770	1,497	1,516	1,776
		Total	8,060	6,509	4,926	3,849	5,358	3,450	1,770	1,497	1,516	1,776

UK: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1,231	1,146	1,033	766	924	813	398	305	269	360
		Total	1,231	1,146	1,033	766	924	813	398	305	269	360
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	-	-	151,572	153,409	148,078	141,861	143,727	141,544
		-of which MSM	-	-	-	-	9,952	10,083	11,334	12,605	12,318	12,592
		Female	-	-	-	-	185,799	184,799	180,450	172,289	182,599	186,747
		Total	-	-	-	-	337,371	338,208	328,528	314,150	326,326	328,291
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	-	-	163,154	214,136	281,948	330,660	397,655	446,728
		-of which MSM	-	-	-	-	16,313	19,549	25,998	31,424	35,132	41,440
		Female	-	-	-	-	159,738	207,426	272,944	318,192	390,675	444,289
		Total	-	-	-	-	322,892	421,562	554,892	648,852	788,330	891,017
Total new STI diagnoses												
	Male (all)	154,666	168,694	179,732	191,948	203,047	214,059	218,869	224,185	232,783	232,031	
	-of which MSM	10,647	13,105	15,601	16,099	18,231	18,804	20,919	22,266	21,603	21,764	
	Female	106,942	115,561	123,809	132,692	143,715	149,977	150,519	152,416	165,126	167,707	
	Total	261,608	284,255	303,541	324,640	346,762	364,036	369,388	376,601	397,909	399,738	
Total other STI diagnoses												
	Male (all)	86,552	91,966	97,597	104,140	114,059	123,749	135,877	139,758	146,907	153,390	
	-of which MSM	4,903	4,219	4,695	5,210	6,249	7,322	8,189	9,175	9,695	10,189	
	Female	67,476	71,660	77,230	81,382	85,476	93,182	101,510	103,724	110,983	114,719	
	Total	154,028	163,626	174,827	185,522	199,535	216,931	237,387	243,482	257,890	268,109	
Total other GUM clinic diagnoses												
	Male (all)	87,324	88,877	90,176	94,153	100,356	99,694	100,727	102,907	117,321	131,022	
	-of which MSM	1,292	1,094	1,032	1,448	1,700	1,051	913	926	817	976	
	Female	213,090	222,452	231,398	237,319	234,645	230,456	226,304	228,919	258,502	280,555	
	Total	300,414	311,329	321,574	331,472	335,001	330,150	327,031	331,826	375,823	411,577	
Total services provided												
	Male (all)	216,362	235,007	289,895	374,940	639,728	752,061	823,352	878,051	982,314	1,067,039	
	-of which MSM	19,538	20,835	26,354	31,881	52,114	58,479	66,713	73,125	76,603	84,882	
	Female	229,963	245,604	299,289	385,754	692,619	793,657	875,948	929,191	1,073,662	1,186,170	
	Total	446,325	480,611	589,184	760,694	1,332,347	1,545,718	1,699,300	1,807,242	2,055,976	2,253,209	

Source: HPA Centre for Infections

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11a or C11b treated for more than 3 months

- data not available/applicable



England:

Number of new STI episodes seen at genitourinary medicine clinics by sex, 1999 - 2008

Diagnoses and/or treatment of infection or disease			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
A1,A2	Primary and secondary infectious syphilis	Male (all)	156	247	618	1,059	1,355	1,768	2,146	2,144	2,110	1,965
		-of which MSM	52	119	351	613	759	903	1,268	1,233	1,248	1,185
		Female	55	75	99	137	183	266	348	277	278	253
		Total	211	322	717	1,196	1,538	2,034	2,494	2,421	2,388	2,218
A3	Early latent syphilis (first 2 years)	Male (all)	122	126	268	330	425	526	666	667	805	786
		-of which MSM	30	48	114	177	194	271	387	411	453	439
		Female	75	93	99	121	150	197	178	175	160	149
		Total	197	219	367	451	575	723	844	842	965	935
A4,A5,A6	Other acquired syphilis	Male (all)	671	778	847	866	1,017	1,111	1,078	1,091	1,065	1,017
		-of which MSM	93	87	122	130	157	157	179	241	220	265
		Female	405	470	638	746	805	864	841	719	807	741
		Total	1,076	1,248	1,485	1,612	1,822	1,975	1,919	1,810	1,872	1,758
A7	Congenital syphilis, aged under 2	Male (all)	1	5	2	2	3	4	6	3	1	4
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	4	0	5	0	3	8	3	3	0
Total	2	9	2	7	3	7	14	6	4	4		
A8	Congenital syphilis, aged 2 or over	Male (all)	5	9	3	5	1	5	9	4	3	3
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	12	4	6	9	4	4	13	4	5	2
Total	17	13	9	14	5	9	22	8	8	5		
A9	Epidemiological treatment of suspected syphilis	Male (all)	24	26	61	157	255	366	486	653	739	808
		-of which MSM	7	6	23	81	148	210	283	413	467	528
		Female	34	26	30	49	66	89	122	124	157	142
Total	58	52	91	206	321	455	608	777	896	950		
B1,B2	Uncomplicated gonorrhoea	Male (all)	10,669	14,272	15,654	16,940	16,268	14,514	12,694	12,267	11,724	10,077
		-of which MSM	1,812	2,867	3,454	3,298	3,653	3,619	3,927	4,074	3,354	2,962
		Female	4,880	6,222	6,744	7,417	7,224	6,265	5,025	5,007	5,423	5,100
		Total	15,549	20,494	22,398	24,357	23,492	20,779	17,719	17,274	17,147	15,177
B3	Gonococcal ophthalmia neonatorum	Male (all)	5	7	7	8	7	10	9	20	12	14
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	7	6	9	9	15	14	81	8	16	22
Total	12	13	16	17	22	24	90	28	28	36		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	1,557	2,112	2,588	3,012	3,301	3,449	3,518	3,731	4,203	4,380
		-of which MSM	429	597	815	879	1,080	1,103	1,235	1,337	1,328	1,329
		Female	1,789	2,189	2,563	2,944	3,134	3,064	3,078	2,918	3,227	3,437
Total	3,346	4,301	5,151	5,956	6,435	6,513	6,596	6,649	7,430	7,817		
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	74	79	95	106	99	135	103	88	97	96
		-of which MSM	12	8	14	25	10	19	31	18	18	19
		Female	237	260	273	272	315	262	225	235	255	238
Total	311	339	368	378	414	397	328	323	352	334		

England: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	28	30	29	34	60	86	258	212	255	276
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	23	31	33	37	76	63	59	66	131	91
		Total	51	61	62	71	136	149	317	278	386	367
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	21,776	26,579	29,562	34,331	37,996	42,681	45,220	49,018	53,469	54,194
		-of which MSM	621	962	1,365	1,422	1,725	1,789	2,261	2,808	3,117	3,726
		Female	29,184	34,791	38,618	43,786	47,520	50,267	50,798	50,192	53,496	53,671
		Total	50,960	61,370	68,180	78,117	85,516	92,948	96,018	99,210	106,965	107,865
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	432	528	545	571	637	653	723	780	798	921
		-of which MSM	25	23	35	32	52	45	70	69	66	85
		Female	1,763	2,114	2,074	2,134	2,091	2,311	2,238	2,176	2,523	2,608
		Total	2,195	2,642	2,619	2,705	2,728	2,964	2,961	2,956	3,321	3,529
C4D	Chlamydial ophthalmia neonatorum	Male (all)	15	19	15	15	18	42	15	13	13	18
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	12	23	28	25	25	34	22	19	8	16
		Total	27	42	43	40	43	76	37	32	21	34
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	11,291	14,023	17,345	21,502	22,531	24,371	25,992	25,200	28,618	29,437
		-of which MSM	243	405	620	1,032	1,282	1,372	1,627	1,917	2,161	2,339
		Female	8,243	10,118	12,096	14,207	14,651	16,668	18,505	18,616	20,444	21,568
		Total	19,534	24,141	29,441	35,709	37,182	41,039	44,497	43,816	49,062	51,005
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	53,214	56,007	59,467	62,040	67,292	69,870	70,394	69,723	67,398	63,391
		-of which MSM	3,375	3,847	4,566	4,470	5,000	5,051	5,195	5,730	5,176	4,910
		Female	0	0	0	0	6,636	7,417	7,389	6,731	6,372	6,600
		Total	53,214	56,007	59,467	62,040	73,928	77,287	77,783	76,454	73,770	69,991
C4I	Epidemiological treatment of NSGI	Male (all)	6,053	6,141	5,907	5,899	5,958	6,213	5,751	5,623	5,931	6,070
		-of which MSM	720	608	674	605	734	791	738	819	720	746
		Female	18,504	18,773	19,558	18,534	16,915	17,097	16,925	15,983	16,015	14,723
		Total	24,557	24,914	25,465	24,433	22,873	23,310	22,676	21,606	21,946	20,793
C5	Complicated infection (non-chlamydial/non-gonococcal) including PID and epididymitis	Male (all)	3,008	3,498	3,713	3,874	3,956	4,364	5,025	4,728	5,183	5,832
		-of which MSM	206	198	237	216	239	275	340	327	347	478
		Female	10,228	10,799	10,943	11,324	10,752	12,512	11,966	12,392	14,159	14,918
		Total	13,236	14,297	14,656	15,198	14,708	16,876	16,991	17,120	19,342	20,750
C6A	Trichomoniasis	Male (all)	641	731	733	568	285	297	330	313	384	363
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	5,448	6,036	6,279	6,628	6,224	5,924	5,074	5,179	5,332	5,247
		Total	6,089	6,767	7,012	7,196	6,509	6,221	5,404	5,492	5,716	5,610
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	1,388	1,314	1,392	1,391	1,444	1,420	1,472	1,637	1,664	1,965
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	55,518	60,667	64,884	70,909	71,917	69,426	70,646	73,497	85,482	90,485
		Total	56,906	61,981	66,276	72,300	73,361	70,846	72,118	75,134	87,146	92,450
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	11,996	12,049	11,972	11,660	11,827	11,644	11,075	10,896	12,666	13,554
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	7,643	7,079	6,753	5,680	5,382	5,816	4,852	4,312	4,603	5,113
		Total	19,639	19,128	18,725	17,340	17,209	17,460	15,927	15,208	17,269	18,667

England: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
C7A	Anogenital candidosis	Male (all)	7,219	6,958	7,165	6,828	5,970	5,947	5,593	6,377	7,610	8,087	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	64,196	65,686	69,918	70,401	66,661	62,919	59,338	58,237	65,186	69,642	
	Total	71,415	72,644	77,083	77,229	72,631	68,866	64,931	64,614	72,796	77,729		
C7B	Epidemiological treatment of C6 & C7	Male (all)	3,621	3,351	3,729	4,177	4,263	4,197	4,374	4,158	4,668	5,032	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	6,395	5,773	6,676	6,674	7,062	7,920	7,306	7,232	7,980	8,988	
	Total	10,016	9,124	10,405	10,851	11,325	12,117	11,680	11,390	12,648	14,020		
C8,C9	Scabies/pediculosis pubis	Male (all)	4,491	4,611	4,178	3,592	3,058	2,420	1,885	1,861	1,767	1,728	
		-of which MSM	741	826	805	708	597	501	449	470	409	443	
		Female	1,320	1,150	948	707	504	393	296	257	267	234	
	Total	5,811	5,761	5,126	4,299	3,562	2,813	2,181	2,118	2,034	1,962		
C10A	Anogenital herpes simplex – first attack	Male (all)	6,031	6,175	6,489	6,522	6,449	6,438	6,810	7,541	9,123	10,187	
		-of which MSM	338	384	399	479	468	458	550	526	616	750	
		Female	9,849	9,972	10,565	10,988	10,708	10,514	10,826	11,985	14,620	16,057	
	Total	15,880	16,147	17,054	17,510	17,157	16,952	17,636	19,526	23,743	26,244		
C10B	Anogenital herpes simplex – recurrence	Male (all)	5,983	6,095	6,204	6,376	6,373	6,072	6,248	6,624	7,185	8,035	
		-of which MSM	396	442	397	391	462	433	498	514	555	654	
		Female	7,226	7,345	8,049	8,142	8,009	7,724	8,014	8,378	9,705	10,885	
	Total	13,209	13,440	14,253	14,518	14,382	13,796	14,262	15,002	16,890	18,920		
C11A	Anogenital warts – first attack	Male (all)	31,846	31,975	32,700	33,987	34,642	36,418	36,410	37,490	40,202	41,610	
		-of which MSM	1,565	1,673	1,771	1,963	2,108	2,032	2,258	2,318	2,376	2,438	
		Female	29,311	28,686	29,723	29,951	30,637	31,799	32,403	33,234	36,119	37,568	
	Total	61,157	60,661	62,423	63,938	65,279	68,217	68,813	70,724	76,321	79,178		
C11B	Anogenital warts – recurrence	Male (all)	22,141	22,226	22,579	22,061	21,813	22,228	23,312	24,639	27,156	28,274	
		-of which MSM	1,211	1,213	1,376	1,387	1,548	1,615	1,546	1,731	1,952	1,855	
		Female	12,293	12,419	12,170	12,137	12,501	12,792	13,714	14,985	16,871	17,812	
	Total	34,434	34,645	34,749	34,198	34,314	35,020	37,026	39,624	44,027	46,086		
C11C	Anogenital warts – re-registered cases	Male (all)	9,314	9,047	8,988	8,464	8,803	8,499	8,768	9,078	9,072	9,528	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	7,597	7,219	7,148	6,517	6,616	6,450	6,806	6,730	6,652	6,674	
	Total	16,911	16,266	16,136	14,981	15,419	14,949	15,574	15,808	15,724	16,202		
C12	Molluscum contagiosum	Male (all)	4,763	5,100	5,571	6,187	6,940	8,045	8,853	9,620	10,519	11,048	
		-of which MSM	215	206	225	209	250	258	286	311	330	321	
		Female	2,637	2,691	3,003	3,250	3,485	3,892	4,239	4,506	4,815	4,850	
	Total	7,400	7,791	8,574	9,437	10,425	11,937	13,092	14,126	15,334	15,898		
C13	Antigen positive viral hepatitis B	Male (all)	527	635	704	931	0	0	0	0	0	0	
		-of which MSM	102	128	155	195	0	0	0	0	0	0	
		Female	249	230	352	472	0	0	0	0	0	0	
	Total	776	865	1,056	1,403	0	0	0	0	0	0		
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	-	-	602	649	637	622	675	729	
		-of which MSM	-	-	-	-	111	88	90	60	57	67	
		Female	-	-	-	-	271	325	325	341	353	380	
	Total	-	-	-	-	873	974	962	963	1,028	1,109		

England: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	-	-	46	54	44	40	43	59
		-of which MSM	-	-	-	-	14	19	14	7	5	17
		Female	-	-	-	-	10	12	25	26	20	12
		Total	-	-	-	-	56	66	69	66	63	71
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	-	-	342	459	414	430	488	557
		-of which MSM	-	-	-	-	121	205	206	228	146	164
		Female	-	-	-	-	150	168	143	192	192	236
		Total	-	-	-	-	492	627	557	622	680	793
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	918	929	918	1,044	1,004	1,039	941	860	805	858
		-of which MSM	83	103	104	140	110	123	115	117	80	120
		Female	425	422	380	484	462	438	398	355	290	337
		Total	1,343	1,351	1,298	1,528	1,466	1,477	1,339	1,215	1,095	1,195
E1A	New HIV diagnosis - asymptomatic	Male (all)	1,063	1,184	1,655	2,302	2,656	2,769	2,850	2,653	3,321	2,965
		-of which MSM	518	590	740	737	1,094	1,201	1,294	1,221	1,356	1,189
		Female	573	670	1,109	1,977	2,106	2,054	2,025	1,684	2,178	1,619
		Total	1,636	1,854	2,764	4,279	4,762	4,823	4,875	4,337	5,499	4,584
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	6,505	7,588	8,375	10,058	0	18	0	0	0	0
		-of which MSM	0	0	0	0	0	0	0	0	0	0
		Female	2,096	2,860	3,792	5,532	0	16	0	0	0	0
		Total	8,601	10,448	12,167	15,590	0	34	0	0	0	0
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	-	-	25,773	30,935	37,629	39,166	39,266	41,560
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	-	-	12,209	16,402	20,525	21,561	22,965	23,954
		Total	-	-	-	-	37,982	47,337	58,154	60,727	62,231	65,514
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	514	470	641	744	664	669	735	616	473	965
		-of which MSM	278	222	300	244	263	246	253	224	141	294
		Female	195	198	326	445	441	385	443	354	304	381
		Total	709	668	967	1,189	1,105	1,054	1,178	970	777	1,346
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	7,788	8,880	9,139	9,265	0	3	0	0	0	0
		-of which MSM	0	0	0	0	0	0	0	0	0	0
		Female	1,854	2,230	2,623	3,310	0	2	0	0	0	0
		Total	9,642	11,110	11,762	12,575	0	5	0	0	0	0
E3A*	AIDS – first presentation	Male (all)	377	360	430	436	0	0	0	0	0	0
		-of which MSM	151	178	167	155	0	0	0	0	0	0
		Female	110	139	214	227	0	0	0	0	0	0
		Total	487	499	644	663	0	0	0	0	0	0
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	-	-	176	217	227	189	239	224
		-of which MSM	-	-	-	-	69	61	76	64	74	68
		Female	-	-	-	-	137	116	150	88	96	117
		Total	-	-	-	-	313	333	377	277	335	341
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	-	-	236	153	161	157	110	108
		-of which MSM	-	-	-	-	94	74	60	82	48	56
		Female	-	-	-	-	133	63	70	69	50	54
		Total	-	-	-	-	369	216	231	226	160	162

England: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
E3B	AIDS – subsequent presentation	Male (all)	5,396	6,119	6,683	7,605	9,539	10,273	11,636	11,563	11,115	11,447	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	1,229	1,551	2,001	2,584	3,999	4,319	4,967	5,157	5,343	5,519	
	Total	6,625	7,670	8,684	10,189	13,538	14,592	16,603	16,720	16,458	16,966		
D2A	Urinary tract infection	Male (all)	2,040	1,910	1,981	2,522	2,483	2,479	2,029	1,990	2,208	2,530	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	9,064	9,351	9,431	9,152	8,717	9,772	9,346	9,692	12,087	14,193	
	Total	11,104	11,261	11,412	11,674	11,200	12,251	11,375	11,682	14,295	16,723		
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	52,418	54,692	54,839	57,505	64,250	64,697	66,968	68,400	78,547	89,066	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	48,791	53,560	54,402	54,721	53,126	54,501	55,660	56,843	64,010	72,046	
	Total	101,209	108,252	109,241	112,226	117,376	119,198	122,628	125,243	142,557	161,112		
D3	Other episodes not requiring treatment	Male (all)	100,901	106,796	125,321	141,931	133,675	160,686	188,431	208,127	235,346	260,602	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	110,295	113,953	131,498	146,874	135,640	158,352	190,100	207,497	243,585	268,608	
	Total	211,196	220,749	256,819	288,805	269,315	319,038	378,531	415,624	478,931	529,210		
P1A	HIV antibody test (no sexual health screen)	Male (all)	73,635	83,890	109,903	145,623	48,164	42,841	39,892	33,782	31,248	30,721	
		-of which MSM	10,869	12,125	15,980	18,223	8,154	7,455	7,747	7,281	6,743	6,734	
		Female	59,051	67,569	92,977	129,933	37,024	30,537	29,860	25,224	24,208	23,960	
	Total	132,686	151,459	202,880	275,556	85,188	73,378	69,752	59,006	55,456	54,681		
P1B	HIV antibody test offered and refused	Male (all)	12,962	14,326	19,265	42,360	95,065	116,163	112,959	107,708	111,805	116,444	
		-of which MSM	1,396	1,137	1,643	3,102	5,937	6,237	6,280	6,288	6,044	6,157	
		Female	12,432	13,612	19,130	47,485	109,531	133,388	131,486	127,902	141,017	152,883	
	Total	25,394	27,938	38,395	89,845	204,596	249,551	244,445	235,610	252,822	269,327		
P2	Hepatitis B vaccination	Male (all)	9,989	10,885	11,859	13,313	14,464	15,858	15,512	15,983	16,741	18,216	
		-of which MSM	4,707	5,218	5,850	6,604	7,838	8,348	8,433	8,354	8,255	8,997	
		Female	2,491	2,630	3,142	3,909	4,092	4,982	5,252	5,558	6,113	6,954	
	Total	12,480	13,515	15,001	17,222	18,556	20,840	20,764	21,541	22,854	25,170		
P3	Contraception (excluding condom provision)	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	30,263	32,771	33,646	31,444	32,774	33,858	34,563	37,987	45,436	52,121	
	Total	30,263	32,771	33,646	31,444	32,774	33,858	34,563	37,987	45,436	52,121		
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	7,583	6,214	4,744	3,663	5,212	3,351	1,675	1,362	1,359	1,552	
	Total	7,583	6,214	4,744	3,663	5,212	3,351	1,675	1,362	1,359	1,552		

England: Number of new STI episodes by sex, 1999 - 2008 continued

			1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1,184	1,120	1,024	751	912	793	383	292	256	333
		Total	1,184	1,120	1,024	751	912	793	383	292	256	333
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	-	-	141,976	142,890	136,146	131,824	135,365	133,594
		-of which MSM	-	-	-	-	9,684	9,833	10,736	11,977	11,637	11,919
		Female	-	-	-	-	176,335	174,285	168,263	161,169	172,261	175,952
		Total	-	-	-	-	318,311	317,175	304,409	292,993	307,626	309,546
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	-	-	158,950	208,016	255,141	298,546	356,410	400,092
		-of which MSM	-	-	-	-	15,882	18,964	22,830	27,686	30,502	36,339
		Female	-	-	-	-	156,028	201,819	251,417	292,160	359,851	408,955
		Total	-	-	-	-	314,978	409,835	506,558	590,706	716,261	809,047
Total new STI diagnoses												
	Male (all)	138,828	151,612	161,918	173,187	182,958	191,870	195,329	199,910	207,867	206,628	
	-of which MSM	9,788	11,973	14,376	14,593	16,481	16,729	18,645	19,804	19,081	19,307	
	Female	95,778	103,788	110,836	119,174	129,189	134,637	133,682	134,538	146,528	149,701	
	Total	234,606	255,400	272,754	292,361	312,147	326,507	329,011	334,448	354,395	356,329	
Total other STI diagnoses												
	Male (all)	77,126	83,435	89,173	95,731	105,628	113,752	124,618	127,565	134,489	140,703	
	-of which MSM	3,250	3,536	4,194	4,660	5,505	5,755	6,166	7,054	7,451	7,772	
	Female	61,412	65,376	70,925	74,977	79,082	85,605	93,691	95,274	102,268	105,549	
	Total	138,538	148,811	160,098	170,708	184,710	199,357	218,309	222,839	236,757	246,252	
Total other GUM clinic diagnoses												
	Male (all)	80,127	81,838	82,700	86,058	92,231	92,585	93,547	95,410	109,374	122,437	
	-of which MSM	185	231	259	335	356	435	425	412	288	368	
	Female	201,048	210,102	218,564	222,907	219,882	215,441	210,097	212,381	241,818	263,317	
	Total	281,175	291,940	301,264	308,965	312,113	308,026	303,644	307,791	351,192	385,754	
Total services provided												
	Male (all)	197,487	215,897	266,348	343,227	592,294	686,454	748,081	795,970	886,915	959,669	
	-of which MSM	16,972	18,480	23,473	27,929	47,495	50,837	56,026	61,586	63,181	70,146	
	Female	214,532	230,535	280,393	359,645	651,424	737,221	810,941	857,497	992,471	1,089,433	
	Total	412,019	446,432	546,741	702,872	1,243,718	1,423,675	1,559,022	1,653,467	1,879,386	2,049,102	

Source: HPA Centre for Infections

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11a or C11b treated for more than 3 months

- data not available/applicable

**Supplementary Tables of Sexually
Transmitted Infections in the United Kingdom**



Code Groups
New STI Diagnoses
Chlamydial infection (uncomplicated and complicated)
Gonorrhoea (uncomplicated and complicated)
Infectious and early latent syphilis
Genital Herpes simplex (first attack)
Genital warts (first attack)
New HIV diagnosis
Non-specific genital infection (uncomplicated and complicated)
Chancroid/lymphogranuloma venerum (LGV)/Donovanosis
Molluscum contagiosum
Trichomoniasis
Scabies
Pediculus pubis
Other STI Diagnoses
Congenital and other acquired syphilis
Recurrent genital Herpes simplex
Recurrent and re-registered genital warts
Subsequent HIV presentations (including AIDS)
Ophthalmia neonatorum (chlamydial or gonococcal)
Epidemiological treatment of suspected STIs (syphilis, chlamydia, gonorrhoea, non-specific genital infection)
Other diagnoses made at GUM clinics
Viral hepatitis B
Viral hepatitis C
Vaginosis and balanitis (including epidemiological treatment)
Anogenital candidiasis (including epidemiological treatment)
Urinary tract infection
Cervical abnormalities
Other conditions requiring treatment at a GUM clinic
Services provided
HIV antibody test
Sexual health screen
Hepatitis B vaccination
Contraception (excluding condom provision)
Other episode not requiring treatment