



Numbers of persons accessing HIV-related care

**Survey of Prevalent HIV Infections Diagnosed
(SOPHID)**

London SHA tables

Table Index

London SHA tables

<u>Table ExpLon:</u>	Diagnosed HIV-infected individuals in London by exposure category: 1999-2008
<u>Table EthnLon:</u>	Diagnosed HIV-infected individuals in London by ethnic group: 1999-2008
<u>Table AgeLon:</u>	Diagnosed HIV-infected individuals in London by age-group and sex: 1999-2008
<u>Table ARTLon:</u>	Diagnosed HIV-infected individuals in London by level of antiretroviral therapy and CD4 count: 2000-2008

SOPHID Table ExpLon: Diagnosed HIV-infected individuals in London by exposure category.¹

Exposure category ²	Year of survey									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Sex between men³	6,403	6,820	7,653	8,288	8,944	9,662	10,426	11,169	11,771	12,317
Injecting drug use	492	466	455	454	497	497	486	522	507	505
Sex between men & women	3,775	4,364	5,838	7,203	8,291	9,265	10,353	11,046	11,639	12,314
Blood/blood products recipient	139	123	128	128	119	124	125	129	114	105
Mother-to-child transmission	385	366	406	472	509	561	627	597	619	678
Other/not known	178	120	176	197	219	162	167	342	517	987
Total	11,372	12,259	14,656	16,742	18,579	20,271	22,184	23,805	25,167	26,906

¹ Figures may be different to those released previously due to improvements in the methods used to identify patients seen more than once within the survey year. This method has been applied to all survey years.

² Patients with missing fields (e.g. ethnicity/age-group) may have been assigned values based on subsequent years data.

³ Including those who also injected drugs.

SOPHID Table EthnLon: Diagnosed HIV-infected individuals in London by ethnic group.¹

Ethnic group ²	Year of survey									
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
White	6,605	6,870	7,610	8,202	9,137	9,801	10,458	11,194	11,781	12,244
Black-Caribbean	332	355	542	623	759	843	935	1,031	1,138	1,230
Black-African	3,191	3,617	4,885	6,020	6,891	7,580	8,305	8,813	9,115	9,620
Black-other/Black-unspecified	230	250	324	366	365	429	554	592	670	778
Indian/Pakistani/Bangladeshi	167	170	170	197	217	258	309	350	397	456
Other/mixed	764	773	851	975	772	871	1,135	1,227	1,339	1,637
Other Asian/Oriental³	9	123	153	195	262	326	304	354	400	447
Not known	74	101	121	164	176	163	184	244	327	494
Total	11,372	12,259	14,656	16,742	18,579	20,271	22,184	23,805	25,167	26,906

¹ Figures may be different to those released previously due to improvements in the methods used to identify patients seen more than once within the survey year. This method has been applied to all survey years.

² Patients with missing fields (e.g. ethnicity/age-group) may have been assigned values based on subsequent years data.

³ This category was not recorded before 2000.

SOPHID Table AgeLon: Diagnosed HIV-infection in London by age-group and sex.¹

Age-group (yrs) ^{2,3}	Year of survey										
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
<1	<i>Boys</i>	19	9	7	4	1	3	12	5	3	2
	<i>Girls</i>	13	5	8	3	7	7	19	4	2	3
	Sub total	32	14	15	7	8	10	31	9	5	5
1-14	<i>Boys</i>	191	187	205	225	232	248	273	238	226	212
	<i>Girls</i>	190	157	176	204	225	234	239	222	223	200
	Sub total	381	344	381	429	457	482	513	460	449	412
15-24	<i>Men</i>	181	174	220	263	280	297	325	391	422	487
	<i>Women</i>	163	175	257	343	373	442	463	486	508	523
	Sub total	344	349	477	606	653	739	788	877	930	1,010
25-29	<i>Men</i>	870	833	834	836	867	891	948	995	1,093	1,169
	<i>Women</i>	477	511	635	784	805	813	897	911	860	804
	Sub total	1,347	1,344	1,469	1,621	1,672	1,704	1,845	1,906	1,953	1,973
30-34	<i>Men</i>	2,079	2,041	2,162	2,149	2,143	2,088	2,114	2,141	2,025	2,058
	<i>Women</i>	814	853	1,102	1,244	1,364	1,390	1,458	1,544	1,520	1,512
	Sub total	2,893	2,894	3,264	3,393	3,507	3,478	3,572	3,685	3,545	3,570
35-39	<i>Men</i>	2,249	2,412	2,880	3,219	3,452	3,494	3,548	3,473	3,448	3,339
	<i>Women</i>	672	809	1,030	1,255	1,434	1,533	1,684	1,738	1,823	1,900
	Sub total	2,921	3,221	3,910	4,474	4,886	5,027	5,232	5,211	5,271	5,239
40-44	<i>Men</i>	1,357	1,584	1,966	2,288	2,689	3,174	3,496	3,896	4,203	4,367
	<i>Women</i>	233	336	483	648	870	1,082	1,275	1,457	1,527	1,743
	Sub total	1,591	1,920	2,449	2,936	3,559	4,256	4,771	5,353	5,730	6,110
45-49	<i>Men</i>	767	870	1,053	1,252	1,430	1,735	2,102	2,424	2,740	3,181
	<i>Women</i>	102	147	215	283	354	442	579	690	870	1,068
	Sub total	869	1,017	1,268	1,535	1,784	2,177	2,681	3,114	3,610	4,249
50-54	<i>Men</i>	463	515	615	738	809	932	1,028	1,200	1,386	1,640
	<i>Women</i>	36	55	87	126	173	214	263	331	370	448
	Sub total	500	570	702	864	982	1,146	1,291	1,531	1,756	2,088
55-59	<i>Men</i>	244	296	338	394	484	553	605	680	770	866
	<i>Women</i>	21	30	49	58	76	113	120	144	170	220
	Sub total	265	326	387	452	560	666	725	824	940	1,086
60+	<i>Men</i>	191	223	279	354	435	491	607	682	797	948
	<i>Women</i>	27	37	54	65	76	95	128	153	181	216
	Sub total	218	260	333	419	511	586	735	835	978	1,164
Age not reported	11	0	1	6	0	0	0	0	0	0	
Total	11,372	12,259	14,656	16,742	18,579	20,271	22,184	23,805	25,167	26,906	

¹ Figures may be different to those released previously due to improvements in the methods used to identify patients seen more than once within the survey year. This method has been applied to all survey years.

² Patients with missing fields (e.g. ethnicity/age-group) may have been assigned values based on subsequent years data.

³ Totals may include those with sex not reported.

SOPHID Table ARTLon: Diagnosed HIV-infected individuals in London by level of antiretroviral therapy and CD4 count.¹

Level of antiretroviral therapy and CD4 count/mm ³	Year of survey								
	2000	2001	2002	2003	2004	2005	2006	2007	2008
No antiretroviral therapy									
0-100	148	125	147	211	182	175	158	139	101
101-200	195	197	217	239	245	274	218	218	178
201-350	545	810	892	1,142	1,418	1,565	1,463	1,425	1,011
351-500	586	948	926	1,260	1,544	1,694	1,633	1,891	1,874
>500	825	1,200	1,166	1,622	1,762	1,961	1,984	2,261	2,301
Not reported	1,178	1,230	803	931	687	741	519	690	785
Sub total	3,477	4,510	4,151	5,405	5,838	6,410	5,975	6,624	6,250
Mono									
0-100	8	9	6	9	10	10	15	16	11
101-200	5	22	10	10	11	14	21	15	15
201-350	12	45	16	39	30	48	73	34	48
351-500	15	55	29	37	32	59	75	38	43
>500	30	83	52	65	58	69	160	71	103
Not reported	18	40	64	88	14	8	10	7	21
Sub total	88	254	177	248	155	208	354	181	241
Dual									
0-100	20	13	15	12	10	23	24	21	14
101-200	28	28	20	26	21	54	51	18	45
201-350	77	69	54	67	46	99	116	90	101
351-500	48	66	52	40	55	96	148	83	130
>500	56	98	62	53	72	147	204	134	289
Not reported	81	57	34	38	11	7	18	26	56
Sub total	310	331	237	236	215	426	561	372	635
Triple									
0-100	384	452	507	524	526	467	366	387	365
101-200	682	894	1,003	1,021	1,120	999	932	910	863
201-350	1,206	1,759	2,136	2,344	2,884	2,946	2,876	2,990	3,102
351-500	1,004	1,508	1,816	2,140	2,740	2,948	3,092	3,417	4,020
>500	1,156	2,001	2,543	2,867	3,714	3,970	4,380	5,289	6,476
Not reported	2,110	1,492	948	1,163	313	311	525	879	1,065
Sub total	6,542	8,106	8,953	10,059	11,297	11,641	12,171	13,872	15,891
Quadruple +									
0-100	101	103	143	139	164	195	214	185	121
101-200	129	150	242	278	310	386	386	367	285
201-350	214	256	516	547	655	909	910	906	763
351-500	155	260	411	463	589	774	973	997	882
>500	194	353	575	615	761	999	1,309	1,442	1,462
Not reported	126	50	193	259	49	88	95	171	331
Sub total	919	1,172	2,080	2,301	2,528	3,351	3,887	4,068	3,844
ART - level not reported									
0-100	0	0	1	1	0	0	1	0	5
101-200	1	1	0	1	1	0	2	0	1
201-350	0	0	0	2	2	1	3	0	2
351-500	0	0	0	2	0	1	1	0	3
>500	0	0	0	2	0	0	1	0	12
Not reported	3	2	5	1	2	0	0	0	3
Sub total	4	3	6	9	5	2	8	0	26
Not reported									
0-100	48	11	42	7	15	4	11	13	1
101-200	48	10	60	4	18	3	33	1	2
201-350	117	34	253	17	36	19	232	0	1
351-500	111	30	258	17	33	27	202	2	1
>500	144	119	356	37	31	30	258	1	3
Not reported	451	76	169	239	100	63	113	33	11
Sub total	919	280	1,138	321	233	146	849	50	19
Total	12,259	14,656	16,742	18,579	20,271	22,184	23,805	25,167	26,906

¹ Figures may be different to those released previously due to improvements in the methods used to identify patients seen more than once within the survey year. This method has been applied to all survey years.