
Table 4: Number of selected STI diagnoses made at genitourinary medicine clinics in England: 2001 - 2010



Data are presented by country, SHA, gender, sexual orientation & age group

Notes on using these tables:

Numbers of diagnoses are adjusted for missing clinic data
Data on unknown age-groups are included in the 'Total'
Data on unknown gender are included in the 'Total'
Data are presented for England only.

An improved STI surveillance system (GUMCAD) was introduced in England in 2009. Unlike the KC60 return, duplicate records in GUMCAD can be identified and removed resulting in a slight reduction in the number of diagnoses reported. England data from 2000-2008 have therefore been statistically adjusted to enable fair comparisons of STI diagnoses over time.

Definitions of conditions presented:

Chlamydia	uncomplicated genital chlamydial infection, KC60 code C4A & C4C
Gonorrhoea	uncomplicated gonorrhoea, KC60 code B1& B2
Syphilis	primary and secondary infectious syphilis, KC60 code A1& A2
Herpes	anogenital herpes (first episode), KC60 code C10A
Warts	anogenital warts (first episode), KC60 code C11A

Definitions of data totals:

Total new STI diagnoses:	} Refer to 'Code Groups' for conditions included
Total other STI diagnoses:	
Total other GUM clinic diagnoses:	
Total services provided:	

Data source

England: GUM clinic KC60 returns (2001-2008)
GUM clinic GUMCAD returns (2009-2010)

Data supplied by

Health Protection Agency, Health Protection Services Colindale

Data prepared by:

HIV and Sexually Transmitted Infections Department
Health Protection Agency
Health Protection Services Colindale
61 Colindale Avenue
London
NW9 5EQ

Tel. 020-8200-6868

Email: hiv/sti@hpa.org.uk

Website: www.hpa.org.uk

Table 4: Number of selected STI diagnoses made at genitourinary medicine clinics in England: 2001 - 2010

Data are presented by country, SHA, gender, sexual orientation & age group



Country tables

England 4 a

Condition tables Diagnoses by SHA Diagnoses by age group (England)

Infectious syphilis	4 b	(i)
Gonorrhoea	4 c	(i)
Chlamydia	4 d	(i)
Genital herpes	4 e	(i)
Genital warts	4 f	(i)

Code Groups: Definitions of conditions presented

Further Information

England

Data on selected diagnoses, 1971 to 2010, are available upon request

HPA Health Protection Services Colindale

Contact: Geraldine Leong

Tel: +44 (0)20 8200 6868

Email: geraldine.leong@hpa.org.uk

Web: www.hpa.org.uk

Table 4 a: Summary tables of selected conditions by gender for England between 2001 - 2010

		2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% change	
												2009-2010	2001-2010
Syphilis (primary and secondary)	Total Males	586	997	1,278	1,676	2,041	2,040	2,006	1,774	1,851	1,671	-10%	185%
	- of which MSM	329	574	710	847	1,195	1,163	1,173	989	1,229	1,070	-13%	225%
	Total Females	93	128	171	251	329	263	267	224	188	172	-9%	85%
	Total	679	1,125	1,449	1,927	2,370	2,303	2,273	1,998	2,048	1,858	-9%	174%
Gonorrhoea (uncomplicated)	Total Males	15,309	16,570	15,930	14,190	12,429	12,006	11,493	9,729	10,572	11,289	7%	-26%
	- of which MSM	3,333	3,184	3,527	3,491	3,788	3,928	3,228	2,597	3,379	4,503	33%	35%
	Total Females	6,550	7,205	7,032	6,108	4,899	4,885	5,299	4,940	5,004	4,825	-4%	-26%
	Total	21,859	23,775	22,962	20,298	17,328	16,891	16,792	14,669	15,606	16,145	3%	-26%
Chlamydia (uncomplicated)	Total Males	29,071	33,774	37,397	42,049	44,604	48,301	52,791	53,697	49,276	47,127	-4%	62%
	- of which MSM	1,265	1,328	1,616	1,676	2,116	2,626	2,918	3,569	4,084	4,975	22%	293%
	Total Females	37,810	42,882	46,557	49,297	49,816	49,209	52,496	53,426	47,475	43,843	-8%	16%
	Total	66,881	76,656	83,954	91,346	94,420	97,510	105,287	107,123	96,869	91,075	-6%	36%
Herpes (first episode)	Total Males	6,411	6,445	6,369	6,356	6,730	7,456	9,055	10,104	10,824	11,584	7%	81%
	- of which MSM	391	468	457	448	538	515	600	735	818	973	19%	149%
	Total Females	10,395	10,814	10,534	10,338	10,649	11,798	14,432	15,990	16,703	18,090	8%	74%
	Total	16,806	17,259	16,903	16,694	17,379	19,254	23,487	26,094	27,564	29,703	8%	77%
Warts (first episode)	Total Males	32,285	33,542	34,198	35,972	35,975	37,021	39,723	41,094	41,934	40,953	-2%	27%
	- of which MSM	1,744	1,931	2,074	2,003	2,225	2,280	2,339	2,488	2,557	2,528	-1%	45%
	Total Females	29,220	29,440	30,121	31,279	31,877	32,679	35,549	37,062	35,904	34,582	-4%	18%
	Total	61,505	62,982	64,319	67,251	67,852	69,700	75,272	78,156	77,900	75,615	-3%	23%
Total new STI diagnoses	Total Males	158,494	169,482	179,048	187,907	191,335	195,667	203,825	201,614	194,771	187,332	-4%	18%
	- of which MSM	13,863	14,082	15,876	16,118	17,953	19,041	18,308	18,546	20,726	22,428	8%	62%
	Total Females	108,297	116,388	126,249	131,695	130,775	131,596	143,516	147,704	139,979	135,169	-3%	25%
	Total	266,791	285,870	305,297	319,602	322,110	327,263	347,341	349,318	335,139	322,920	-4%	21%
Total other STI diagnoses	Total Males	84,235	90,520	93,024	99,654	108,673	111,783	118,548	125,040	126,117	128,701	2%	53%
	- of which MSM	4,027	4,488	5,297	5,530	5,962	6,826	7,185	7,376	8,260	9,047	10%	125%
	Total Females	68,069	71,943	72,356	77,717	84,584	86,193	92,711	96,607	93,486	90,944	-3%	34%
	Total	152,304	162,463	165,380	177,371	193,257	197,976	211,259	221,647	219,906	220,049	0%	44%
Total other GUM diagnoses	Total Males	76,782	79,806	85,333	85,837	86,758	88,304	101,176	113,046	117,261	119,255	2%	55%
	- of which MSM	245	316	337	408	397	385	272	537	679	415	-39%	69%
	Total Females	210,428	214,686	211,768	207,625	202,640	204,352	233,180	255,507	266,569	268,246	1%	27%
	Total	287,210	294,492	297,101	293,462	289,398	292,656	334,356	368,553	384,559	388,220	1%	35%
Total services provided	Total Males	265,968	342,816	591,863	686,581	749,679	795,495	889,059	970,958	994,688	1,029,189	3%	287%
	- of which MSM	23,275	27,712	47,247	50,546	55,955	61,322	62,899	69,860	82,813	91,701	11%	294%
	Total Females	280,300	359,527	651,303	738,157	813,214	857,330	994,540	1,105,587	1,137,272	1,184,375	4%	323%
	Total	546,268	702,343	1,243,166	1,424,738	1,562,893	1,652,825	1,883,599	2,076,545	2,134,801	2,216,494	4%	306%

Data totals include 'Unknown' gender. As such, totals may exceed the sum of male and females

Table 4 b: New episodes of infectious syphilis (primary and secondary) by SHA & gender: England 2001 - 2010

MALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	15	21	55	80	139	122	87	93	116	114	7%	-2%	660%
East of England	16	27	64	70	69	71	59	54	49	50	3%	2%	213%
London	271	514	566	669	692	667	790	594	765	735	44%	-4%	171%
North East	2	17	63	88	74	72	104	82	86	58	3%	-33%	2800%
North West	168	202	192	258	383	353	301	310	252	249	15%	-1%	48%
South Central	11	20	60	67	103	101	69	78	64	67	4%	5%	509%
South East Coast	25	60	88	128	75	93	102	99	94	85	5%	-10%	240%
South West	10	32	53	80	95	131	140	133	93	69	4%	-26%	590%
West Midlands	40	51	85	168	308	293	249	181	164	132	8%	-20%	230%
Yorkshire & the Humber	28	53	52	68	103	137	105	150	168	112	7%	-33%	300%
England	586	997	1,278	1,676	2,041	2,040	2,006	1,774	1,851	1,671	100%	-10%	185%

MALES (MSM cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	5	8	15	28	48	44	39	42	66	66	6%	0%	1220%
East of England	9	12	28	23	28	30	24	17	26	27	3%	4%	200%
London	160	292	331	362	481	445	530	330	575	501	47%	-13%	213%
North East	1	15	45	65	56	39	66	49	60	41	4%	-32%	4000%
North West	111	149	131	130	283	248	209	223	193	191	18%	-1%	72%
South Central	2	3	18	17	38	54	25	34	31	40	4%	29%	1900%
South East Coast	12	42	69	100	50	69	49	75	45	39	4%	-13%	225%
South West	6	16	16	44	55	61	91	73	48	42	4%	-13%	600%
West Midlands	13	21	36	50	108	104	86	80	74	61	6%	-18%	369%
Yorkshire & the Humber	10	16	21	28	48	69	54	66	111	62	6%	-44%	520%
England	329	574	710	847	1,195	1,163	1,173	989	1,229	1,070	100%	-13%	225%

FEMALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	2	3	8	18	43	40	39	20	14	18	10%	29%	800%
East of England	3	9	14	18	13	9	10	17	8	8	5%	0%	167%
London	46	67	93	100	111	72	77	52	52	61	35%	17%	33%
North East	2	0	4	4	6	15	17	9	19	11	6%	-42%	450%
North West	10	14	13	11	25	23	18	21	26	16	9%	-38%	60%
South Central	5	6	9	11	15	4	5	7	7	5	3%	-29%	0%
South East Coast	2	10	7	5	14	8	8	12	7	5	3%	-29%	150%
South West	3	3	4	17	16	11	12	26	8	12	7%	50%	300%
West Midlands	14	5	11	52	69	71	65	47	28	16	9%	-43%	14%
Yorkshire & the Humber	6	11	8	15	17	10	16	13	19	20	12%	5%	233%
England	93	128	171	251	329	263	267	224	188	172	100%	-9%	85%

TOTAL ('Male', 'Female' and 'Not Known' gender)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	17	24	63	98	182	162	126	113	130	132	7%	2%	676%
East of England	19	36	78	88	82	80	69	71	57	58	3%	2%	205%
London	317	581	659	769	803	739	867	646	826	811	44%	-2%	156%
North East	4	17	67	92	80	87	121	91	105	69	4%	-34%	1625%
North West	178	216	205	269	408	376	319	331	278	265	14%	-5%	49%
South Central	16	26	69	78	118	105	74	85	71	72	4%	1%	350%
South East Coast	27	70	95	133	89	101	110	111	101	90	5%	-11%	233%
South West	13	35	57	97	111	142	152	159	101	81	4%	-20%	523%
West Midlands	54	56	96	220	377	364	314	228	192	148	8%	-23%	174%
Yorkshire & the Humber	34	64	60	83	120	147	121	163	187	132	7%	-29%	288%
England	679	1,125	1,449	1,927	2,370	2,303	2,273	1,998	2,048	1,858	100%	-9%	174%

[‡] Percentage of total cases in England during 2010 which were in this SHA

Table 4 b (i): New episodes of infectious syphilis (primary and secondary) by age group & gender: England 2001 - 2010

MALES (total cases)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	0	1	0	0	3	0	0	1	0	0	0%	-	-
15-19	12	26	27	33	67	56	44	48	48	39	2%	-19%	225%
20-24	67	93	122	164	266	236	229	213	221	188	11%	-15%	181%
25-34	224	360	472	552	660	608	599	502	552	514	31%	-7%	129%
35-44	179	347	435	616	656	732	650	590	578	514	31%	-11%	187%
45+	93	163	220	310	387	397	419	420	443	410	25%	-7%	341%
Total	586	997	1,278	1,676	2,041	2,040	2,006	1,774	1,851	1,671	100%	-10%	185%

FEMALES (total cases)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	0	1	0	0	7	1	2	3	0	0	0%	-	-
15-19	11	17	25	29	35	52	44	28	35	19	11%	-46%	73%
20-24	24	31	38	58	79	62	62	54	37	48	28%	30%	100%
25-34	37	49	67	90	98	78	86	80	66	53	31%	-20%	43%
35-44	15	19	26	52	70	51	38	36	28	37	22%	32%	147%
45+	5	10	15	22	40	18	30	22	20	15	9%	-25%	200%
Total	93	128	171	251	329	263	267	224	188	172	100%	-9%	85%

TOTAL ('Male', 'Female' and 'Not Known' gender)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	0	2	0	0	10	1	2	4	0	0	0%	-	-
15-19	23	43	52	62	102	108	88	76	83	58	3%	-30%	152%
20-24	91	124	160	222	345	298	291	267	259	237	13%	-8%	160%
25-34	261	409	539	642	758	686	685	582	621	573	31%	-8%	120%
35-44	194	366	461	668	726	783	688	626	609	556	30%	-9%	187%
45+	98	173	235	332	427	415	449	442	465	428	23%	-8%	337%
Total	679	1,125	1,449	1,927	2,370	2,303	2,273	1,998	2,048	1,858	100%	-9%	174%

† Percentage of total cases in England during 2010 which were in this SHA

Table 4 c: New episodes of gonorrhoea (uncomplicated) by SHA & gender: England 2001 - 2010

MALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	884	1,011	945	848	715	783	746	543	631	564	5%	-11%	-36%
East of England	640	650	810	783	644	557	536	512	605	579	5%	-4%	-10%
London	6,507	6,841	5,982	5,375	4,890	4,700	4,681	3,905	4,450	4,973	44%	12%	-24%
North East	371	525	448	423	415	348	320	327	390	367	3%	-6%	-1%
North West	1,959	2,013	2,069	1,757	1,615	1,715	1,640	1,389	1,225	1,427	13%	16%	-27%
South Central	635	681	833	671	547	487	430	438	452	479	4%	6%	-25%
South East Coast	683	706	635	700	619	616	601	417	486	526	5%	8%	-23%
South West	802	1,045	950	765	674	626	497	502	549	475	4%	-13%	-41%
West Midlands	1,821	1,689	1,748	1,451	1,139	1,101	1,093	940	972	1,080	10%	11%	-41%
Yorkshire & the Humber	1,007	1,409	1,510	1,417	1,171	1,073	949	756	812	819	7%	1%	-19%
England	15,309	16,570	15,930	14,190	12,429	12,006	11,493	9,729	10,572	11,289	100%	7%	-26%

MALES (MSM cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	104	84	126	124	108	124	86	54	88	148	3%	68%	42%
East of England	107	98	130	138	126	141	116	95	129	191	4%	48%	79%
London	1,777	1,564	1,701	1,850	2,097	2,098	1,864	1,275	1,947	2,445	54%	26%	38%
North East	74	134	107	107	146	110	74	74	97	94	2%	-3%	27%
North West	404	491	558	442	468	548	432	448	485	719	16%	48%	78%
South Central	104	84	94	92	109	130	91	85	119	135	3%	13%	30%
South East Coast	342	266	291	318	323	307	196	167	113	179	4%	58%	-48%
South West	143	154	157	134	111	139	90	144	117	179	4%	53%	25%
West Midlands	161	163	216	148	159	151	130	116	114	212	5%	86%	32%
Yorkshire & the Humber	117	146	147	138	141	180	149	139	170	201	4%	18%	72%
England	3,333	3,184	3,527	3,491	3,788	3,928	3,228	2,597	3,379	4,503	100%	33%	35%

FEMALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	477	528	492	401	338	381	476	391	443	419	9%	-5%	-12%
East of England	355	385	433	432	298	277	258	305	298	266	6%	-11%	-25%
London	2,597	2,885	2,503	1,987	1,518	1,489	1,767	1,666	1,580	1,630	34%	3%	-37%
North East	174	217	217	164	139	134	130	149	195	247	5%	27%	42%
North West	745	771	767	786	765	782	801	738	679	623	13%	-8%	-16%
South Central	307	340	398	305	205	195	218	180	193	239	5%	24%	-22%
South East Coast	153	171	190	193	149	181	154	146	174	141	3%	-19%	-8%
South West	307	382	436	320	287	298	262	243	253	179	4%	-29%	-42%
West Midlands	940	858	856	736	589	597	673	625	673	563	12%	-16%	-40%
Yorkshire & the Humber	495	668	740	784	611	551	560	497	516	518	11%	0%	5%
England	6,550	7,205	7,032	6,108	4,899	4,885	5,299	4,940	5,004	4,825	100%	-4%	-26%

TOTAL ('Male', 'Female' and 'Not Known' gender)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	1,361	1,539	1,437	1,249	1,053	1,164	1,222	934	1,082	985	6%	-9%	-28%
East of England	995	1,035	1,243	1,215	942	834	794	817	903	845	5%	-6%	-15%
London	9,104	9,726	8,485	7,362	6,408	6,189	6,448	5,571	6,052	6,632	41%	10%	-27%
North East	545	742	665	587	554	482	450	476	585	614	4%	5%	13%
North West	2,704	2,784	2,836	2,543	2,380	2,497	2,441	2,127	1,904	2,050	13%	8%	-24%
South Central	942	1,021	1,231	976	752	682	648	618	645	718	4%	11%	-24%
South East Coast	836	877	825	893	768	797	755	563	660	667	4%	1%	-20%
South West	1,109	1,427	1,386	1,085	961	924	759	745	802	654	4%	-18%	-41%
West Midlands	2,761	2,547	2,604	2,187	1,728	1,698	1,766	1,565	1,645	1,643	10%	0%	-40%
Yorkshire & the Humber	1,502	2,077	2,250	2,201	1,782	1,624	1,509	1,253	1,328	1,337	8%	1%	-11%
England	21,859	23,775	22,962	20,298	17,328	16,891	16,792	14,669	15,606	16,145	100%	3%	-26%

[‡] Percentage of total cases in England during 2010 which were in this SHA

Table 4 c (i): New episodes of gonorrhoea (uncomplicated) by age group & gender: England 2001 - 2010

MALES (total cases)											% change		
Age	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	10	15	20	10	12	9	6	4	4	4	0%	0%	-60%
15-19	1,987	2,185	1,990	1,870	1,484	1,460	1,555	1,283	1,250	1,108	10%	-11%	-44%
20-24	4,141	4,563	4,518	3,935	3,440	3,181	3,141	2,727	2,939	3,035	27%	3%	-27%
25-34	5,445	5,695	5,473	4,722	4,199	4,121	3,838	3,273	3,583	4,097	36%	14%	-25%
35-44	2,644	2,888	2,839	2,606	2,372	2,225	1,937	1,565	1,775	1,922	17%	8%	-27%
45+	972	1,057	1,072	1,032	918	989	922	823	1,011	1,095	10%	8%	13%
Total	15,309	16,570	15,930	14,190	12,429	12,006	11,493	9,729	10,572	11,289	100%	7%	-26%

MALES (MSM cases)											% change		
Age	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	2	0	1	2	1	0	0	0	0	1	0%	-	-50%
15-19	134	168	155	162	181	202	188	182	175	208	5%	19%	55%
20-24	540	576	641	606	654	707	608	538	654	910	20%	39%	69%
25-34	1,493	1,257	1,310	1,408	1,548	1,577	1,274	970	1,340	1,797	40%	34%	20%
35-44	869	820	973	976	1,057	995	794	609	819	1,048	23%	28%	21%
45+	253	231	293	319	326	394	313	248	387	522	12%	35%	106%
Total	3,333	3,184	3,527	3,491	3,788	3,928	3,228	2,597	3,379	4,503	100%	33%	35%

FEMALES (total cases)											% change		
Age	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	62	81	65	52	59	40	54	43	43	40	1%	-7%	-35%
15-19	2,681	2,843	2,873	2,508	2,006	1,885	2,076	2,004	1,863	1,749	36%	-6%	-35%
20-24	1,953	2,138	2,143	1,904	1,538	1,610	1,811	1,607	1,719	1,665	35%	-3%	-15%
25-34	1,239	1,442	1,301	1,169	887	923	942	919	971	952	20%	-2%	-23%
35-44	447	525	513	357	300	301	268	235	269	270	6%	0%	-40%
45+	118	114	134	110	107	126	124	130	136	140	3%	3%	19%
Total	6,550	7,205	7,032	6,108	4,899	4,885	5,299	4,940	5,004	4,825	100%	-4%	-26%

TOTAL ('Male', 'Female' and 'Not Known' gender)											% change		
Age	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	72	96	85	62	71	49	60	47	47	44	0%	-6%	-39%
15-19	4,668	5,028	4,863	4,378	3,490	3,345	3,631	3,287	3,121	2,861	18%	-8%	-39%
20-24	6,094	6,701	6,661	5,839	4,978	4,791	4,952	4,334	4,666	4,707	29%	1%	-23%
25-34	6,684	7,137	6,774	5,891	5,086	5,044	4,780	4,192	4,560	5,054	31%	11%	-24%
35-44	3,091	3,413	3,352	2,963	2,672	2,526	2,205	1,800	2,051	2,200	14%	7%	-29%
45+	1,090	1,171	1,206	1,142	1,025	1,115	1,046	953	1,148	1,242	8%	8%	14%
Total	21,859	23,775	22,962	20,298	17,328	16,891	16,792	14,669	15,606	16,145	100%	3%	-26%

† Percentage of total cases in England during 2010 which were in this SHA

Table 4 d: New episodes of genital chlamydia infection (uncomplicated) by SHA & gender: England 2001 - 2010

MALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	2,341	2,558	2,811	3,418	3,627	3,945	4,049	4,210	3,541	3,229	7%	-9%	38%
East of England	2,048	2,363	2,719	3,337	3,481	3,724	4,484	4,776	4,100	3,656	8%	-11%	79%
London	8,292	9,110	9,567	9,834	10,323	10,262	11,297	11,112	11,003	11,818	25%	7%	43%
North East	850	1,026	1,419	1,973	2,197	2,660	2,892	2,968	2,782	2,785	6%	0%	228%
North West	3,964	4,906	6,073	6,619	6,947	7,777	8,004	8,191	7,474	6,652	14%	-11%	68%
South Central	1,810	1,989	2,314	2,705	3,097	3,139	3,697	3,779	3,233	2,953	6%	-9%	63%
South East Coast	1,160	1,664	1,781	2,506	2,540	2,889	3,282	3,400	3,125	2,957	6%	-5%	155%
South West	2,298	2,517	2,589	3,085	3,305	4,273	4,590	4,468	3,783	3,732	8%	-1%	62%
West Midlands	3,043	3,432	3,568	3,447	4,032	4,384	4,755	4,781	5,005	4,635	10%	-7%	52%
Yorkshire & the Humber	3,265	4,209	4,556	5,125	5,055	5,248	5,731	6,012	5,230	4,710	10%	-10%	44%
England	29,071	33,774	37,397	42,049	44,604	48,301	52,791	53,697	49,276	47,127	100%	-4%	62%

MALES (MSM cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	28	27	52	65	80	71	99	92	120	167	3%	39%	496%
East of England	30	48	44	72	61	131	126	154	171	201	4%	18%	570%
London	760	681	770	755	1,052	1,282	1,425	1,531	1,908	2,479	50%	30%	226%
North East	20	22	29	72	106	132	109	133	163	218	4%	34%	990%
North West	194	249	337	293	302	401	477	633	722	723	15%	0%	273%
South Central	17	14	28	28	43	89	80	152	176	188	4%	7%	1006%
South East Coast	63	100	138	174	214	174	171	243	147	151	3%	3%	140%
South West	57	61	54	68	79	101	133	210	199	291	6%	46%	411%
West Midlands	54	64	85	76	78	114	139	195	187	275	6%	47%	409%
Yorkshire & the Humber	42	62	79	73	101	131	159	226	291	282	6%	-3%	571%
England	1,265	1,328	1,616	1,676	2,116	2,626	2,918	3,569	4,084	4,975	100%	22%	293%

FEMALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	3,370	3,792	3,794	4,354	4,470	4,458	4,860	4,697	3,622	3,067	7%	-15%	-9%
East of England	3,442	4,059	4,407	4,636	4,828	4,520	5,172	5,233	4,162	3,672	8%	-12%	7%
London	8,913	9,504	9,915	9,999	10,009	9,582	10,587	10,181	9,682	9,712	22%	0%	9%
North East	1,496	1,727	1,984	2,440	2,462	2,682	2,600	2,757	2,611	2,648	6%	1%	77%
North West	5,761	6,554	7,547	7,461	7,485	7,851	7,837	7,866	7,031	6,471	15%	-8%	12%
South Central	2,539	2,760	3,320	3,722	3,932	3,473	3,599	3,477	2,973	2,952	7%	-1%	16%
South East Coast	1,486	2,061	2,349	2,925	2,741	2,864	3,111	3,534	3,198	2,907	7%	-9%	96%
South West	2,775	2,964	3,134	3,330	3,606	3,933	3,962	4,224	3,687	3,384	8%	-8%	22%
West Midlands	3,532	4,026	4,344	4,167	4,328	4,298	4,667	5,271	5,190	4,464	10%	-14%	26%
Yorkshire & the Humber	4,496	5,435	5,763	6,263	5,955	5,548	6,101	6,186	5,319	4,566	10%	-14%	2%
England	37,810	42,882	46,557	49,297	49,816	49,209	52,496	53,426	47,475	43,843	100%	-8%	16%

TOTAL ('Male', 'Female' and 'Not Known' gender)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	5,711	6,350	6,605	7,772	8,097	8,403	8,909	8,907	7,178	6,304	7%	-12%	10%
East of England	5,490	6,422	7,126	7,973	8,309	8,244	9,666	10,009	8,262	7,328	8%	-11%	33%
London	17,205	18,614	19,482	19,833	20,332	19,844	21,884	21,293	20,776	21,624	24%	4%	26%
North East	2,346	2,753	3,403	4,413	4,659	5,342	5,492	5,725	5,393	5,433	6%	1%	132%
North West	9,725	11,460	13,620	14,080	14,432	15,628	15,841	16,057	14,507	13,123	14%	-10%	35%
South Central	4,349	4,749	5,634	6,427	7,029	6,612	7,296	7,256	6,209	5,905	6%	-5%	36%
South East Coast	2,646	3,725	4,130	5,431	5,281	5,753	6,393	6,934	6,323	5,864	6%	-7%	122%
South West	5,073	5,481	5,723	6,415	6,911	8,206	8,552	8,692	7,472	7,117	8%	-5%	40%
West Midlands	6,575	7,458	7,912	7,614	8,360	8,682	9,422	10,052	10,197	9,100	10%	-11%	38%
Yorkshire & the Humber	7,761	9,644	10,319	11,388	11,010	10,796	11,832	12,198	10,552	9,277	10%	-12%	20%
England	66,881	76,656	83,954	91,346	94,420	97,510	105,287	107,123	96,869	91,075	100%	-6%	36%

[‡] Percentage of total cases in England during 2010 which were in this SHA

Table 4 d (i): New episodes of genital chlamydia infection (uncomplicated) by age group & gender: England 2001 - 2010

MALES (total cases)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	21	26	29	27	26	32	29	32	25	26	0%	4%	24%
15-19	4,032	5,078	5,822	6,703	7,117	7,714	8,726	8,721	7,594	6,653	14%	-12%	65%
20-24	10,507	12,697	14,837	17,011	18,003	19,278	20,839	21,144	19,241	18,028	38%	-6%	72%
25-34	10,473	11,378	11,897	13,213	13,986	15,193	16,452	17,040	15,770	15,650	33%	-1%	49%
35-44	3,185	3,539	3,770	3,921	4,131	4,554	4,864	4,790	4,603	4,534	10%	-1%	42%
45+	831	938	999	1,154	1,338	1,522	1,771	1,946	2,010	2,190	5%	9%	164%
Total	29,071	33,774	37,397	42,049	44,604	48,301	52,791	53,697	49,276	47,127	100%	-4%	62%

FEMALES (total cases)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	215	217	271	277	291	296	303	324	234	234	1%	0%	9%
15-19	13,262	15,502	17,336	18,672	18,858	18,401	19,753	19,891	17,079	14,988	34%	-12%	13%
20-24	13,928	16,168	17,607	18,842	19,050	18,855	19,949	20,658	18,430	17,451	40%	-5%	25%
25-34	8,274	8,629	9,001	9,105	9,195	9,341	9,828	10,083	9,395	8,824	20%	-6%	7%
35-44	1,743	1,896	1,931	1,915	1,920	1,813	1,980	1,879	1,758	1,696	4%	-4%	-3%
45+	353	364	407	475	474	468	539	554	558	618	1%	11%	75%
Total	37,810	42,882	46,557	49,297	49,816	49,209	52,496	53,426	47,475	43,843	100%	-8%	16%

TOTAL ('Male', 'Female' and 'Not Known' gender)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	236	243	300	304	317	328	332	356	259	260	0%	0%	10%
15-19	17,294	20,580	23,158	25,375	25,975	26,115	28,479	28,612	24,693	21,655	24%	-12%	25%
20-24	24,435	28,865	32,444	35,853	37,053	38,133	40,788	41,802	37,710	35,505	39%	-6%	45%
25-34	18,747	20,007	20,898	22,318	23,181	24,534	26,280	27,123	25,203	24,519	27%	-3%	31%
35-44	4,928	5,435	5,701	5,836	6,051	6,367	6,844	6,669	6,376	6,246	7%	-2%	27%
45+	1,184	1,302	1,406	1,629	1,812	1,990	2,310	2,500	2,574	2,812	3%	9%	138%
Total	66,881	76,656	83,954	91,346	94,420	97,510	105,287	107,123	96,869	91,075	100%	-6%	36%

† Percentage of total cases in England during 2010 which were in this SHA

Table 4 e: New episodes of genital herpes (first episode) by SHA & gender: England 2001 - 2010

MALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	344	371	366	377	391	536	677	682	746	804	7%	8%	134%
East of England	595	577	561	554	556	638	606	721	838	984	8%	17%	65%
London	1,887	1,945	2,007	1,912	1,946	2,045	2,391	2,447	2,726	2,856	25%	5%	51%
North East	169	154	174	156	230	268	370	400	484	524	5%	8%	210%
North West	703	683	652	730	933	977	1,220	1,412	1,536	1,572	14%	2%	124%
South Central	513	445	505	492	463	583	674	801	700	848	7%	21%	65%
South East Coast	427	492	464	476	437	506	653	769	821	912	8%	11%	114%
South West	612	598	550	549	605	540	716	898	913	966	8%	6%	58%
West Midlands	611	611	540	579	615	616	827	924	999	1,031	9%	3%	69%
Yorkshire & the Humber	550	569	550	531	554	747	921	1,050	1,061	1,087	9%	2%	98%
England	6,411	6,445	6,369	6,356	6,730	7,456	9,055	10,104	10,824	11,584	100%	7%	81%

MALES (MSM cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	9	15	11	9	9	15	13	21	12	41	4%	242%	356%
East of England	14	21	14	18	22	19	12	34	25	38	4%	52%	171%
London	238	272	279	254	293	280	358	365	472	514	53%	9%	116%
North East	6	8	9	6	15	9	18	22	20	21	2%	5%	250%
North West	32	53	35	47	72	56	57	95	96	121	12%	26%	278%
South Central	14	4	11	20	13	29	23	29	31	35	4%	13%	150%
South East Coast	26	43	47	45	36	41	37	53	58	53	5%	-9%	104%
South West	27	14	21	22	25	17	27	38	23	48	5%	109%	78%
West Midlands	16	16	22	15	31	23	30	47	38	47	5%	24%	194%
Yorkshire & the Humber	9	22	8	12	22	26	25	31	43	55	6%	28%	511%
England	391	468	457	448	538	515	600	735	818	973	100%	19%	149%

FEMALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	673	723	722	724	695	791	1,033	1,042	1,101	1,249	7%	13%	86%
East of England	1,073	1,038	951	955	1,014	1,086	1,268	1,365	1,433	1,656	9%	16%	54%
London	2,729	2,772	2,853	2,762	2,762	3,018	3,629	3,669	3,907	4,177	23%	7%	53%
North East	295	326	293	318	389	465	632	599	735	783	4%	7%	165%
North West	1,235	1,204	1,225	1,313	1,425	1,657	2,022	2,404	2,471	2,473	14%	0%	100%
South Central	807	816	851	769	813	916	1,080	1,218	1,192	1,361	8%	14%	69%
South East Coast	683	823	800	831	770	838	1,047	1,209	1,321	1,498	8%	13%	119%
South West	965	942	816	773	830	835	1,042	1,326	1,426	1,495	8%	5%	55%
West Midlands	979	1,100	994	949	985	1,037	1,212	1,502	1,541	1,698	9%	10%	73%
Yorkshire & the Humber	956	1,070	1,029	944	966	1,155	1,467	1,656	1,576	1,700	9%	8%	78%
England	10,395	10,814	10,534	10,338	10,649	11,798	14,432	15,990	16,703	18,090	100%	8%	74%

TOTAL ('Male', 'Female' and 'Not Known' gender)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	1,017	1,094	1,088	1,101	1,086	1,327	1,710	1,724	1,850	2,055	7%	11%	102%
East of England	1,668	1,615	1,512	1,509	1,570	1,724	1,874	2,086	2,271	2,640	9%	16%	58%
London	4,616	4,717	4,860	4,674	4,708	5,063	6,020	6,116	6,665	7,060	24%	6%	53%
North East	464	480	467	474	619	733	1,002	999	1,219	1,307	4%	7%	182%
North West	1,938	1,887	1,877	2,043	2,358	2,634	3,242	3,816	4,007	4,045	14%	1%	109%
South Central	1,320	1,261	1,356	1,261	1,276	1,499	1,754	2,019	1,892	2,209	7%	17%	67%
South East Coast	1,110	1,315	1,264	1,307	1,207	1,344	1,700	1,978	2,142	2,410	8%	13%	117%
South West	1,577	1,540	1,366	1,322	1,435	1,375	1,758	2,224	2,340	2,461	8%	5%	56%
West Midlands	1,590	1,711	1,534	1,528	1,600	1,653	2,039	2,426	2,541	2,729	9%	7%	72%
Yorkshire & the Humber	1,506	1,639	1,579	1,475	1,520	1,902	2,388	2,706	2,637	2,787	9%	6%	85%
England	16,806	17,259	16,903	16,694	17,379	19,254	23,487	26,094	27,564	29,703	100%	8%	77%

[‡] Percentage of total cases in England during 2010 which were in this SHA

Table 4 e (j): New episodes of genital herpes (first episode) by age group & gender: England 2001 - 2010

MALES (total cases)											% change		
Age	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	2	3	1	1	2	1	0	5	2	2	0%	0%	0%
15-19	386	384	355	397	504	542	642	709	695	757	7%	9%	96%
20-24	1,423	1,403	1,368	1,477	1,596	1,782	2,254	2,511	2,739	2,846	25%	4%	100%
25-34	2,657	2,470	2,557	2,433	2,555	2,713	3,213	3,640	3,965	4,220	36%	6%	59%
35-44	1,250	1,431	1,443	1,375	1,334	1,539	1,789	1,926	2,026	2,140	18%	6%	71%
45+	684	725	644	672	733	874	1,112	1,308	1,389	1,604	14%	15%	135%
Total	6,411	6,445	6,369	6,356	6,730	7,456	9,055	10,104	10,824	11,584	100%	7%	81%

FEMALES (total cases)											% change		
Age	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	45	35	45	30	33	32	56	62	56	46	0%	-18%	2%
15-19	2,101	2,020	2,089	2,145	2,143	2,533	3,093	3,374	3,249	3,388	19%	4%	61%
20-24	2,875	3,115	2,969	2,905	3,152	3,554	4,227	4,562	4,951	5,311	29%	7%	85%
25-34	3,333	3,430	3,307	3,095	3,207	3,416	4,150	4,632	4,917	5,374	30%	9%	61%
35-44	1,354	1,463	1,427	1,413	1,365	1,488	1,799	2,065	2,072	2,313	13%	12%	71%
45+	665	707	697	747	740	772	1,050	1,292	1,441	1,648	9%	14%	148%
Total	10,395	10,814	10,534	10,338	10,649	11,798	14,432	15,990	16,703	18,090	100%	8%	74%

TOTAL ('Male', 'Female' and 'Not Known' gender)											% change		
Age	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	47	38	46	31	35	33	56	67	58	48	0%	-17%	2%
15-19	2,487	2,404	2,444	2,542	2,647	3,075	3,735	4,083	3,945	4,145	14%	5%	67%
20-24	4,298	4,518	4,337	4,382	4,748	5,336	6,481	7,073	7,703	8,162	27%	6%	90%
25-34	5,990	5,900	5,864	5,528	5,762	6,129	7,363	8,272	8,898	9,605	32%	8%	60%
35-44	2,604	2,894	2,870	2,788	2,699	3,027	3,588	3,991	4,100	4,459	15%	9%	71%
45+	1,349	1,432	1,341	1,419	1,473	1,646	2,162	2,600	2,835	3,259	11%	15%	142%
Total	16,806	17,259	16,903	16,694	17,379	19,254	23,487	26,094	27,564	29,703	100%	8%	77%

† Percentage of total cases in England during 2010 which were in this SHA

Table 4 f: New episodes of genital warts (first episode) by SHA & gender: England 2001 - 2010

MALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	2,406	2,359	2,365	2,620	2,654	2,699	2,916	2,714	2,997	2,766	7%	-8%	15%
East of England	3,085	3,109	3,244	3,369	3,478	3,464	3,670	3,813	3,913	3,771	9%	-4%	22%
London	6,457	7,007	6,807	7,131	6,785	6,964	7,609	7,567	7,792	7,615	19%	-2%	18%
North East	1,609	1,708	1,792	2,018	2,019	2,220	2,432	2,410	2,408	2,520	6%	5%	57%
North West	4,814	4,873	5,360	5,493	5,735	5,896	6,220	6,462	6,600	6,135	15%	-7%	27%
South Central	2,353	2,474	2,479	2,472	2,492	2,647	2,742	2,918	2,966	2,944	7%	-1%	25%
South East Coast	2,473	2,785	2,499	2,599	2,655	2,794	2,901	3,126	3,376	3,330	8%	-1%	35%
South West	3,057	3,121	3,296	3,449	3,471	3,514	3,588	3,808	3,714	3,803	9%	2%	24%
West Midlands	2,861	2,841	2,856	3,150	3,099	3,174	3,474	3,812	3,828	3,797	9%	-1%	33%
Yorkshire & the Humber	3,170	3,265	3,500	3,671	3,587	3,649	4,171	4,464	4,340	4,272	10%	-2%	35%
England	32,285	33,542	34,198	35,972	35,975	37,021	39,723	41,094	41,934	40,953	100%	-2%	27%

MALES (MSM cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	74	51	79	78	86	97	73	66	106	100	4%	-6%	35%
East of England	81	97	101	129	126	116	109	109	134	142	6%	6%	75%
London	865	960	959	887	967	1,054	1,141	1,148	1,102	1,001	40%	-9%	16%
North East	57	53	81	82	100	100	117	111	148	160	6%	8%	181%
North West	212	236	292	311	326	323	330	360	393	358	14%	-9%	69%
South Central	39	49	64	68	84	69	91	71	115	141	6%	23%	262%
South East Coast	142	193	192	160	190	175	143	185	155	140	6%	-10%	-1%
South West	113	91	97	108	122	133	114	161	117	163	6%	39%	44%
West Midlands	86	101	121	100	130	126	119	144	127	146	6%	15%	70%
Yorkshire & the Humber	75	100	88	80	94	87	102	133	160	177	7%	11%	136%
England	1,744	1,931	2,074	2,003	2,225	2,280	2,339	2,488	2,557	2,528	100%	-1%	45%

FEMALES (total cases)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	2,193	2,051	2,083	2,353	2,365	2,387	2,658	2,642	2,501	2,435	7%	-3%	11%
East of England	2,878	2,864	2,932	3,179	3,277	3,186	3,432	3,504	3,535	3,355	10%	-5%	17%
London	5,352	5,529	5,348	5,504	5,462	5,605	6,086	6,080	5,991	5,849	17%	-2%	9%
North East	1,549	1,631	1,597	1,795	1,849	1,959	2,293	2,255	2,206	2,134	6%	-3%	38%
North West	4,200	4,065	4,696	4,623	4,946	5,174	5,322	5,670	5,465	5,246	15%	-4%	25%
South Central	2,193	2,243	2,343	2,237	2,327	2,429	2,694	2,730	2,650	2,531	7%	-4%	15%
South East Coast	2,127	2,365	2,205	2,504	2,506	2,540	2,863	2,969	2,979	2,880	8%	-3%	35%
South West	2,865	2,999	3,018	3,038	3,221	3,117	3,173	3,481	3,424	3,281	9%	-4%	15%
West Midlands	2,742	2,548	2,646	2,716	2,845	2,878	3,213	3,570	3,385	3,337	10%	-1%	22%
Yorkshire & the Humber	3,121	3,145	3,253	3,330	3,079	3,404	3,815	4,161	3,768	3,534	10%	-6%	13%
England	29,220	29,440	30,121	31,279	31,877	32,679	35,549	37,062	35,904	34,582	100%	-4%	18%

TOTAL ('Male', 'Female' and 'Not Known' gender)											% change		
SHA	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total[‡]	2009-2010	2001-2010
East Midlands	4,599	4,410	4,448	4,973	5,019	5,086	5,574	5,356	5,510	5,212	7%	-5%	13%
East of England	5,963	5,973	6,176	6,548	6,755	6,650	7,102	7,317	7,448	7,126	9%	-4%	20%
London	11,809	12,536	12,155	12,635	12,247	12,569	13,695	13,647	13,829	13,531	18%	-2%	15%
North East	3,158	3,339	3,389	3,813	3,868	4,179	4,725	4,665	4,614	4,654	6%	1%	47%
North West	9,014	8,938	10,056	10,116	10,681	11,070	11,542	12,132	12,066	11,381	15%	-6%	26%
South Central	4,546	4,717	4,822	4,709	4,819	5,076	5,436	5,648	5,617	5,475	7%	-3%	20%
South East Coast	4,600	5,150	4,704	5,103	5,161	5,334	5,764	6,095	6,355	6,210	8%	-2%	35%
South West	5,922	6,120	6,314	6,487	6,692	6,631	6,761	7,289	7,139	7,085	9%	-1%	20%
West Midlands	5,603	5,389	5,502	5,866	5,944	6,052	6,687	7,382	7,214	7,134	9%	-1%	27%
Yorkshire & the Humber	6,291	6,410	6,753	7,001	6,666	7,053	7,986	8,625	8,108	7,807	10%	-4%	24%
England	61,505	62,982	64,319	67,251	67,852	69,700	75,272	78,156	77,900	75,615	100%	-3%	23%

[‡] Percentage of total cases in England during 2010 which were in this SHA

Table 4 f (j): New episodes of genital warts (first episode) by age group & gender: United Kingdom 2000 - 2009

MALES (total cases)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	31	29	35	29	28	26	26	14	25	26	0%	4%	-16%
15-19	3,050	3,122	3,378	3,799	3,759	3,974	4,446	4,695	4,650	4,306	11%	-7%	41%
20-24	10,713	11,327	11,737	12,579	12,697	12,935	13,869	14,473	14,636	14,464	35%	-1%	35%
25-34	12,303	12,366	12,311	12,489	12,553	12,744	13,564	13,825	14,254	13,928	34%	-2%	13%
35-44	4,297	4,671	4,672	4,814	4,638	4,913	5,057	5,086	5,219	4,987	12%	-4%	16%
45+	1,851	1,906	2,060	2,252	2,294	2,425	2,613	2,995	3,126	3,213	8%	3%	74%
Total	32,285	33,542	34,198	35,972	35,975	37,021	39,723	41,094	41,934	40,953	100%	-2%	27%

FEMALES (total cases)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	112	126	120	117	111	112	127	135	158	120	0%	-24%	7%
15-19	8,333	8,396	8,814	9,421	9,731	10,251	11,207	11,669	11,225	10,101	29%	-10%	21%
20-24	9,912	9,850	10,156	10,470	10,835	11,194	11,945	12,636	12,143	12,303	36%	1%	24%
25-34	7,453	7,519	7,378	7,488	7,461	7,413	8,029	8,316	7,927	7,689	22%	-3%	3%
35-44	2,303	2,426	2,535	2,704	2,667	2,565	2,869	2,873	2,980	2,856	8%	-4%	24%
45+	1,054	1,031	1,114	1,068	1,066	1,143	1,291	1,427	1,449	1,491	4%	3%	41%
Total	29,220	29,440	30,121	31,279	31,877	32,679	35,549	37,062	35,904	34,582	100%	-4%	18%

TOTAL ('Male', 'Female' and 'Not Known' gender)

Age											% change		
	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	% of total†	2009-2010	2001-2010
<15	143	155	155	146	139	138	153	149	183	146	0%	-20%	2%
15-19	11,383	11,518	12,192	13,220	13,490	14,225	15,653	16,364	15,887	14,417	19%	-9%	27%
20-24	20,625	21,177	21,893	23,049	23,532	24,129	25,814	27,109	26,796	26,791	35%	0%	30%
25-34	19,756	19,885	19,689	19,977	20,014	20,157	21,593	22,141	22,203	21,649	29%	-2%	10%
35-44	6,600	7,097	7,207	7,518	7,305	7,478	7,926	7,959	8,205	7,853	10%	-4%	19%
45+	2,905	2,937	3,174	3,320	3,360	3,568	3,904	4,422	4,580	4,708	6%	3%	62%
Total	61,505	62,982	64,319	67,251	67,852	69,700	75,272	78,156	77,900	75,615	100%	-3%	23%

† Percentage of total cases in England during 2010 which were in this SHA

Supplementary Tables of Sexually Transmitted Infections in England



Code Groups
New STI diagnoses
Chlamydial infection (uncomplicated and complicated)
Gonorrhoea (uncomplicated and complicated)
Infectious and early latent syphilis
Genital herpes simplex (first episode)
Genital warts (first episode)
New HIV diagnosis
Non-specific genital infection (uncomplicated and complicated)
Chancroid/lymphogranuloma venerum (LGV)/Donovanosis
Molluscum contagiosum
Trichomoniasis
Scabies
Pediculus pubis
Other STI diagnoses
Congenital and other acquired syphilis
Recurrent genital herpes simplex
Recurrent and re-registered genital warts
Subsequent HIV presentations (including AIDS)
Ophthalmia neonatorum (chamydial or gonococcal)
Epidemiological treatment of suspected STIs (syphilis, chlamydia, gonorrhoea, non-specific genital infection)
Other diagnoses made at GUM clinics
Viral hepatitis B
Viral hepatitis C
Vaginosis and balanitis (including epidemiological treatment)
Anogenital candidiasis (including epidemiological treatment)
Urinary tract infection
Cervical abnormalities
Other conditions requiring treatment at a GUM clinic
Services provided
HIV antibody test
Sexual health screen
Hepatitis B vaccination
Contraception (excluding condom provision)
Other episode not requiring treatment