
Table 3: Total number of STI diagnoses & other episodes of care seen at genitourinary medicine clinics in England: 2001 - 2010



Data are presented by SHA, gender & sexual orientation

Notes on using these tables:

Numbers of diagnoses are adjusted for missing clinic data
Data on unknown gender are included in the 'Total'

An improved STI surveillance system (GUMCAD) was introduced in England in 2009. Unlike the KC60 return, duplicate records in GUMCAD can be identified and removed resulting in a slight reduction in the number of diagnoses reported. England (SHA) data from 2000-2008 have therefore been statistically adjusted to enable fair comparisons of STI diagnoses over time.

Definitions of data totals:

Total new STI diagnoses:	}	Refer to 'Code Groups' for conditions included
Total other STI diagnoses:		
Total other GUM clinic diagnoses:		
Total services provided:		

Data source:

England: GUM clinic KC60 returns (2001-2008)
 GUM clinic GUMCAD returns (2009-2010)

Data supplied by:

Health Protection Agency, Health Protection Services Colindale

Data prepared by:

HIV and Sexually Transmitted Infections Department
Health Protection Agency
Health Protection Services Colindale
61 Colindale Avenue
London
NW9 5EQ

Tel. 020-8200-6868
Email: hiv/sti@hpa.org.uk
Website: www.hpa.org.uk

Table 3: Total number of STI diagnoses & other episodes of care seen at genitourinary medicine clinics in England: 2001 - 2010
Data are presented by SHA, gender & sexual orientation



Data tables

East Midlands	3 b:
East of England	3 c:
London	3 d:
North East	3 e:
North West	3 f:
South Central	3 g:
South East Coast	3 h:
South West	3 i:
West Midlands	3 j:
Yorkshire & the Humber	3 k:

Code groups: Definitions of conditions presented

Further Information

England

Data on selected diagnoses, 1971 to 2010, are available upon request

HPA Health Protection Services Colindale

Contact: Geraldine Leong

Tel: +44 (0)20 8200 6868

Email: geraldine.leong@hpa.org.uk

Web: www.hpa.org.uk



3 b: East Midlands SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	15	21	55	80	139	122	87	93	116	114
		-of which MSM	5	8	15	28	48	44	39	42	66	66
		Female	2	3	8	18	43	40	39	20	14	18
		Total	17	24	63	98	182	162	126	113	130	132
A3	Early latent syphilis (first 2 years)	Male (all)	13	4	6	17	36	26	23	32	20	31
		-of which MSM	4	0	2	5	14	10	7	12	12	17
		Female	4	0	3	16	14	18	15	11	17	12
		Total	17	4	9	33	50	44	38	43	37	43
A4,A5,A6	Other acquired syphilis	Male (all)	24	22	35	44	54	63	47	34	47	44
		-of which MSM	1	1	1	1	10	8	10	3	13	12
		Female	17	23	23	37	50	29	38	33	36	36
		Total	41	45	58	81	104	92	85	67	83	80
A7	Congenital syphilis, aged under 2	Male (all)	0	0	0	0	0	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	0	0	0	0	0
Total	0	0	0	0	0	0	0	0	0	0		
A8	Congenital syphilis, aged 2 or over	Male (all)	0	0	0	0	1	0	0	0	1	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	2	1	1	0	0	0	0	1	2
Total	1	2	1	1	1	0	0	0	0	2	2	
A9	Epidemiological treatment of suspected syphilis	Male (all)	2	4	5	10	26	40	34	26	40	27
		-of which MSM	2	3	3	4	2	17	13	10	20	11
		Female	0	0	8	12	27	19	15	8	9	11
		Total	2	4	13	22	53	59	49	34	49	38
B1,B2	Uncomplicated gonorrhoea	Male (all)	884	1,011	945	848	715	783	746	543	631	564
		-of which MSM	104	84	126	124	108	124	86	54	88	148
		Female	477	528	492	401	338	381	476	391	443	419
		Total	1,361	1,539	1,437	1,249	1,053	1,164	1,222	934	1,082	985
B3	Gonococcal ophthalmia neonatorum	Male (all)	0	1	1	0	1	1	3	1	0	1
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	1	0	0	0	0	6	1	1	0
Total	0	2	1	0	1	1	9	2	1	1		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	121	137	165	169	164	167	210	192	263	261
		-of which MSM	26	25	43	28	26	33	28	21	30	58
		Female	162	204	205	217	206	249	263	220	240	216
		Total	283	341	370	386	370	416	473	412	505	477
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	6	8	8	9	8	7	9	5	13	10
		-of which MSM	0	0	0	0	1	2	2	0	1	3
		Female	13	15	24	14	18	13	29	22	16	19
		Total	19	23	32	23	26	20	38	27	29	29

East Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	1	2	4	3	2	0	6	9	8	3
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	1	2	1	1	5	11	8	5	5
		Total	1	3	6	4	3	5	17	17	13	8
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	2,341	2,558	2,811	3,418	3,627	3,945	4,049	4,210	3,541	3,229
		-of which MSM	28	27	52	65	80	71	99	92	120	167
		Female	3,370	3,792	3,794	4,354	4,470	4,458	4,860	4,697	3,622	3,067
		Total	5,711	6,350	6,605	7,772	8,097	8,403	8,909	8,907	7,178	6,304
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	36	34	37	69	67	94	64	50	36	66
		-of which MSM	1	0	1	3	1	2	3	2	1	4
		Female	210	263	271	267	244	205	277	225	183	161
		Total	246	297	308	336	311	299	341	275	219	227
C4D	Chlamydial ophthalmia neonatorum	Male (all)	0	0	0	2	0	1	0	2	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	2	1	4	2	0	0	1	0	2	1
		Total	2	1	4	4	0	1	1	2	2	1
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	1,400	1,715	1,739	2,014	2,259	2,061	2,468	2,365	1,922	1,860
		-of which MSM	13	27	28	43	55	49	51	34	69	99
		Female	689	946	1,046	1,278	1,565	1,541	1,536	1,603	1,377	1,407
		Total	2,089	2,661	2,785	3,292	3,824	3,602	4,004	3,968	3,310	3,271
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	2,823	3,026	3,649	3,926	3,906	4,074	4,335	3,617	3,398	3,202
		-of which MSM	92	69	101	121	134	174	158	141	159	202
		Female	0	0	233	215	186	216	296	219	211	207
		Total	2,823	3,026	3,882	4,141	4,092	4,290	4,631	3,836	3,615	3,410
C4I	Epidemiological treatment of NSGI	Male (all)	384	397	409	361	307	296	369	247	228	250
		-of which MSM	36	25	39	43	31	37	28	30	26	29
		Female	1,092	1,048	1,076	1,080	979	1,113	1,277	989	889	770
		Total	1,476	1,445	1,485	1,441	1,286	1,409	1,646	1,236	1,120	1,021
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	216	235	235	267	321	322	357	334	401	338
		-of which MSM	6	4	9	13	7	9	7	14	24	22
		Female	663	712	637	640	627	722	901	783	721	758
		Total	879	947	872	907	948	1,044	1,258	1,117	1,123	1,096
C6A	Trichomoniasis	Male (all)	6	12	9	5	6	10	15	29	21	29
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	325	382	389	396	344	394	389	376	333	340
		Total	331	394	398	401	350	404	404	405	354	369
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	112	121	113	46	64	104	139	92	102	131
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3,600	4,122	4,110	4,047	3,917	3,950	5,110	5,042	5,041	4,896
		Total	3,712	4,243	4,223	4,093	3,981	4,054	5,249	5,134	5,158	5,034
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	1,210	1,068	1,103	1,233	983	957	870	903	797	658
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1,142	1,034	992	882	593	497	481	508	329	198
		Total	2,352	2,102	2,095	2,115	1,576	1,454	1,351	1,411	1,126	856

East Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C7A	Anogenital candidosis	Male (all)	422	400	259	280	351	338	490	530	505	650
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	4,048	4,300	4,122	3,912	4,074	4,373	5,291	4,985	4,766	4,736
		Total	4,470	4,700	4,381	4,192	4,425	4,711	5,781	5,515	5,282	5,394
C7B	Epidemiological treatment of C6 & C7	Male (all)	270	269	251	243	314	244	392	270	306	305
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	372	312	377	446	488	433	388	346	348	348
		Total	642	581	628	689	802	677	780	616	654	653
C8,C9	Scabies/pediculosis pubis	Male (all)	298	237	223	185	116	126	100	106	94	100
		-of which MSM	44	22	18	23	17	21	18	13	19	24
		Female	81	68	35	35	16	17	23	14	12	15
		Total	379	305	258	220	132	143	123	120	106	115
C10A	Anogenital herpes simplex – first episode	Male (all)	344	371	366	377	391	536	677	682	746	804
		-of which MSM	9	15	11	9	9	15	13	21	12	41
		Female	673	723	722	724	695	791	1,033	1,042	1,101	1,249
		Total	1,017	1,094	1,088	1,101	1,086	1,327	1,710	1,724	1,850	2,055
C10B	Anogenital herpes simplex – recurrence	Male (all)	266	297	281	232	244	277	404	417	494	579
		-of which MSM	7	6	12	6	9	12	8	6	18	17
		Female	357	421	394	413	376	431	471	522	628	736
		Total	623	718	675	645	620	708	875	939	1,122	1,315
C11A	Anogenital warts – first episode	Male (all)	2,406	2,359	2,365	2,620	2,654	2,699	2,916	2,714	2,997	2,766
		-of which MSM	74	51	79	78	86	97	73	66	106	100
		Female	2,193	2,051	2,083	2,353	2,365	2,387	2,658	2,642	2,501	2,435
		Total	4,599	4,410	4,448	4,973	5,019	5,086	5,574	5,356	5,510	5,212
C11B	Anogenital warts – recurrence	Male (all)	1,585	1,542	1,399	1,523	1,449	1,576	1,878	1,815	1,475	1,374
		-of which MSM	48	38	63	44	38	64	48	37	38	69
		Female	870	812	806	830	872	920	1,144	1,138	972	933
		Total	2,455	2,354	2,205	2,353	2,321	2,496	3,022	2,953	2,449	2,308
C11C	Anogenital warts – re-registered cases	Male (all)	529	472	473	472	488	512	586	502	526	487
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	450	499	409	416	384	420	430	394	422	330
Total	979	971	882	888	872	932	1,016	896	948	817		
C12	Molluscum contagiosum	Male (all)	354	357	440	572	572	674	663	711	669	590
		-of which MSM	10	10	8	8	11	11	10	6	7	10
		Female	182	194	228	226	240	283	284	309	245	243
		Total	536	551	668	798	812	957	947	1,020	914	834
C13	Antigen positive viral hepatitis B	Male (all)	37	41	-	-	-	-	-	-	-	-
		-of which MSM	3	5	-	-	-	-	-	-	-	-
		Female	10	14	-	-	-	-	-	-	-	-
		Total	47	55	-	-	-	-	-	-	-	-
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	24	31	32	21	31	28	43	37
		-of which MSM	-	-	2	2	3	1	0	1	1	5
		Female	-	-	5	15	10	8	12	19	19	15
		Total	-	-	29	46	42	29	43	47	63	52

East Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	0	1	0	0	3	0	3	1
		-of which MSM	-	-	0	0	0	0	0	0	0	0
		Female	-	-	0	0	1	1	0	0	1	1
		Total	-	-	0	1	1	1	3	0	4	2
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	5	3	2	4	8	11	22	4
		-of which MSM	-	-	1	1	1	0	0	1	3	0
		Female	-	-	0	1	2	0	5	13	2	4
		Total	-	-	5	4	4	4	13	24	24	8
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	72	97	83	79	64	78	76	83	51	32
		-of which MSM	1	1	0	5	1	2	2	3	3	2
		Female	18	19	19	12	23	10	13	15	13	9
		Total	90	116	102	91	87	88	89	98	64	41
E1A	New HIV diagnosis - asymptomatic	Male (all)	38	73	176	158	139	136	233	114	85	100
		-of which MSM	11	15	28	33	38	42	61	30	47	53
		Female	47	84	203	213	202	140	334	105	58	67
		Total	85	157	379	371	341	276	567	219	143	167
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	219	360	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	108	258	-	-	-	-	-	-	-	-
		Total	327	618	-	-	-	-	-	-	-	-
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	536	841	996	1,053	1,191	896	745	626
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	304	649	896	960	1,242	918	709	621
		Total	-	-	840	1,490	1,892	2,013	2,433	1,814	1,454	1,247
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	26	24	41	16	34	15	26	21	15	20
		-of which MSM	7	6	12	5	6	1	1	4	4	7
		Female	7	20	29	14	26	21	22	24	18	9
		Total	33	44	70	30	60	36	48	45	33	29
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	313	375	-	-	-	-	-	-	-	-
		-of which MSM	0	0	-	-	-	-	-	-	-	-
		Female	63	148	-	-	-	-	-	-	-	-
		Total	376	523	-	-	-	-	-	-	-	-
E3A*	AIDS – first presentation	Male (all)	22	22	-	-	-	-	-	-	-	-
		-of which MSM	6	5	-	-	-	-	-	-	-	-
		Female	3	8	-	-	-	-	-	-	-	-
		Total	25	30	-	-	-	-	-	-	-	-
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	7	6	7	7	9	10	18	4
		-of which MSM	-	-	0	0	1	0	0	1	1	0
		Female	-	-	6	5	7	4	3	11	12	4
		Total	-	-	13	11	14	11	12	21	30	8
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	3	2	1	3	0	7	2	2
		-of which MSM	-	-	0	0	0	0	0	2	1	0
		Female	-	-	1	0	3	3	1	1	0	13
		Total	-	-	4	2	4	6	1	8	2	15

East Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
E3B	AIDS – subsequent presentation	Male (all)	426	143	158	147	196	198	282	157	92	104
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	215	38	56	69	65	44	183	97	57	59
		Total	641	181	214	216	261	242	465	254	149	163
D2A	Urinary tract infection	Male (all)	124	135	120	108	77	79	85	86	111	107
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	466	517	436	494	426	446	563	590	568	567
		Total	590	652	556	602	503	525	648	676	680	676
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	3,887	4,060	4,094	4,118	3,927	4,044	4,900	4,132	4,721	4,475
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3,303	3,482	3,467	3,341	3,165	3,116	3,443	3,167	3,627	3,689
		Total	7,190	7,542	7,561	7,459	7,092	7,160	8,343	7,299	8,362	8,167
D3	Other episodes not requiring treatment	Male (all)	8,281	9,837	8,276	9,891	12,176	13,766	15,571	15,709	16,796	15,622
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	7,968	9,456	8,432	9,981	12,074	13,207	16,230	16,666	16,294	15,756
		Total	16,249	19,293	16,708	19,872	24,250	26,973	31,801	32,375	33,136	31,416
P1A	HIV antibody counselling - with testing	Male (all)	8,113	10,499	1,946	1,657	1,665	1,687	1,860	1,763	1,399	1,304
		-of which MSM	511	507	194	185	169	162	169	133	182	218
		Female	6,632	9,561	1,498	1,330	1,485	1,249	1,721	1,620	1,385	1,367
		Total	14,745	20,060	3,444	2,987	3,150	2,936	3,581	3,383	2,785	2,675
P1B	HIV antibody counselling - without testing	Male (all)	2,618	2,952	5,571	7,229	8,348	6,947	7,846	8,080	6,956	6,433
		-of which MSM	54	46	157	147	205	203	162	175	154	208
		Female	2,738	3,370	6,117	7,900	8,840	7,127	8,990	9,867	8,735	8,186
		Total	5,356	6,322	11,688	15,129	17,188	14,074	16,836	17,947	15,729	14,635
P2	Hepatitis B vaccination	Male (all)	719	833	876	941	1,015	1,105	1,147	1,034	1,141	1,168
		-of which MSM	202	180	294	297	335	350	326	258	414	480
		Female	212	268	295	476	504	439	459	462	499	677
		Total	931	1,101	1,171	1,417	1,519	1,544	1,606	1,496	1,641	1,845
P3	Family Planning	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	851	809	749	952	853	913	907	1,068	918	1,010
		Total	851	809	749	952	853	913	907	1,068	918	1,010
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	340	184	95	99	66	60	87	62	40	29
		Total	340	184	95	99	66	60	87	62	40	29

East Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	63	51	27	32	17	11	17	9	12	3
		Total	63	51	27	32	17	11	17	9	12	3
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	9,237	10,518	8,858	7,738	8,038	8,143	7,285	6,688
		-of which MSM	-	-	243	248	216	238	208	186	218	301
		Female	-	-	10,801	11,949	10,210	8,880	10,335	10,956	9,718	9,028
		Total	-	-	20,038	22,467	19,068	16,618	18,373	19,099	17,056	15,729
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	10,270	16,027	18,973	22,196	26,045	25,728	25,767	26,119
		-of which MSM	-	-	593	791	850	1,126	1,051	990	1,558	2,139
		Female	-	-	10,141	15,079	18,254	21,316	26,197	25,986	24,699	25,579
		Total	-	-	20,411	31,106	37,227	43,512	52,242	51,714	50,526	51,749
Total new STI diagnoses (for GUM clinics only)	Male (all)	9,807	10,332	11,377	12,576	12,740	13,576	14,315	13,280	12,809	11,970	
	-of which MSM	395	311	462	515	561	623	577	498	667	864	
	Female	8,247	8,836	9,159	9,892	9,836	10,095	11,650	10,899	9,512	9,028	
	Total	18,054	19,168	20,536	22,468	22,576	23,671	25,965	24,179	22,366	21,023	
Total other STI diagnoses	Male (all)	5,291	5,487	5,204	5,817	6,186	6,248	7,472	6,661	5,835	5,615	
	-of which MSM	139	130	189	169	171	220	186	143	215	295	
	Female	4,029	4,409	4,333	5,004	5,423	5,729	6,607	5,924	5,343	5,135	
	Total	9,320	9,896	9,537	10,821	11,609	11,977	14,079	12,585	11,196	10,756	
Total other GUM clinic diagnoses	Male (all)	6,134	6,191	6,052	6,142	5,814	5,869	6,994	6,135	6,661	6,400	
	-of which MSM	4	6	3	8	5	3	2	5	7	7	
	Female	13,362	14,035	13,650	13,281	12,782	12,905	15,410	14,756	14,766	14,495	
	Total	19,496	20,226	19,702	19,423	18,596	18,774	22,404	20,891	21,469	20,915	
Total services provided	Male (all)	19,731	24,121	36,176	46,263	51,035	53,439	60,507	60,457	59,344	57,334	
	-of which MSM	767	733	1,481	1,668	1,775	2,079	1,916	1,742	2,526	3,346	
	Female	18,401	23,464	38,033	47,667	52,220	53,131	64,839	66,625	62,248	61,603	
	Total	38,132	47,585	74,209	93,930	103,255	106,570	125,346	127,082	121,791	119,059	

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 c: East of England SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	16	27	64	70	69	71	59	54	49	50
		-of which MSM	9	12	28	23	28	30	24	17	26	27
		Female	3	9	14	18	13	9	10	17	8	8
		Total	19	36	78	88	82	80	69	71	57	58
A3	Early latent syphilis (first 2 years)	Male (all)	14	12	28	30	36	31	39	44	27	33
		-of which MSM	4	5	9	13	14	9	18	16	9	17
		Female	4	6	10	12	8	14	9	11	9	10
		Total	18	18	38	42	44	45	48	55	36	43
A4,A5,A6	Other acquired syphilis	Male (all)	49	56	66	60	86	64	69	54	63	41
		-of which MSM	2	7	7	8	7	10	6	12	13	8
		Female	28	58	49	63	64	56	51	59	45	37
		Total	77	114	115	123	150	120	120	113	108	78
A7	Congenital syphilis, aged under 2	Male (all)	0	0	0	1	2	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	1	0	0	0	0	0	0
		Total	0	0	0	2	2	0	0	0	0	0
A8	Congenital syphilis, aged 2 or over	Male (all)	0	0	0	1	1	0	0	0	1	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	1	0	0	2	1	0	1	1	0
		Total	0	1	0	1	3	1	0	1	2	0
A9	Epidemiological treatment of suspected syphilis	Male (all)	2	6	5	9	7	11	9	18	10	7
		-of which MSM	2	2	2	2	1	2	6	6	6	2
		Female	3	1	2	4	5	7	4	4	3	3
		Total	5	7	7	13	12	18	13	22	13	10
B1,B2	Uncomplicated gonorrhoea	Male (all)	640	650	810	783	644	557	536	512	605	579
		-of which MSM	107	98	130	138	126	141	116	95	129	191
		Female	355	385	433	432	298	277	258	305	298	266
		Total	995	1,035	1,243	1,215	942	834	794	817	903	845
B3	Gonococcal ophthalmia neonatorum	Male (all)	0	0	0	0	0	0	2	3	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	0	0	0	0	0	2	1	0	0
		Total	1	0	0	0	0	0	4	4	0	0
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	96	132	153	192	198	208	197	213	213	214
		-of which MSM	21	23	35	27	43	56	43	45	57	62
		Female	155	166	202	209	255	174	175	188	171	146
		Total	251	298	355	401	453	382	372	401	384	360
B5	Complicated gonococcal infection - including PID and epididymitis	Male (all)	2	3	6	4	4	3	4	4	3	6
		-of which MSM	2	1	0	0	1	0	0	1	1	3
		Female	13	12	18	20	8	4	6	4	3	8
		Total	15	15	24	24	12	7	10	8	6	14

East of England SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	3	1	4	11	11	5	9	17	7	13
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3	3	8	7	5	9	6	7	13	12
		Total	6	4	12	18	16	14	15	24	20	25
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	2,048	2,363	2,719	3,337	3,481	3,724	4,494	4,776	4,100	3,656
		-of which MSM	30	48	44	72	61	131	126	154	171	201
		Female	3,442	4,059	4,407	4,636	4,828	4,520	5,172	5,233	4,162	3,672
		Total	5,490	6,422	7,126	7,973	8,309	8,244	9,666	10,009	8,262	7,328
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	30	43	36	41	49	48	46	63	61	50
		-of which MSM	0	2	1	1	0	1	4	9	5	2
		Female	156	164	139	172	178	147	169	170	188	188
		Total	186	207	175	213	227	195	215	233	249	238
C4D	Chlamydial ophthalmia neonatorum	Male (all)	0	1	1	1	4	0	1	2	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	1	1	3	4	1	1	2	1
		Total	0	1	2	2	7	4	2	3	2	1
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	1,363	1,559	1,658	1,852	2,007	1,881	1,909	1,867	1,789	1,674
		-of which MSM	20	23	24	31	42	67	61	78	87	96
		Female	928	922	920	1,284	1,381	1,348	1,405	1,414	1,489	1,206
		Total	2,291	2,481	2,578	3,136	3,388	3,229	3,314	3,281	3,278	2,880
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	5,351	6,256	6,727	6,843	6,807	7,296	7,505	7,428	7,186	6,444
		-of which MSM	194	275	285	322	271	289	311	336	405	353
		Female	0	0	1,636	1,551	1,133	930	761	630	621	660
		Total	5,351	6,256	8,363	8,394	7,940	8,226	8,266	8,058	7,807	7,104
C4I	Epidemiological treatment of NSGI	Male (all)	693	644	728	793	735	760	790	775	704	692
		-of which MSM	19	40	47	41	45	44	45	51	63	64
		Female	2,581	2,630	2,084	2,151	2,164	2,293	2,507	2,546	2,227	1,975
		Total	3,274	3,274	2,812	2,944	2,899	3,053	3,297	3,321	2,931	2,667
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	331	381	396	367	455	461	436	498	522	496
		-of which MSM	9	19	8	15	9	11	11	20	21	23
		Female	785	824	869	922	819	904	1,010	1,177	1,240	1,336
		Total	1,116	1,205	1,265	1,289	1,274	1,365	1,446	1,675	1,762	1,832
C6A	Trichomoniasis	Male (all)	7	12	7	10	16	16	13	15	17	13
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	258	297	258	292	315	283	279	303	296	332
		Total	265	309	265	302	331	299	292	318	313	345
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	132	88	82	89	72	85	125	119	111	119
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	4,770	5,460	5,371	5,387	5,306	5,567	5,953	6,512	6,502	6,663
		Total	4,902	5,548	5,453	5,476	5,378	5,652	6,078	6,631	6,613	6,782
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	896	895	902	778	817	899	930	842	811	724
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	354	534	506	428	449	251	216	243	241	204
		Total	1,250	1,429	1,408	1,206	1,266	1,150	1,146	1,085	1,052	928

East of England SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C7A	Anogenital candidosis	Male (all)	638	635	568	483	414	498	607	719	828	930	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	7,136	7,648	6,649	6,488	6,222	6,174	6,336	6,805	6,732	6,547	
		Total	7,774	8,283	7,217	6,971	6,636	6,672	6,943	7,524	7,560	7,477	
C7B	Epidemiological treatment of C6 & C7	Male (all)	398	410	351	437	533	415	496	487	368	359	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	516	480	448	443	607	643	593	763	575	534	
		Total	914	890	799	880	1,140	1,058	1,089	1,250	943	893	
C8,C9	Scabies/pediculosis pubis	Male (all)	299	255	208	156	121	141	106	92	104	112	
		-of which MSM	49	36	35	25	23	37	28	18	14	35	
		Female	73	58	36	18	21	17	12	13	12	10	
		Total	372	313	244	174	142	158	118	105	116	122	
C10A	Anogenital herpes simplex – first episode	Male (all)	595	577	561	554	556	638	606	721	838	984	
		-of which MSM	14	21	14	18	22	19	12	34	25	38	
		Female	1,073	1,038	951	955	1,014	1,086	1,268	1,365	1,433	1,656	
		Total	1,668	1,615	1,512	1,509	1,570	1,724	1,874	2,086	2,271	2,640	
C10B	Anogenital herpes simplex – recurrence	Male (all)	606	661	688	665	615	585	633	599	692	823	
		-of which MSM	24	23	22	33	35	22	30	32	22	19	
		Female	950	979	1,002	813	800	732	884	927	1,112	1,185	
		Total	1,556	1,640	1,690	1,478	1,415	1,317	1,517	1,526	1,804	2,008	
C11A	Anogenital warts – first episode	Male (all)	3,085	3,109	3,244	3,369	3,478	3,464	3,670	3,813	3,913	3,771	
		-of which MSM	81	97	101	129	126	116	109	109	134	142	
		Female	2,878	2,864	2,932	3,179	3,277	3,186	3,432	3,504	3,535	3,355	
		Total	5,963	5,973	6,176	6,548	6,755	6,650	7,102	7,317	7,448	7,126	
C11B	Anogenital warts – recurrence	Male (all)	2,172	2,078	2,116	2,143	2,100	2,302	2,378	2,547	2,752	2,910	
		-of which MSM	61	61	70	82	68	85	68	78	110	102	
		Female	1,266	1,261	1,293	1,316	1,376	1,581	1,550	1,758	1,828	1,841	
		Total	3,438	3,339	3,409	3,459	3,476	3,883	3,928	4,305	4,580	4,751	
C11C	Anogenital warts – re-registered cases	Male (all)	727	729	651	602	606	628	515	530	474	441	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	573	529	509	461	520	508	433	409	403	321	
		Total	1,300	1,258	1,160	1,063	1,126	1,136	948	939	877	762	
C12	Molluscum contagiosum	Male (all)	560	641	670	750	837	963	926	986	919	882	
		-of which MSM	3	17	9	8	15	10	14	22	17	12	
		Female	296	317	352	404	436	455	495	526	409	383	
		Total	856	958	1,022	1,154	1,273	1,418	1,421	1,512	1,328	1,265	
C13	Antigen positive viral hepatitis B	Male (all)	59	51	-	-	-	-	-	-	-	-	
		-of which MSM	7	8	-	-	-	-	-	-	-	-	
		Female	22	21	-	-	-	-	-	-	-	-	
		Total	81	72	-	-	-	-	-	-	-	-	
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	34	32	50	44	48	37	39	48	
		-of which MSM	-	-	5	2	2	1	2	1	3	4	
		Female	-	-	32	29	34	34	26	37	41	38	
		Total	-	-	66	61	84	78	74	74	80	86	

East of England SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	5	2	5	4	1	2	5	2
		-of which MSM	-	-	0	0	1	1	0	1	0	0
		Female	-	-	0	3	4	3	1	2	5	3
		Total	-	-	5	5	9	7	2	4	10	5
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	3	8	13	14	20	26	37	36
		-of which MSM	-	-	0	5	7	6	3	5	8	8
		Female	-	-	2	9	5	2	12	19	17	30
		Total	-	-	5	17	18	16	32	45	54	66
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	102	76	79	73	55	45	45	54	56	38
		-of which MSM	4	3	2	3	2	1	4	1	8	3
		Female	42	27	40	35	30	35	32	27	32	19
		Total	144	103	119	108	85	80	77	81	88	57
E1A	New HIV diagnosis - asymptomatic	Male (all)	89	110	152	139	145	162	177	155	109	151
		-of which MSM	19	27	34	41	35	34	44	44	45	61
		Female	111	172	202	182	188	192	152	150	134	106
		Total	200	282	354	321	333	354	329	305	243	257
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	377	611	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	238	596	-	-	-	-	-	-	-	-
		Total	615	1,207	-	-	-	-	-	-	-	-
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	1,153	1,435	1,861	2,063	1,799	1,869	1,897	1,933
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	1,127	1,520	2,042	2,347	2,337	2,219	2,172	2,233
		Total	-	-	2,280	2,955	3,903	4,410	4,136	4,088	4,069	4,166
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	30	54	53	72	72	58	51	58	38	55
		-of which MSM	5	10	10	15	14	13	7	14	9	16
		Female	36	48	46	83	74	56	49	45	33	30
		Total	66	102	99	155	146	114	100	103	71	85
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	355	501	-	-	-	-	-	-	-	-
		-of which MSM	0	0	-	-	-	-	-	-	-	-
		Female	115	277	-	-	-	-	-	-	-	-
		Total	470	778	-	-	-	-	-	-	-	-
E3A*	AIDS – first presentation	Male (all)	29	30	-	-	-	-	-	-	-	-
		-of which MSM	3	4	-	-	-	-	-	-	-	-
		Female	17	25	-	-	-	-	-	-	-	-
		Total	46	55	-	-	-	-	-	-	-	-
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	11	25	21	20	19	24	13	18
		-of which MSM	-	-	3	6	5	2	3	3	0	9
		Female	-	-	13	16	22	9	11	16	11	8
		Total	-	-	24	41	43	29	30	40	24	26
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	7	10	14	12	7	7	2	7
		-of which MSM	-	-	1	2	3	3	0	2	0	2
		Female	-	-	5	14	12	7	7	11	8	2
		Total	-	-	12	24	26	19	14	18	10	9

East of England SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	263	327	409	455	575	675	519	559	529	484	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	61	140	195	269	374	440	404	409	397	413	413
		Total	324	467	604	724	949	1,115	923	968	926	897	897
D2A	Urinary tract infection	Male (all)	143	150	139	95	102	121	116	105	153	164	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	749	770	655	650	647	780	838	927	996	1,024	1,024
		Total	892	920	794	745	749	901	954	1,032	1,149	1,188	1,188
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	3,627	3,624	4,067	4,417	4,892	5,875	5,860	6,542	6,904	7,090	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	4,923	4,892	4,180	4,710	5,154	6,125	6,090	7,072	7,006	7,308	7,308
		Total	8,550	8,516	8,247	9,127	10,046	12,000	11,950	13,614	13,910	14,398	14,398
D3	Other episodes not requiring treatment	Male (all)	11,095	12,758	12,880	16,009	18,472	19,036	20,388	25,066	25,431	24,937	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	12,359	13,755	14,171	17,155	20,425	21,486	24,068	29,500	29,693	29,506	29,506
		Total	23,454	26,513	27,051	33,164	38,897	40,522	44,456	54,566	55,124	54,443	54,443
P1A	HIV antibody test (no sexual health screen)	Male (all)	9,303	11,972	3,428	2,684	2,682	2,391	2,665	2,339	2,161	2,133	
		-of which MSM	774	961	343	326	295	244	254	252	311	298	
		Female	7,861	10,918	2,915	2,198	2,314	2,198	2,470	2,230	2,093	1,963	
		Total	17,164	22,890	6,343	4,882	4,996	4,589	5,135	4,569	4,254	4,096	4,096
P1B	HIV antibody test offered and refused	Male (all)	1,438	3,512	8,881	11,606	11,408	11,118	11,297	11,527	11,447	11,005	
		-of which MSM	63	111	302	267	271	257	271	251	333	306	
		Female	1,780	4,312	10,817	13,626	13,694	13,934	14,945	16,030	15,855	15,386	
		Total	3,218	7,824	19,698	25,232	25,102	25,052	26,242	27,557	27,302	26,391	26,391
P2	Hepatitis B vaccination (1st dose only)	Male (all)	966	998	1,156	1,130	986	1,171	1,113	1,347	1,321	1,365	
		-of which MSM	372	457	557	506	425	472	511	602	615	649	
		Female	352	435	495	486	479	521	558	663	636	622	
		Total	1,318	1,433	1,651	1,616	1,465	1,692	1,671	2,010	1,957	1,987	1,987
P3	Contraception (excluding condom provision)	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	4,508	4,448	3,628	2,923	2,855	2,935	2,682	3,239	4,021	4,779	
		Total	4,508	4,448	3,628	2,923	2,855	2,935	2,682	3,239	4,021	4,779	4,779
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	391	340	333	253	205	195	188	235	270	239	
		Total	391	340	333	253	205	195	188	235	270	239	239

East of England SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	92	83	97	76	65	46	44	30	42	28
		Total	92	83	97	76	65	46	44	30	42	28
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	13,509	12,725	11,664	11,189	11,187	11,127	10,665	9,989
		-of which MSM	-	-	390	342	300	323	324	351	451	453
		Female	-	-	17,278	16,115	14,900	14,462	15,355	16,113	15,214	14,480
		Total	-	-	30,787	28,840	26,564	25,651	26,542	27,240	25,879	24,469
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	14,954	20,203	23,951	27,318	31,701	36,434	36,305	35,750
		-of which MSM	-	-	922	1,115	1,134	1,357	1,512	1,850	2,299	2,664
		Female	-	-	15,974	21,553	26,145	28,758	34,439	39,769	39,095	37,841
		Total	-	-	30,928	41,756	50,096	56,076	66,140	76,203	75,400	73,591
Total new STI diagnoses (for GUM clinics only)		Male (all)	13,100	14,494	15,696	16,561	16,802	17,658	18,696	19,260	18,511	17,313
		-of which MSM	526	668	711	826	750	843	827	892	1,011	1,130
		Female	9,486	10,256	12,324	12,899	12,637	12,098	13,099	13,476	12,405	12,040
		Total	22,586	24,750	28,020	29,460	29,439	29,756	31,795	32,736	30,916	29,353
Total other STI diagnoses		Male (all)	6,732	7,335	7,635	8,219	8,811	9,189	8,828	9,043	9,126	9,226
		-of which MSM	152	183	208	226	244	289	259	304	358	355
		Female	6,916	7,585	7,389	8,106	8,998	9,498	9,760	9,947	9,858	9,363
		Total	13,648	14,920	15,024	16,325	17,809	18,687	18,588	18,990	18,984	18,589
Total other GUM clinic diagnoses		Male (all)	5,995	5,929	6,230	6,414	6,953	8,000	8,248	8,933	9,312	9,510
		-of which MSM	11	11	7	10	12	9	9	8	19	15
		Female	18,995	20,255	18,313	18,511	18,728	19,855	20,329	22,672	22,459	22,637
		Total	24,990	26,184	24,543	24,925	25,681	27,855	28,577	31,605	31,771	32,147
Total services provided		Male (all)	22,802	29,240	54,808	64,357	69,163	72,223	78,351	87,840	87,330	85,179
		-of which MSM	1,209	1,529	2,514	2,556	2,425	2,653	2,872	3,306	4,009	4,370
		Female	26,860	33,868	65,278	74,056	80,812	84,294	94,517	107,544	106,607	104,577
		Total	49,662	63,108	120,086	138,413	149,975	156,517	172,868	195,384	193,937	189,756

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 d: London SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	271	514	566	669	692	667	790	594	765	735
		-of which MSM	160	292	331	362	481	445	530	330	575	501
		Female	46	67	93	100	111	72	77	52	52	61
		Total	317	581	659	769	803	739	867	646	826	811
A3	Early latent syphilis (first 2 years)	Male (all)	96	156	165	197	215	236	303	260	267	297
		-of which MSM	47	77	65	96	138	152	193	123	196	204
		Female	37	61	72	67	46	50	45	47	47	36
		Total	133	217	237	264	261	286	348	307	314	333
A4,A5,A6	Other acquired syphilis	Male (all)	504	493	542	622	508	548	529	489	479	546
		-of which MSM	75	62	85	81	85	109	105	114	127	151
		Female	415	452	484	466	404	369	431	329	353	336
		Total	919	945	1,026	1,088	912	917	960	818	833	891
A7	Congenital syphilis, aged under 2	Male (all)	0	2	2	2	2	1	1	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	5	0	1	3	0	1	1	0	0
Total	0	7	2	3	5	1	2	1	0	0		
A8	Congenital syphilis, aged 2 or over	Male (all)	1	4	0	2	1	1	1	2	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3	2	0	1	5	1	2	0	3	0
Total	4	6	0	3	6	2	3	2	3	0		
A9	Epidemiological treatment of suspected syphilis	Male (all)	24	105	186	252	271	335	420	453	516	547
		-of which MSM	7	51	111	149	188	230	288	235	381	356
		Female	17	38	37	39	40	51	62	61	51	66
		Total	41	143	223	291	311	386	482	514	569	615
B1,B2	Uncomplicated gonorrhoea	Male (all)	6,507	6,841	5,982	5,375	4,890	4,700	4,681	3,905	4,450	4,973
		-of which MSM	1,777	1,564	1,701	1,850	2,097	2,098	1,864	1,275	1,947	2,445
		Female	2,597	2,885	2,503	1,987	1,518	1,489	1,767	1,666	1,580	1,630
		Total	9,104	9,726	8,485	7,362	6,408	6,189	6,448	5,571	6,052	6,632
B3	Gonococcal ophthalmia neonatorum	Male (all)	2	2	3	5	5	9	4	5	11	8
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	6	10	6	5	5	5	7	9	6
Total	3	8	13	11	10	14	9	12	20	14		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	1,283	1,367	1,457	1,551	1,490	1,512	1,871	1,822	1,880	2,182
		-of which MSM	501	473	594	661	683	748	804	595	902	985
		Female	1,095	1,278	1,216	1,129	1,090	1,001	1,296	1,356	1,025	874
		Total	2,378	2,645	2,673	2,680	2,580	2,513	3,167	3,178	2,916	3,069
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	44	37	30	31	36	25	26	34	49	76
		-of which MSM	5	8	4	8	16	10	9	9	14	45
		Female	107	99	97	64	69	58	84	75	96	73
		Total	151	136	127	95	105	83	110	109	145	149

London SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	14	14	27	37	173	139	180	145	153	264
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	19	16	28	20	23	15	54	25	11	32
		Total	33	30	55	57	196	154	234	170	166	297
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	8,292	9,110	9,567	9,834	10,323	10,262	11,297	11,112	11,003	11,818
		-of which MSM	760	681	770	755	1,052	1,282	1,425	1,531	1,908	2,479
		Female	8,913	9,504	9,915	9,999	10,009	9,582	10,587	10,181	9,682	9,712
		Total	17,205	18,614	19,482	19,833	20,332	19,844	21,884	21,293	20,776	21,624
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	180	147	179	145	159	156	176	196	220	178
		-of which MSM	23	21	26	26	27	38	35	42	57	63
		Female	526	495	461	426	450	377	471	463	410	414
		Total	706	642	640	571	609	533	647	659	632	595
C4D	Chlamydial ophthalmia neonatorum	Male (all)	4	5	7	7	6	3	4	3	2	2
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	2	6	8	5	8	5	2	2	4	1
		Total	6	11	15	12	14	8	6	5	6	3
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	4,189	5,879	6,411	6,575	6,308	6,300	7,517	7,255	7,613	7,794
		-of which MSM	314	581	747	783	895	1,046	1,271	1,218	1,432	1,558
		Female	3,902	4,678	4,859	4,974	4,974	4,764	5,506	5,440	5,004	4,835
		Total	8,091	10,557	11,270	11,549	11,282	11,064	13,023	12,695	12,654	12,670
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	23,439	23,144	24,292	25,289	25,065	25,156	24,643	19,881	18,914	18,304
		-of which MSM	2,675	2,368	2,668	2,746	2,836	3,314	3,121	2,666	2,614	2,501
		Female	0	0	1,111	838	902	929	934	882	958	937
		Total	23,439	23,144	25,403	26,127	25,967	26,085	25,577	20,763	19,938	19,331
C4I	Epidemiological treatment of NSGI	Male (all)	2,094	2,057	1,993	2,216	1,947	2,069	1,935	1,794	1,846	1,808
		-of which MSM	371	305	391	411	391	496	443	299	330	322
		Female	6,468	5,410	5,189	5,429	5,393	4,865	4,716	3,493	3,280	3,054
		Total	8,562	7,467	7,182	7,645	7,340	6,934	6,651	5,287	5,142	4,866
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	1,242	1,163	1,217	1,534	1,572	1,487	1,581	1,730	1,992	2,196
		-of which MSM	144	111	137	150	200	180	199	262	305	281
		Female	4,913	4,981	4,653	5,400	4,867	4,968	5,567	5,496	5,923	5,395
		Total	6,155	6,144	5,870	6,934	6,439	6,455	7,148	7,226	7,937	7,623
C6A	Trichomoniasis	Male (all)	544	363	162	149	176	132	156	157	181	173
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3,069	3,320	3,097	2,978	2,285	2,358	2,591	2,418	1,800	1,765
		Total	3,613	3,683	3,259	3,127	2,461	2,490	2,747	2,575	1,992	1,943
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	197	248	273	229	249	253	238	256	264	259
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	24,750	26,823	26,218	24,905	25,309	26,653	31,189	30,976	32,810	32,981
		Total	24,947	27,071	26,491	25,134	25,558	26,906	31,427	31,232	33,300	33,477
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	2,521	2,550	2,858	2,680	2,687	2,675	3,210	3,144	3,257	3,248
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1,771	1,434	1,384	1,758	1,129	1,266	1,215	1,318	1,624	1,290
		Total	4,292	3,984	4,242	4,438	3,816	3,941	4,425	4,462	4,949	4,582

London SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C7A	Anogenital candidosis	Male (all)	2,465	2,377	2,085	2,124	1,729	1,969	2,434	2,509	2,904	2,951	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	23,285	22,093	20,623	20,000	17,866	17,706	20,489	21,518	22,817	22,817	23,802
		Total	25,750	24,470	22,708	22,124	19,595	19,675	22,923	24,027	25,853	26,899	
C7B	Epidemiological treatment of C6 & C7	Male (all)	1,340	1,554	1,544	1,475	1,489	1,357	1,368	1,418	1,533	1,180	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	2,230	2,173	2,393	3,009	2,639	2,769	3,001	3,157	2,991	2,614	
		Total	3,570	3,727	3,937	4,484	4,128	4,126	4,369	4,575	4,536	3,806	
C8,C9	Scabies/pediculosis pubis	Male (all)	1,259	1,093	1,009	798	591	591	531	519	517	491	
		-of which MSM	359	300	280	230	197	183	175	166	226	185	
		Female	280	217	178	147	108	79	78	67	64	59	
		Total	1,539	1,310	1,187	945	699	670	609	586	581	552	
C10A	Anogenital herpes simplex – first episode	Male (all)	1,887	1,945	2,007	1,912	1,946	2,045	2,391	2,447	2,726	2,856	
		-of which MSM	238	272	279	254	293	280	358	365	472	514	
		Female	2,729	2,772	2,853	2,762	2,762	3,018	3,629	3,669	3,907	4,177	
		Total	4,616	4,717	4,860	4,674	4,708	5,063	6,020	6,116	6,665	7,060	
C10B	Anogenital herpes simplex – recurrence	Male (all)	1,725	1,729	1,767	1,748	1,716	1,733	1,910	2,005	2,362	2,470	
		-of which MSM	189	198	234	195	211	232	293	311	383	398	
		Female	2,056	1,963	1,891	1,980	2,185	2,160	2,450	2,395	2,848	3,097	
		Total	3,781	3,692	3,658	3,728	3,901	3,893	4,360	4,400	5,257	5,634	
C11A	Anogenital warts – first episode	Male (all)	6,457	7,007	6,807	7,131	6,785	6,964	7,609	7,567	7,792	7,615	
		-of which MSM	865	960	959	887	967	1,054	1,141	1,148	1,102	1,001	
		Female	5,352	5,529	5,348	5,504	5,462	5,605	6,086	6,080	5,991	5,849	
		Total	11,809	12,536	12,155	12,635	12,247	12,569	13,695	13,647	13,829	13,531	
C11B	Anogenital warts – recurrence	Male (all)	4,800	4,760	4,828	4,935	4,969	5,193	5,813	5,571	6,762	6,911	
		-of which MSM	711	723	788	857	775	830	1,062	898	1,000	952	
		Female	2,273	2,297	2,441	2,523	2,622	2,631	3,040	3,060	3,451	3,519	
		Total	7,073	7,057	7,269	7,458	7,591	7,824	8,853	8,631	10,270	10,528	
C11C	Anogenital warts – re-registered cases	Male (all)	1,908	1,802	1,811	1,761	1,743	1,857	1,967	2,027	2,026	1,870	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	1,098	1,063	1,051	968	1,032	1,018	1,008	1,049	972	956	
		Total	3,006	2,865	2,862	2,729	2,775	2,875	2,975	3,076	3,012	2,849	
C12	Molluscum contagiosum	Male (all)	1,289	1,396	1,602	1,806	1,923	1,999	2,239	2,276	2,106	1,914	
		-of which MSM	109	103	118	119	118	153	158	226	136	121	
		Female	679	734	757	835	823	905	1,019	954	865	807	
		Total	1,968	2,130	2,359	2,641	2,746	2,904	3,258	3,230	2,984	2,738	
C13	Antigen positive viral hepatitis B	Male (all)	403	576	-	-	-	-	-	-	-	-	
		-of which MSM	94	115	-	-	-	-	-	-	-	-	
		Female	231	308	-	-	-	-	-	-	-	-	
		Total	634	884	-	-	-	-	-	-	-	-	
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	305	358	300	316	339	413	449	419	
		-of which MSM	-	-	60	53	47	28	28	47	47	55	
		Female	-	-	164	190	176	198	210	217	234	204	
		Total	-	-	469	548	476	514	549	630	684	624	

London SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	22	32	21	15	17	20	24	39
		-of which MSM	-	-	8	14	9	3	1	7	12	13
		Female	-	-	5	7	13	9	6	5	14	10
		Total	-	-	27	39	34	24	23	25	38	49
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	237	351	271	266	287	304	354	300
		-of which MSM	-	-	76	167	152	164	82	122	87	52
		Female	-	-	112	126	86	124	124	136	192	204
		Total	-	-	349	477	357	390	411	440	550	504
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	210	288	229	295	229	239	175	365	577	292
		-of which MSM	51	76	53	62	58	54	37	178	344	109
		Female	89	147	134	94	93	82	57	88	139	76
		Total	299	435	363	389	322	321	232	453	717	368
E1A	New HIV diagnosis - asymptomatic	Male (all)	815	1,124	1,135	1,232	1,182	1,054	1,487	1,242	1,333	1,343
		-of which MSM	384	341	544	595	568	521	662	545	871	794
		Female	534	960	726	687	691	581	949	593	553	520
		Total	1,349	2,084	1,861	1,919	1,873	1,635	2,436	1,835	1,888	1,866
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	4,575	4,792	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	2,365	2,839	-	-	-	-	-	-	-	-
		Total	6,940	7,631	-	-	-	-	-	-	-	-
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	9,927	10,894	12,077	11,283	12,110	13,286	13,967	16,037
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	4,498	5,295	6,059	5,747	6,415	6,911	7,063	7,531
		Total	-	-	14,425	16,189	18,136	17,030	18,525	20,197	21,113	23,695
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	268	380	306	278	240	300	175	209	242	218
		-of which MSM	135	131	144	113	93	117	67	68	126	87
		Female	179	212	197	126	141	126	98	93	116	124
		Total	447	592	503	404	381	426	273	302	358	343
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	4,876	4,580	-	-	-	-	-	-	-	-
		-of which MSM	0	0	-	-	-	-	-	-	-	-
		Female	1,734	1,958	-	-	-	-	-	-	-	-
		Total	6,610	6,538	-	-	-	-	-	-	-	-
E3A*	AIDS – first presentation	Male (all)	210	215	-	-	-	-	-	-	-	-
		-of which MSM	86	84	-	-	-	-	-	-	-	-
		Female	135	122	-	-	-	-	-	-	-	-
		Total	345	337	-	-	-	-	-	-	-	-
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	78	63	68	57	70	36	95	92
		-of which MSM	-	-	40	19	26	21	23	5	32	24
		Female	-	-	47	40	37	31	36	34	50	35
		Total	-	-	125	103	105	88	106	70	145	127
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	189	74	79	71	55	61	36	27
		-of which MSM	-	-	78	42	31	40	28	31	24	11
		Female	-	-	109	35	25	30	27	21	9	17
		Total	-	-	298	109	104	101	82	82	45	44

London SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	2,253	2,774	3,735	3,797	4,086	3,582	3,613	3,722	3,368	3,474	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	745	1,103	1,828	1,760	1,922	1,812	1,863	1,984	1,817	1,877	
		Total	2,998	3,877	5,563	5,557	6,008	5,394	5,476	5,706	5,186	5,351	
D2A	Urinary tract infection	Male (all)	833	1,355	1,499	1,095	1,160	999	1,130	1,153	1,200	1,165	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	4,647	4,475	4,418	4,865	4,762	4,626	5,687	6,189	6,267	6,124	
		Total	5,480	5,830	5,917	5,960	5,922	5,625	6,817	7,342	7,485	7,321	
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	18,290	19,796	21,810	22,103	21,997	21,650	27,378	29,915	34,326	39,073	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	16,145	16,767	16,569	16,921	16,420	16,147	19,778	21,095	28,941	31,336	
		Total	34,435	36,563	38,379	39,024	38,417	37,797	47,156	51,010	63,464	70,627	
D3	Other episodes not requiring treatment	Male (all)	46,905	50,015	45,725	51,910	56,528	62,294	65,119	69,653	80,830	105,721	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	50,269	52,273	44,115	49,725	55,724	60,451	66,127	64,788	72,725	89,876	
		Total	97,174	102,288	89,840	101,635	112,252	122,745	131,246	134,441	154,200	196,355	
P1A	HIV antibody counselling - with testing	Male (all)	40,085	49,836	20,975	18,801	16,687	12,837	10,888	10,166	9,258	12,609	
		-of which MSM	8,408	9,387	4,690	4,267	4,404	3,987	3,857	1,820	4,567	6,660	
		Female	34,693	46,519	17,178	13,155	12,087	8,702	6,962	6,133	4,509	4,649	
		Total	74,778	96,355	38,153	31,956	28,774	21,539	17,850	16,299	13,791	17,278	
P1B	HIV antibody counselling - without testing	Male (all)	5,397	12,431	30,327	38,611	32,046	28,165	26,614	24,795	25,022	25,273	
		-of which MSM	881	1,730	3,328	3,583	3,582	3,483	3,423	3,705	3,876	3,467	
		Female	4,848	14,175	35,996	46,318	38,796	34,193	33,426	31,445	32,741	34,004	
		Total	10,245	26,606	66,323	84,929	70,842	62,358	60,040	56,240	58,013	59,390	
P2	Hepatitis B vaccination	Male (all)	4,731	4,798	4,525	5,905	5,413	5,118	5,149	5,427	5,573	5,871	
		-of which MSM	2,737	2,844	2,951	3,695	3,520	3,282	3,162	2,576	3,624	3,543	
		Female	965	1,312	1,195	1,688	1,647	1,734	1,813	2,013	1,966	1,874	
		Total	5,696	6,110	5,720	7,593	7,060	6,852	6,962	7,440	7,559	7,773	
P3	Family Planning	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	18,068	16,774	16,395	19,825	20,080	22,249	26,572	26,849	34,848	41,364	
		Total	18,068	16,774	16,395	19,825	20,080	22,249	26,572	26,849	34,848	41,364	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	1,361	940	2,579	838	405	372	363	297	490	548	
		Total	1,361	940	2,579	838	405	372	363	297	490	548	

London SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	259	165	293	167	59	65	39	54	59	48
		Total	259	165	293	167	59	65	39	54	59	48
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	39,625	41,286	39,333	39,112	39,741	38,176	38,631	41,114
		-of which MSM	-	-	5,621	5,924	6,590	7,319	7,371	7,276	9,321	10,687
		Female	-	-	50,465	51,855	49,836	48,103	49,485	46,282	46,520	47,185
		Total	-	-	90,090	93,141	89,169	87,215	89,226	84,458	85,613	88,744
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	45,962	57,552	73,381	84,711	101,487	113,750	127,704	135,754
		-of which MSM	-	-	7,524	8,724	11,166	13,675	15,283	18,122	22,492	23,244
		Female	-	-	48,143	60,172	77,031	86,952	106,442	120,171	135,113	141,407
		Total	-	-	94,105	117,724	150,412	171,663	207,929	233,921	263,866	278,520
Total new STI diagnoses (for GUM clinics only)	Male (all)	52,604	54,434	55,131	56,480	56,036	55,970	58,335	52,310	52,805	53,543	
	-of which MSM	7,681	7,229	8,066	8,210	9,109	9,848	9,960	8,761	10,581	11,245	
	Female	29,980	31,852	32,136	31,980	30,304	30,243	34,072	32,795	32,105	31,626	
	Total	82,584	86,286	87,267	88,460	86,340	86,213	92,407	85,105	85,228	85,555	
Total other STI diagnoses	Male (all)	28,448	30,566	32,858	34,441	35,208	34,497	37,750	38,495	40,868	43,676	
	-of which MSM	2,254	2,477	3,028	3,179	3,259	3,731	4,294	3,701	4,579	4,733	
	Female	22,309	23,220	23,621	24,611	25,767	24,459	26,824	26,109	25,889	26,169	
	Total	50,757	53,786	56,479	59,052	60,975	58,956	64,574	64,604	67,026	70,229	
Total other GUM clinic diagnoses	Male (all)	26,259	28,744	30,862	30,742	30,132	29,739	36,576	39,497	44,888	48,926	
	-of which MSM	145	191	197	296	266	249	148	354	490	229	
	Female	74,768	75,325	74,892	72,880	68,957	70,017	82,158	85,050	96,578	99,237	
	Total	101,027	104,069	105,754	103,622	99,089	99,756	118,734	124,547	142,125	148,853	
Total services provided	Male (all)	97,118	117,080	187,139	214,065	223,388	232,237	248,998	261,967	287,018	326,342	
	-of which MSM	12,026	13,961	24,114	26,193	29,262	31,746	33,096	33,499	43,880	47,601	
	Female	108,843	131,053	213,487	242,738	255,201	262,384	290,827	297,681	328,422	360,359	
	Total	205,961	248,133	400,626	456,803	478,589	494,621	539,825	559,648	617,890	689,424	

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 e: North East SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	2	17	63	88	74	72	104	82	86	58
		-of which MSM	1	15	45	65	56	39	66	49	60	41
		Female	2	0	4	4	6	15	17	9	19	11
		Total	4	17	67	92	80	87	121	91	105	69
A3	Early latent syphilis (first 2 years)	Male (all)	5	8	21	24	36	38	28	24	27	23
		-of which MSM	3	6	16	20	30	32	22	20	18	20
		Female	3	0	4	2	3	3	4	1	4	10
		Total	8	8	25	26	39	41	32	25	31	33
A4,A5,A6	Other acquired syphilis	Male (all)	10	8	18	10	15	19	24	29	22	37
		-of which MSM	1	3	2	4	5	9	11	14	10	21
		Female	5	6	11	12	9	8	13	16	5	14
		Total	15	14	29	22	24	27	37	45	27	51
A7	Congenital syphilis, aged under 2	Male (all)	0	0	0	0	0	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	0	0	0	0	0
Total	0	0	0	0	0	0	0	0	0	0		
A8	Congenital syphilis, aged 2 or over	Male (all)	1	0	0	0	0	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	3	1	0	0	1	0	0	0	0
Total	1	3	1	0	0	1	0	0	0	0		
A9	Epidemiological treatment of suspected syphilis	Male (all)	1	1	3	5	8	21	20	8	31	25
		-of which MSM	0	0	1	5	6	13	10	5	15	7
		Female	0	0	0	0	1	5	4	3	8	1
		Total	1	1	3	5	9	26	24	11	39	26
B1,B2	Uncomplicated gonorrhoea	Male (all)	371	525	448	423	415	348	320	327	390	367
		-of which MSM	74	134	107	107	146	110	74	74	97	94
		Female	174	217	217	164	139	134	130	149	195	247
		Total	545	742	665	587	554	482	450	476	585	614
B3	Gonococcal ophthalmia neonatorum	Male (all)	0	0	1	0	0	1	0	1	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	0	0	0	0	0
Total	0	0	1	0	0	1	0	1	0	0		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	23	97	93	89	118	108	106	176	138	186
		-of which MSM	3	39	40	33	46	49	33	44	33	45
		Female	44	65	89	94	100	80	75	108	126	132
		Total	67	162	182	183	218	188	181	284	264	318
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	2	5	5	5	2	1	3	8	4	2
		-of which MSM	0	3	0	0	1	0	0	2	2	0
		Female	3	8	7	7	8	5	4	8	7	7
		Total	5	13	12	12	10	6	7	16	11	9

North East SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	1	1	1	0	1	1	1	1	0	4
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	1	3	1	0	3	0	0	0
		Total	1	1	2	3	2	1	4	1	0	4
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	850	1,026	1,419	1,973	2,197	2,660	2,892	2,968	2,782	2,785
		-of which MSM	20	22	29	72	106	132	109	133	163	218
		Female	1,496	1,727	1,984	2,440	2,462	2,682	2,600	2,757	2,611	2,648
		Total	2,346	2,753	3,403	4,413	4,659	5,342	5,492	5,725	5,393	5,433
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	19	27	26	27	38	38	34	47	33	43
		-of which MSM	1	2	1	0	0	1	0	1	1	4
		Female	87	108	98	78	95	101	107	128	114	127
		Total	106	135	124	105	133	139	141	175	147	170
C4D	Chlamydial ophthalmia neonatorum	Male (all)	1	0	0	0	0	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	1	0	2	0	0	1	0	0
		Total	1	0	1	0	2	0	0	1	0	0
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	803	959	956	1,093	1,257	1,255	1,279	1,460	1,449	1,418
		-of which MSM	10	19	19	40	57	54	56	68	94	107
		Female	388	374	474	642	765	882	979	1,032	1,054	1,041
		Total	1,191	1,333	1,430	1,735	2,022	2,137	2,258	2,492	2,503	2,459
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	1,356	1,304	1,545	1,566	1,754	1,997	2,191	1,992	2,056	1,780
		-of which MSM	53	62	81	79	100	106	117	131	144	132
		Female	0	0	262	312	346	308	179	120	127	144
		Total	1,356	1,304	1,807	1,878	2,100	2,305	2,370	2,112	2,183	1,924
C4I	Epidemiological treatment of NSGI	Male (all)	48	73	79	95	124	94	141	147	149	158
		-of which MSM	9	9	12	13	18	21	21	29	23	29
		Female	428	359	280	351	380	420	532	502	440	375
		Total	476	432	359	446	504	514	673	649	589	533
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	50	46	55	89	167	95	102	158	163	195
		-of which MSM	2	3	1	6	8	5	4	9	19	14
		Female	96	114	114	116	195	175	223	334	368	318
		Total	146	160	169	205	362	270	325	492	531	513
C6A	Trichomoniasis	Male (all)	0	2	1	1	3	2	3	5	2	3
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	68	65	42	61	51	62	67	58	54	48
		Total	68	67	43	62	54	64	70	63	56	51
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	70	43	67	64	64	85	78	47	79	73
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	2,009	2,218	2,466	2,433	2,647	2,803	3,311	3,797	4,090	4,191
		Total	2,079	2,261	2,533	2,497	2,711	2,888	3,389	3,844	4,169	4,264
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	596	637	679	628	664	646	870	949	838	773
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	323	279	197	203	208	215	221	232	218	202
		Total	919	916	876	831	872	861	1,091	1,181	1,056	975

North East SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C7A	Anogenital candidosis	Male (all)	137	104	94	102	149	227	290	231	317	244	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	1,702	1,445	1,510	1,464	1,781	1,871	2,186	2,933	3,079	2,694	
		Total	1,839	1,549	1,604	1,566	1,930	2,098	2,476	3,164	3,396	2,938	
C7B	Epidemiological treatment of C6 & C7	Male (all)	65	58	64	56	54	68	55	42	46	91	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	86	122	94	120	148	70	57	61	51	60	
		Total	151	180	158	176	202	138	112	103	97	151	
C8,C9	Scabies/pediculosis pubis	Male (all)	150	127	127	114	109	93	114	124	122	97	
		-of which MSM	29	20	32	24	30	32	40	46	40	33	
		Female	28	31	14	26	12	9	13	11	12	15	
		Total	178	158	141	140	121	102	127	135	134	112	
C10A	Anogenital herpes simplex – first episode	Male (all)	169	154	174	156	230	268	370	400	484	524	
		-of which MSM	6	8	9	6	15	9	18	22	20	21	
		Female	295	326	293	318	389	465	632	599	735	783	
		Total	464	480	467	474	619	733	1,002	999	1,219	1,307	
C10B	Anogenital herpes simplex – recurrence	Male (all)	48	51	50	48	65	66	81	126	155	174	
		-of which MSM	2	2	3	1	4	4	0	10	9	5	
		Female	96	71	52	69	73	79	112	167	237	266	
		Total	144	122	102	117	138	145	193	293	392	440	
C11A	Anogenital warts – first episode	Male (all)	1,609	1,708	1,792	2,018	2,019	2,220	2,432	2,410	2,408	2,520	
		-of which MSM	57	53	81	82	100	100	117	111	148	160	
		Female	1,549	1,631	1,597	1,795	1,849	1,959	2,293	2,255	2,206	2,134	
		Total	3,158	3,339	3,389	3,813	3,868	4,179	4,725	4,665	4,614	4,654	
C11B	Anogenital warts – recurrence	Male (all)	1,102	1,022	1,042	1,029	1,230	1,213	1,419	1,506	1,216	1,105	
		-of which MSM	22	32	44	26	52	51	58	69	49	68	
		Female	560	600	582	627	650	742	870	900	775	742	
		Total	1,662	1,622	1,624	1,656	1,880	1,955	2,289	2,406	1,991	1,847	
C11C	Anogenital warts – re-registered cases	Male (all)	557	394	421	379	370	355	302	328	300	220	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	492	384	393	406	366	341	275	328	211	176	
		Total	1,049	778	814	785	736	696	577	656	511	396	
C12	Molluscum contagiosum	Male (all)	286	279	337	397	467	529	584	561	509	481	
		-of which MSM	10	3	14	11	15	12	10	4	16	16	
		Female	164	172	183	201	272	242	272	255	203	205	
		Total	450	451	520	598	739	771	856	816	712	686	
C13	Antigen positive viral hepatitis B	Male (all)	10	11	-	-	-	-	-	-	-	-	
		-of which MSM	1	0	-	-	-	-	-	-	-	-	
		Female	3	8	-	-	-	-	-	-	-	-	
		Total	13	19	-	-	-	-	-	-	-	-	
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	8	8	7	6	16	8	16	10	
		-of which MSM	-	-	3	2	0	1	3	2	3	1	
		Female	-	-	3	3	3	1	1	0	3	4	
		Total	-	-	11	11	10	7	17	8	19	14	

North East SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	1	0	1	1	0	0	1	0	
		-of which MSM	-	-	1	0	0	0	0	0	0	0	
		Female	-	-	0	0	0	0	0	0	0	1	0
		Total	-	-	1	0	1	1	0	0	0	2	0
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	0	0	5	1	3	2	12	3	
		-of which MSM	-	-	0	0	4	0	0	0	4	1	
		Female	-	-	1	0	0	1	0	4	3	4	
		Total	-	-	1	0	5	2	3	6	15	7	
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	13	15	16	10	11	12	13	16	21	22	
		-of which MSM	2	0	4	0	0	1	1	1	3	6	
		Female	5	6	9	7	13	6	6	6	5	13	
		Total	18	21	25	17	24	18	19	22	26	35	
E1A	New HIV diagnosis - asymptomatic	Male (all)	12	22	36	38	58	41	55	31	51	43	
		-of which MSM	8	14	17	27	41	25	25	25	40	30	
		Female	10	13	24	30	20	15	20	10	14	11	
		Total	22	35	60	68	78	56	75	41	65	54	
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	103	114	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	15	51	-	-	-	-	-	-	-	-	
		Total	118	165	-	-	-	-	-	-	-	-	
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	116	214	333	391	475	473	443	509	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	-	-	71	93	110	133	188	158	148	163	
		Total	-	-	187	307	443	524	663	631	591	672	
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	1	2	4	6	3	0	8	9	3	13	
		-of which MSM	1	0	2	4	2	0	4	4	2	10	
		Female	1	2	2	0	3	2	0	3	1	1	
		Total	2	4	6	6	6	2	8	12	4	14	
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	13	8	-	-	-	-	-	-	-	-	
		-of which MSM	0	0	-	-	-	-	-	-	-	-	
		Female	4	11	-	-	-	-	-	-	-	-	
		Total	17	19	-	-	-	-	-	-	-	-	
E3A*	AIDS – first presentation	Male (all)	2	1	-	-	-	-	-	-	-	-	
		-of which MSM	1	1	-	-	-	-	-	-	-	-	
		Female	0	0	-	-	-	-	-	-	-	-	
		Total	2	1	-	-	-	-	-	-	-	-	
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	0	1	2	1	2	1	1	1	
		-of which MSM	-	-	0	0	1	0	1	0	0	1	
		Female	-	-	0	1	1	0	1	0	0	0	
		Total	-	-	0	2	3	1	3	1	1	1	
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	2	0	1	0	0	0	0	0	
		-of which MSM	-	-	1	0	1	0	0	0	0	0	
		Female	-	-	1	0	1	0	0	0	1	0	
		Total	-	-	3	0	2	0	0	0	0	1	0

North East SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	5	7	9	10	12	15	10	6	11	15	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	12	5	5	5	7	5	5	5	4	6	8
		Total	17	12	14	15	19	20	15	10	17	23	
D2A	Urinary tract infection	Male (all)	34	29	23	36	30	23	33	36	51	71	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	145	136	117	122	136	168	319	438	516	445	
		Total	179	165	140	158	166	191	352	474	567	516	
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	620	625	1,260	1,229	1,590	1,767	2,318	2,513	2,485	2,604	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	588	740	694	725	923	1,075	1,506	1,745	1,808	1,973	
		Total	1,208	1,365	1,954	1,954	2,513	2,842	3,824	4,258	4,293	4,577	
D3	Other episodes not requiring treatment	Male (all)	4,221	4,959	5,678	7,402	8,439	8,468	10,609	12,277	12,441	11,918	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	3,970	4,548	5,186	6,801	8,122	8,475	11,234	12,425	12,300	13,894	
		Total	8,191	9,507	10,864	14,203	16,561	16,943	21,843	24,702	24,741	25,813	
P1A	HIV antibody counselling - with testing	Male (all)	2,805	6,073	960	976	1,103	1,007	1,005	902	797	716	
		-of which MSM	439	606	218	170	261	235	215	177	186	175	
		Female	2,179	5,057	569	630	738	768	735	726	680	650	
		Total	4,984	11,130	1,529	1,606	1,841	1,775	1,740	1,628	1,477	1,366	
P1B	HIV antibody counselling - without testing	Male (all)	364	2,063	3,751	3,686	3,711	3,616	4,609	4,823	4,934	4,729	
		-of which MSM	119	140	166	138	159	149	164	162	166	181	
		Female	245	2,222	4,137	3,983	4,306	4,244	5,882	6,246	6,189	6,443	
		Total	609	4,285	7,888	7,669	8,017	7,860	10,491	11,069	11,123	11,172	
P2	Hepatitis B vaccination	Male (all)	253	388	485	569	531	540	585	657	772	796	
		-of which MSM	191	283	354	412	391	360	407	439	566	548	
		Female	28	101	96	119	135	134	146	183	228	283	
		Total	281	489	581	688	666	674	731	840	1,000	1,079	
P3	Family Planning	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	788	707	545	549	594	417	576	632	738	2,202	
		Total	788	707	545	549	594	417	576	632	738	2,202	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	156	126	110	66	20	28	31	43	33	53	
		Total	156	126	110	66	20	28	31	43	33	53	

North East SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	23	28	24	14	6	7	4	16	12	23
		Total	23	28	24	14	6	7	4	16	12	23
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	3,923	3,668	3,531	3,299	4,005	4,261	4,231	4,094
		-of which MSM	-	-	199	165	205	205	207	211	240	229
		Female	-	-	4,760	4,288	4,502	4,426	5,622	5,870	5,928	6,788
		Total	-	-	8,683	7,956	8,033	7,725	9,627	10,131	10,159	10,882
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	8,328	10,214	12,582	14,706	17,516	19,135	19,596	19,900
		-of which MSM	-	-	758	884	1,116	1,141	1,319	1,535	1,912	1,972
		Female	-	-	7,179	9,369	11,249	13,239	16,377	18,349	19,141	20,101
		Total	-	-	15,507	19,583	23,831	27,945	33,893	37,484	38,737	40,002
Total new STI diagnoses (for GUM clinics only)	Male (all)	4,883	5,253	6,054	6,926	7,575	8,404	9,243	9,148	9,121	8,939	
	-of which MSM	265	345	435	503	651	603	607	631	770	794	
	Female	3,976	4,414	4,846	5,558	5,852	6,177	6,565	6,697	6,670	6,709	
	Total	8,859	9,667	10,900	12,484	13,427	14,581	15,808	15,845	15,791	15,648	
Total other STI diagnoses	Male (all)	2,717	2,735	2,790	2,972	3,533	3,538	3,857	4,260	3,914	3,847	
	-of which MSM	48	105	122	122	189	201	189	239	233	282	
	Female	2,044	1,929	1,960	2,299	2,464	2,696	3,053	3,219	3,011	2,918	
	Total	4,761	4,664	4,750	5,271	5,997	6,234	6,910	7,479	6,925	6,765	
Total other GUM clinic diagnoses	Male (all)	1,545	1,522	2,212	2,133	2,575	2,836	3,676	3,844	3,866	3,891	
	-of which MSM	3	0	8	2	4	2	4	3	10	8	
	Female	5,040	5,108	5,225	5,157	5,885	6,245	7,642	9,275	9,819	9,662	
	Total	6,585	6,630	7,437	7,290	8,460	9,081	11,318	13,119	13,685	13,553	
Total services provided	Male (all)	7,643	13,483	23,125	26,515	29,897	31,636	38,329	42,055	42,771	42,153	
	-of which MSM	749	1,029	1,695	1,769	2,132	2,090	2,312	2,524	3,070	3,105	
	Female	7,210	12,635	22,472	25,739	29,646	31,703	40,572	44,431	45,204	50,361	
	Total	14,853	26,118	45,597	52,254	59,543	63,339	78,901	86,486	87,975	92,516	

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 f: North West SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	168	202	192	258	383	353	301	310	252	249
		-of which MSM	111	149	131	130	283	248	209	223	193	191
		Female	10	14	13	11	25	23	18	21	26	16
		Total	178	216	205	269	408	376	319	331	278	265
A3	Early latent syphilis (first 2 years)	Male (all)	61	58	94	82	130	118	168	172	138	125
		-of which MSM	36	42	53	52	94	93	115	122	116	98
		Female	5	7	14	23	26	24	22	17	13	12
		Total	66	65	108	105	156	142	190	189	151	137
A4,A5,A6	Other acquired syphilis	Male (all)	62	67	65	73	90	96	82	70	92	113
		-of which MSM	19	17	20	22	24	39	29	32	45	50
		Female	26	33	26	50	47	54	55	51	48	43
		Total	88	100	91	123	137	150	137	121	140	156
A7	Congenital syphilis, aged under 2	Male (all)	0	0	0	0	0	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	4	0	0	0	0	0
		Total	0	0	0	0	4	0	0	0	0	0
A8	Congenital syphilis, aged 2 or over	Male (all)	1	1	0	0	3	3	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	0	0	0	3	1	2	1	0	0
		Total	2	1	0	0	6	4	2	1	0	0
A9	Epidemiological treatment of suspected syphilis	Male (all)	12	16	20	25	41	85	83	94	139	115
		-of which MSM	11	13	13	21	27	62	65	74	116	102
		Female	0	1	4	8	8	3	15	10	25	18
		Total	12	17	24	33	49	88	98	104	164	133
B1,B2	Uncomplicated gonorrhoea	Male (all)	1,959	2,013	2,069	1,757	1,615	1,715	1,640	1,389	1,225	1,427
		-of which MSM	404	491	558	442	468	548	432	448	485	719
		Female	745	771	767	786	765	782	801	738	679	623
		Total	2,704	2,784	2,836	2,543	2,380	2,497	2,441	2,127	1,904	2,050
B3	Gonococcal ophthalmia neonatorum	Male (all)	0	1	1	1	0	2	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	2	2	0	0	0	0	1	4	2	1
		Total	2	3	1	1	0	2	1	4	2	1
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	357	391	427	425	490	582	503	637	631	624
		-of which MSM	92	111	144	135	144	177	169	192	217	226
		Female	345	325	381	338	358	408	347	395	310	317
		Total	702	716	808	763	848	990	850	1,032	941	941
B5	Complicated gonococcal infection - including PID and epididymitis	Male (all)	3	6	7	22	13	16	22	10	13	14
		-of which MSM	0	2	2	4	6	1	3	1	2	8
		Female	22	16	28	32	21	26	31	26	32	33
		Total	25	22	35	54	34	42	53	36	45	47

North West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	2	4	2	6	14	17	13	25	21	51
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	4	2	3	3	5	10	9	6	9	6
		Total	6	6	5	9	19	27	22	31	30	57
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	3,964	4,906	6,073	6,619	6,947	7,777	8,004	8,191	7,474	6,652
		-of which MSM	194	249	337	293	302	401	477	633	722	723
		Female	5,761	6,554	7,547	7,461	7,485	7,851	7,837	7,866	7,031	6,471
		Total	9,725	11,460	13,620	14,080	14,432	15,628	15,841	16,057	14,507	13,123
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	39	61	91	76	104	84	89	132	137	117
		-of which MSM	3	0	9	6	9	9	8	14	25	11
		Female	210	229	277	319	275	286	341	373	380	298
		Total	249	290	368	395	379	370	430	505	517	415
C4D	Chlamydial ophthalmia neonatorum	Male (all)	4	1	1	27	0	2	1	3	1	2
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	6	2	3	21	5	1	1	1	2	0
		Total	10	3	4	48	5	3	2	4	3	2
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	2,704	3,170	3,285	3,237	3,767	3,904	4,225	4,459	4,035	4,133
		-of which MSM	117	199	210	241	285	341	282	432	420	557
		Female	1,533	1,854	1,829	2,097	2,544	2,744	2,779	2,911	2,467	2,339
		Total	4,237	5,024	5,114	5,334	6,311	6,648	7,004	7,370	6,502	6,473
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	4,531	4,567	5,024	5,335	5,750	5,601	5,211	5,534	5,704	4,935
		-of which MSM	253	286	345	317	365	394	337	477	509	550
		Female	0	0	533	608	465	339	362	423	441	418
		Total	4,531	4,567	5,557	5,943	6,215	5,940	5,573	5,957	6,145	5,353
C4I	Epidemiological treatment of NSGI	Male (all)	373	323	357	398	240	228	255	423	467	432
		-of which MSM	73	48	66	63	48	38	42	63	77	63
		Female	1,300	1,132	1,200	1,263	1,187	1,073	1,000	1,055	1,169	977
		Total	1,673	1,455	1,557	1,661	1,427	1,301	1,255	1,478	1,636	1,409
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	211	232	256	239	379	348	367	572	755	709
		-of which MSM	9	15	17	14	22	27	24	45	93	82
		Female	615	703	658	788	793	922	1,036	1,299	1,397	1,246
		Total	826	935	914	1,027	1,172	1,270	1,403	1,871	2,152	1,955
C6A	Trichomoniasis	Male (all)	17	28	26	16	24	31	42	39	31	29
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	496	474	501	433	430	385	373	427	390	430
		Total	513	502	527	449	454	416	415	466	421	459
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	245	178	204	270	299	267	222	229	224	212
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	5,427	6,244	6,977	7,310	8,045	8,442	9,187	10,095	10,139	9,280
		Total	5,672	6,422	7,181	7,580	8,344	8,709	9,409	10,324	10,363	9,492
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	1,335	1,100	1,146	1,142	1,198	1,240	1,339	1,537	1,523	1,272
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	666	394	537	401	571	738	911	1,028	354	273
		Total	2,001	1,494	1,683	1,543	1,769	1,978	2,250	2,565	1,877	1,545

North West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C7A	Anogenital candidosis	Male (all)	639	556	609	545	599	761	876	898	802	795
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	5,755	5,670	5,493	5,793	5,151	4,816	5,037	5,422	5,657	5,042
		Total	6,394	6,226	6,102	6,338	5,750	5,577	5,913	6,320	6,459	5,837
C7B	Epidemiological treatment of C6 & C7	Male (all)	335	305	346	351	325	412	465	636	612	568
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	470	454	416	448	394	444	483	671	738	700
		Total	805	759	762	799	719	856	948	1,307	1,350	1,269
C8,C9	Scabies/pediculosis pubis	Male (all)	484	450	313	283	225	259	227	184	225	217
		-of which MSM	73	82	57	55	45	55	31	42	86	77
		Female	116	78	57	41	31	29	41	31	29	23
		Total	600	528	370	324	256	288	268	215	254	240
C10A	Anogenital herpes simplex – first episode	Male (all)	703	683	652	730	933	977	1,220	1,412	1,536	1,572
		-of which MSM	32	53	35	47	72	56	57	95	96	121
		Female	1,235	1,204	1,225	1,313	1,425	1,657	2,022	2,404	2,471	2,473
		Total	1,938	1,887	1,877	2,043	2,358	2,634	3,242	3,816	4,007	4,045
C10B	Anogenital herpes simplex – recurrence	Male (all)	554	537	551	504	585	733	834	921	1,084	1,216
		-of which MSM	25	25	29	33	61	56	53	84	76	101
		Female	687	639	645	667	764	904	1,140	1,339	1,600	1,650
		Total	1,241	1,176	1,196	1,171	1,349	1,637	1,974	2,260	2,684	2,866
C11A	Anogenital warts – first episode	Male (all)	4,814	4,873	5,360	5,493	5,735	5,896	6,220	6,462	6,600	6,135
		-of which MSM	212	236	292	311	326	323	330	360	393	358
		Female	4,200	4,065	4,696	4,623	4,946	5,174	5,322	5,670	5,465	5,246
		Total	9,014	8,938	10,056	10,116	10,681	11,070	11,542	12,132	12,066	11,381
C11B	Anogenital warts – recurrence	Male (all)	2,387	2,321	2,378	2,537	3,052	3,341	3,637	3,748	3,139	2,996
		-of which MSM	121	112	109	138	163	207	206	209	201	184
		Female	1,328	1,438	1,430	1,519	1,728	2,059	2,361	2,364	1,965	1,976
		Total	3,715	3,759	3,808	4,056	4,780	5,400	5,998	6,112	5,104	4,973
C11C	Anogenital warts – re-registered cases	Male (all)	914	949	1,302	1,090	1,342	1,504	1,407	1,388	1,109	1,034
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	818	811	930	822	1,067	1,110	1,073	1,031	818	715
		Total	1,732	1,760	2,232	1,912	2,409	2,614	2,480	2,419	1,927	1,749
C12	Molluscum contagiosum	Male (all)	528	586	651	797	1,022	1,220	1,415	1,453	1,469	1,221
		-of which MSM	23	21	19	20	28	30	29	28	47	33
		Female	327	339	395	420	469	560	587	647	547	533
		Total	855	925	1,046	1,217	1,491	1,780	2,002	2,100	2,016	1,754
C13	Antigen positive viral hepatitis B	Male (all)	51	84	-	-	-	-	-	-	-	-
		-of which MSM	11	38	-	-	-	-	-	-	-	-
		Female	19	17	-	-	-	-	-	-	-	-
		Total	70	101	-	-	-	-	-	-	-	-
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	49	39	53	61	68	75	63	48
		-of which MSM	-	-	14	11	14	10	11	17	11	8
		Female	-	-	18	26	27	26	31	27	36	29
		Total	-	-	67	65	80	87	99	102	99	77

North West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

		2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	3	7	5	8	7	18	4	5
		-of which MSM	-	-	3	2	2	3	3	6	1	1
		Female	-	-	1	0	3	5	3	3	3	1
		Total	-	-	4	7	8	13	10	21	7	6
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	30	39	46	76	74	104	99	99
		-of which MSM	-	-	25	20	25	40	35	55	48	48
		Female	-	-	3	8	16	19	8	22	33	20
		Total	-	-	33	47	62	95	82	126	132	119
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	109	109	112	84	108	104	115	147	116	112
		-of which MSM	6	9	6	6	10	12	15	37	21	24
		Female	41	45	46	60	45	55	53	86	43	47
		Total	150	154	158	144	153	159	168	233	159	159
E1A	New HIV diagnosis - asymptomatic	Male (all)	161	226	273	293	351	349	330	372	329	291
		-of which MSM	103	117	145	165	199	187	159	177	206	169
		Female	43	92	125	146	175	155	136	182	130	96
		Total	204	318	398	439	526	504	466	554	459	387
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	658	949	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	121	211	-	-	-	-	-	-	-	-
		Total	779	1,160	-	-	-	-	-	-	-	-
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	1,830	2,121	3,496	4,172	4,082	5,069	4,483	4,831
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	451	661	1,044	1,472	1,492	2,045	1,657	1,829
		Total	-	-	2,281	2,782	4,540	5,644	5,574	7,114	6,140	6,660
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	112	47	52	71	72	52	49	55	28	39
		-of which MSM	72	19	32	41	51	30	20	31	16	26
		Female	9	8	18	17	21	17	20	23	9	6
		Total	121	55	70	88	93	69	69	78	37	45
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	853	975	-	-	-	-	-	-	-	-
		-of which MSM	0	0	-	-	-	-	-	-	-	-
		Female	117	160	-	-	-	-	-	-	-	-
		Total	970	1,135	-	-	-	-	-	-	-	-
E3A*	AIDS – first presentation	Male (all)	32	23	-	-	-	-	-	-	-	-
		-of which MSM	13	11	-	-	-	-	-	-	-	-
		Female	13	6	-	-	-	-	-	-	-	-
		Total	45	29	-	-	-	-	-	-	-	-
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	16	26	30	38	35	47	27	17
		-of which MSM	-	-	7	9	14	21	14	19	9	8
		Female	-	-	9	8	12	11	10	12	11	7
		Total	-	-	25	34	42	49	45	59	38	24
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	2	23	23	31	20	12	15	8
		-of which MSM	-	-	2	16	14	21	7	9	10	4
		Female	-	-	1	4	5	12	5	5	2	0
		Total	-	-	3	27	28	43	25	17	17	8

North West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	352	452	477	607	770	866	1,012	1,235	1,015	973	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	97	114	134	177	206	287	336	363	333	332	332
		Total	449	566	611	784	976	1,153	1,348	1,598	1,348	1,305	1,305
D2A	Urinary tract infection	Male (all)	189	146	104	130	124	154	142	203	220	211	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	401	340	309	344	376	430	572	902	954	888	888
		Total	590	486	413	474	500	584	714	1,105	1,174	1,099	1,099
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	4,022	3,983	4,788	5,125	5,794	6,343	6,801	7,807	7,604	7,033	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	4,588	4,355	4,224	4,010	3,993	4,718	5,573	6,162	5,692	5,296	5,296
		Total	8,610	8,338	9,012	9,135	9,787	11,061	12,374	13,969	13,296	12,329	12,329
D3	Other episodes not requiring treatment	Male (all)	11,646	14,055	14,868	18,104	24,389	28,922	33,898	37,858	39,158	38,404	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	11,721	13,903	14,303	16,840	23,110	26,790	32,305	36,249	37,233	37,457	37,457
		Total	23,367	27,958	29,171	34,944	47,499	55,712	66,203	74,107	76,392	75,863	75,863
P1A	HIV antibody test (no sexual health screen)	Male (all)	11,016	15,227	4,522	3,766	3,874	4,148	4,036	3,942	3,594	3,462	
		-of which MSM	1,539	1,976	879	735	824	913	879	862	846	780	
		Female	8,259	12,073	2,765	2,275	2,354	2,858	2,859	3,034	2,698	2,820	
		Total	19,275	27,300	7,287	6,041	6,228	7,006	6,895	6,976	6,292	6,283	6,283
P1B	HIV antibody test offered and refused	Male (all)	1,733	6,139	11,917	14,522	15,550	16,678	17,817	19,086	18,864	18,112	
		-of which MSM	102	283	472	673	758	812	755	755	827	845	
		Female	1,600	6,971	12,912	15,718	17,410	19,046	21,028	22,862	22,239	22,041	
		Total	3,333	13,110	24,829	30,240	32,960	35,724	38,845	41,948	41,104	40,153	40,153
P2	Hepatitis B vaccination (1st dose only)	Male (all)	1,314	1,496	1,641	1,698	1,807	1,860	1,957	2,258	2,074	2,219	
		-of which MSM	621	827	1,003	1,027	1,163	1,156	1,186	1,368	1,330	1,331	
		Female	412	425	453	424	455	502	599	644	590	695	
		Total	1,726	1,921	2,094	2,122	2,262	2,362	2,556	2,902	2,664	2,914	2,914
P3	Contraception (excluding condom provision)	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	1,751	1,977	1,523	1,374	1,299	2,207	3,357	5,443	6,670	9,292	
		Total	1,751	1,977	1,523	1,374	1,299	2,207	3,357	5,443	6,670	9,292	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	404	443	437	424	195	121	142	218	166	110	
		Total	404	443	437	424	195	121	142	218	166	110	

North West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	68	71	118	161	40	41	18	62	45	22
		Total	68	71	118	161	40	41	18	62	45	22
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	-	18,574	18,839	18,988	19,314	20,033	19,806	18,568
		-of which MSM	-	-	-	1,126	1,340	1,526	1,512	1,651	1,845	1,991
		Female	-	-	-	21,412	22,102	21,765	22,672	24,220	23,562	22,621
		Total	-	-	-	39,986	40,941	40,753	41,986	44,253	43,369	41,189
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	-	24,506	30,499	36,130	42,641	48,392	49,702	47,754
		-of which MSM	-	-	-	2,357	2,914	3,533	3,863	4,864	5,754	6,109
		Female	-	-	-	21,173	27,226	32,132	39,793	44,921	46,249	44,152
		Total	-	-	-	45,679	57,725	68,262	82,434	93,313	95,954	91,909
Total new STI diagnoses (for GUM clinics only)		Male (all)	17,757	18,942	21,151	22,103	23,727	24,851	25,353	26,359	25,964	23,800
		-of which MSM	1,525	1,762	2,039	1,906	2,284	2,423	2,245	2,715	2,998	3,174
		Female	13,798	14,556	16,866	17,032	17,369	18,251	18,968	20,165	19,060	17,937
		Total	31,555	33,498	38,017	39,135	41,096	43,102	44,321	46,524	45,027	41,737
Total other STI diagnoses		Male (all)	9,263	10,176	10,696	11,068	13,899	15,549	16,141	18,059	16,210	16,477
		-of which MSM	471	536	593	669	766	941	853	1,095	1,162	1,287
		Female	6,394	6,728	7,034	7,627	8,970	10,128	10,607	11,575	10,398	10,197
		Total	15,657	16,904	17,730	18,695	22,869	25,677	26,748	29,634	26,608	26,676
Total other GUM clinic diagnoses		Male (all)	6,925	6,461	7,391	7,732	8,551	9,426	10,109	11,654	11,267	10,355
		-of which MSM	17	47	48	39	51	65	64	115	81	81
		Female	17,839	18,033	18,579	18,985	18,856	19,855	22,018	24,698	23,860	21,708
		Total	24,764	24,494	25,970	26,717	27,407	29,281	32,127	36,352	35,127	32,064
Total services provided		Male (all)	25,709	36,917	69,488	81,170	94,958	106,726	119,663	131,569	133,198	128,519
		-of which MSM	2,262	3,086	5,064	5,918	6,999	7,940	8,195	9,500	10,602	11,056
		Female	23,743	35,349	68,667	79,216	93,956	105,300	122,613	137,373	139,241	139,078
		Total	49,452	72,266	138,155	160,386	188,914	212,026	242,276	268,942	272,445	267,603

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 g: South Central SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	11	20	60	67	103	101	69	78	64	67
		-of which MSM	2	3	18	17	38	54	25	34	31	40
		Female	5	6	9	11	15	4	5	7	7	5
		Total	16	26	69	78	118	105	74	85	71	72
A3	Early latent syphilis (first 2 years)	Male (all)	16	16	19	28	32	37	35	20	35	23
		-of which MSM	1	1	8	11	13	20	14	8	10	14
		Female	14	13	12	19	9	7	9	4	10	6
		Total	30	29	31	47	41	44	44	24	45	29
A4,A5,A6	Other acquired syphilis	Male (all)	23	27	43	37	54	41	49	53	45	46
		-of which MSM	1	4	2	3	2	5	4	14	4	9
		Female	18	25	34	42	55	37	30	38	31	37
		Total	41	52	77	79	109	78	79	91	76	83
A7	Congenital syphilis, aged under 2	Male (all)	0	0	0	0	0	1	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	2	0	0	0	0
Total	0	0	0	0	0	3	0	0	0	0		
A8	Congenital syphilis, aged 2 or over	Male (all)	0	0	0	1	0	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	0	1	0	0	0
Total	0	0	0	1	0	0	1	0	0	0		
A9	Epidemiological treatment of suspected syphilis	Male (all)	1	0	9	5	23	38	23	34	20	37
		-of which MSM	0	0	3	3	11	16	6	25	10	20
		Female	1	0	3	5	9	2	11	12	4	8
		Total	2	0	12	10	32	40	34	46	24	45
B1,B2	Uncomplicated gonorrhoea	Male (all)	635	681	833	671	547	487	430	438	452	479
		-of which MSM	104	84	94	92	109	130	91	85	119	135
		Female	307	340	398	305	205	195	218	180	193	239
		Total	942	1,021	1,231	976	752	682	648	618	645	718
B3	Gonococcal ophthalmia neonatorum	Male (all)	3	4	0	3	0	0	1	0	2	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3	0	1	3	1	1	0	1	0	1
Total	6	4	1	6	1	1	1	1	1	2	1	
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	80	99	129	118	117	103	141	122	171	188
		-of which MSM	14	19	10	17	41	33	30	26	44	62
		Female	83	90	124	125	114	86	118	128	113	140
		Total	163	189	253	243	231	189	259	250	284	328
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	5	6	5	5	0	2	3	6	3	4
		-of which MSM	1	0	0	0	0	0	1	0	0	0
		Female	13	21	25	11	15	11	4	9	8	10
		Total	18	27	30	16	15	13	7	15	11	14

South Central SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	2	2	9	5	5	11	10	10	8	13
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3	6	5	13	8	6	13	9	3	2
		Total	5	8	14	18	13	17	23	19	11	15
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	1,810	1,989	2,314	2,705	3,097	3,139	3,697	3,779	3,233	2,953
		-of which MSM	17	14	28	28	43	89	80	152	176	188
		Female	2,539	2,760	3,320	3,722	3,932	3,473	3,599	3,477	2,973	2,952
		Total	4,349	4,749	5,634	6,427	7,029	6,612	7,296	7,256	6,209	5,905
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	43	44	41	39	48	48	57	49	45	53
		-of which MSM	1	0	1	1	4	0	0	2	0	4
		Female	150	128	144	193	172	156	178	164	123	82
		Total	193	172	185	232	220	204	235	213	168	135
C4D	Chlamydial ophthalmia neonatorum	Male (all)	3	4	1	0	0	0	1	2	1	1
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	14	7	4	2	0	1	0	3	0	3
		Total	17	11	5	2	0	1	1	5	1	4
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	1,150	1,366	1,494	1,829	1,931	1,720	2,176	2,317	2,069	2,203
		-of which MSM	6	7	12	15	23	41	50	79	96	126
		Female	640	755	952	1,153	1,353	1,371	1,554	1,608	1,380	1,431
		Total	1,790	2,121	2,446	2,982	3,284	3,091	3,730	3,925	3,449	3,634
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	4,096	4,320	5,776	5,871	6,188	5,404	5,022	5,034	4,636	4,258
		-of which MSM	157	126	181	201	248	235	164	227	229	238
		Female	0	0	771	841	958	794	629	661	745	674
		Total	4,096	4,320	6,547	6,712	7,146	6,198	5,651	5,695	5,383	4,932
C4I	Epidemiological treatment of NSGI	Male (all)	665	576	623	665	597	465	542	507	536	575
		-of which MSM	29	22	22	36	42	32	24	37	31	42
		Female	2,084	1,904	1,708	1,878	1,774	1,476	1,478	1,343	1,261	1,146
		Total	2,749	2,480	2,331	2,543	2,371	1,941	2,020	1,850	1,797	1,721
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	388	487	440	440	513	427	503	493	512	487
		-of which MSM	9	6	8	7	4	10	6	29	22	18
		Female	962	909	1,013	1,221	1,159	865	1,047	1,028	1,120	1,106
		Total	1,350	1,396	1,453	1,661	1,672	1,292	1,550	1,521	1,632	1,593
C6A	Trichomoniasis	Male (all)	20	27	17	22	30	19	27	14	16	18
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	221	222	217	231	209	160	165	177	201	148
		Total	241	249	234	253	239	179	192	191	217	166
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	84	113	107	74	75	112	144	214	246	275
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	4,200	3,948	4,737	4,278	4,616	4,203	4,893	5,458	5,420	5,530
		Total	4,284	4,061	4,844	4,352	4,691	4,315	5,037	5,672	5,668	5,805
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	1,150	1,158	1,239	1,136	1,045	1,037	1,249	1,444	1,348	1,214
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	869	784	719	696	643	501	614	629	603	509
		Total	2,019	1,942	1,958	1,832	1,688	1,538	1,863	2,073	1,951	1,723

South Central SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C7A	Anogenital candidosis	Male (all)	472	492	441	463	374	332	411	512	640	660	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	5,160	5,704	6,283	5,028	4,546	4,348	4,916	5,528	5,445	5,546	
		Total	5,632	6,196	6,724	5,491	4,920	4,680	5,327	6,040	6,088	6,206	
C7B	Epidemiological treatment of C6 & C7	Male (all)	168	164	130	164	123	166	260	288	302	236	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	829	575	741	767	276	299	439	468	425	459	
		Total	997	739	871	931	399	465	699	756	727	695	
C8,C9	Scabies/pediculosis pubis	Male (all)	247	214	179	168	111	115	125	126	129	115	
		-of which MSM	32	18	22	23	24	27	19	27	32	16	
		Female	49	32	28	27	21	19	22	21	23	17	
		Total	296	246	207	195	132	134	147	147	152	132	
C10A	Anogenital herpes simplex – first episode	Male (all)	513	445	505	492	463	583	674	801	700	848	
		-of which MSM	14	4	11	20	13	29	23	29	31	35	
		Female	807	816	851	769	813	916	1,080	1,218	1,192	1,361	
		Total	1,320	1,261	1,356	1,261	1,276	1,499	1,754	2,019	1,892	2,209	
C10B	Anogenital herpes simplex – recurrence	Male (all)	573	591	538	496	486	569	588	725	656	623	
		-of which MSM	7	7	8	19	22	24	18	37	27	40	
		Female	827	785	746	700	590	727	797	994	946	880	
		Total	1,400	1,376	1,284	1,196	1,076	1,296	1,385	1,719	1,602	1,503	
C11A	Anogenital warts – first episode	Male (all)	2,353	2,474	2,479	2,472	2,492	2,647	2,742	2,918	2,966	2,944	
		-of which MSM	39	49	64	68	84	69	91	71	115	141	
		Female	2,193	2,243	2,343	2,237	2,327	2,429	2,694	2,730	2,650	2,531	
		Total	4,546	4,717	4,822	4,709	4,819	5,076	5,436	5,648	5,617	5,475	
C11B	Anogenital warts – recurrence	Male (all)	1,751	1,710	1,664	1,474	1,644	1,808	1,820	1,787	2,078	2,116	
		-of which MSM	29	27	32	47	57	63	72	52	71	77	
		Female	953	930	1,083	925	1,035	1,250	1,250	1,273	1,310	1,282	
		Total	2,704	2,640	2,747	2,399	2,679	3,058	3,070	3,060	3,389	3,398	
C11C	Anogenital warts – re-registered cases	Male (all)	516	453	432	423	459	432	493	537	644	768	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	516	383	469	412	402	365	453	418	484	568	
		Total	1,032	836	901	835	861	797	946	955	1,128	1,336	
C12	Molluscum contagiosum	Male (all)	382	442	471	547	587	626	687	748	737	650	
		-of which MSM	6	4	3	7	8	8	11	13	16	11	
		Female	155	198	190	236	264	233	258	302	275	256	
		Total	537	640	661	783	851	859	945	1,050	1,012	906	
C13	Antigen positive viral hepatitis B	Male (all)	27	21	-	-	-	-	-	-	-	-	
		-of which MSM	2	2	-	-	-	-	-	-	-	-	
		Female	11	20	-	-	-	-	-	-	-	-	
		Total	38	41	-	-	-	-	-	-	-	-	
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	23	22	24	17	31	43	36	27	
		-of which MSM	-	-	1	1	2	0	2	3	2	3	
		Female	-	-	7	11	4	12	16	18	17	7	
		Total	-	-	30	33	28	29	47	61	53	34	

South Central SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	1	1	1	5	1	4	2	6
		-of which MSM	-	-	0	0	0	0	0	0	0	0
		Female	-	-	0	0	0	5	3	0	1	0
		Total	-	-	1	1	1	10	4	4	3	6
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	5	3	0	1	15	11	10	6
		-of which MSM	-	-	0	0	0	0	2	0	2	1
		Female	-	-	0	1	1	2	5	8	11	2
		Total	-	-	5	4	1	3	20	19	21	8
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	47	42	57	40	69	47	65	54	61	51
		-of which MSM	0	1	1	2	3	0	2	2	1	4
		Female	15	23	25	16	27	26	20	18	23	10
		Total	62	65	82	56	96	73	85	72	87	61
E1A	New HIV diagnosis - asymptomatic	Male (all)	82	90	137	123	129	114	110	129	108	91
		-of which MSM	21	14	18	31	41	38	40	46	33	38
		Female	84	150	203	149	145	93	103	83	84	62
		Total	166	240	340	272	274	207	213	212	192	153
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	360	495	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	238	423	-	-	-	-	-	-	-	-
		Total	598	918	-	-	-	-	-	-	-	-
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	779	983	1,161	1,355	1,570	1,812	2,346	2,530
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	762	1,009	1,194	1,315	1,394	1,571	2,121	2,152
		Total	-	-	1,541	1,992	2,355	2,670	2,964	3,383	4,467	4,682
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	32	30	39	16	42	43	50	61	45	48
		-of which MSM	4	3	3	2	9	11	11	22	9	17
		Female	28	49	35	34	34	49	47	56	38	32
		Total	60	79	74	50	76	92	97	117	83	80
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	399	401	-	-	-	-	-	-	-	-
		-of which MSM	0	0	-	-	-	-	-	-	-	-
		Female	186	195	-	-	-	-	-	-	-	-
		Total	585	596	-	-	-	-	-	-	-	-
E3A*	AIDS – first presentation	Male (all)	22	29	-	-	-	-	-	-	-	-
		-of which MSM	8	4	-	-	-	-	-	-	-	-
		Female	9	22	-	-	-	-	-	-	-	-
		Total	31	51	-	-	-	-	-	-	-	-
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	14	19	17	16	17	24	15	12
		-of which MSM	-	-	4	1	3	1	4	15	4	5
		Female	-	-	16	12	19	7	11	9	8	7
		Total	-	-	30	31	36	23	28	33	23	19
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	0	0	8	5	6	6	9	4
		-of which MSM	-	-	0	0	1	3	2	6	2	3
		Female	-	-	1	2	7	5	4	3	1	2
		Total	-	-	1	2	15	10	10	9	10	6

South Central SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	167	182	219	252	308	357	353	415	492	548	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	73	91	118	142	172	187	197	207	272	268	268
		Total	240	273	337	394	480	544	550	622	764	816	816
D2A	Urinary tract infection	Male (all)	94	116	111	115	88	94	113	116	116	109	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	683	687	771	668	781	630	900	1,118	1,205	1,171	1,171
		Total	777	803	882	783	869	724	1,013	1,234	1,322	1,280	1,280
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	4,043	3,813	3,925	3,766	3,909	3,676	3,354	4,328	4,540	4,700	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	4,672	4,367	4,469	4,645	5,609	4,691	3,527	4,699	4,295	4,392	4,392
		Total	8,715	8,180	8,394	8,411	9,518	8,367	6,881	9,027	8,837	9,092	9,092
D3	Other episodes not requiring treatment	Male (all)	8,585	9,964	7,577	8,733	11,862	14,116	15,048	16,257	16,852	18,777	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	10,314	11,657	8,941	9,991	13,746	15,968	17,464	19,809	19,642	22,770	22,770
		Total	18,899	21,621	16,518	18,724	25,608	30,084	32,512	36,066	36,511	41,548	41,548
P1A	HIV antibody test (no sexual health screen)	Male (all)	7,705	9,579	2,456	2,162	2,269	2,367	1,915	2,936	1,819	2,193	
		-of which MSM	549	527	176	200	229	273	200	196	238	295	
		Female	6,996	8,962	1,686	1,549	1,762	1,976	1,676	1,801	1,548	1,795	
		Total	14,701	18,541	4,142	3,711	4,031	4,343	3,591	4,737	3,398	3,988	
P1B	HIV antibody test offered and refused	Male (all)	3,923	5,212	5,600	6,299	7,651	7,989	8,098	8,695	8,466	8,815	
		-of which MSM	77	98	94	149	146	182	130	172	223	246	
		Female	4,542	6,040	7,171	7,730	9,201	10,065	11,897	14,787	18,059	19,418	
		Total	8,465	11,252	12,771	14,029	16,852	18,054	19,995	23,482	26,530	28,233	
P2	Hepatitis B vaccination (1st dose only)	Male (all)	464	614	719	803	811	861	986	1,105	1,057	1,138	
		-of which MSM	202	193	218	289	351	361	303	377	535	583	
		Female	113	147	172	198	251	311	366	468	469	456	
		Total	577	761	891	1,001	1,062	1,172	1,352	1,573	1,528	1,594	
P3	Contraception (excluding condom provision)	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	936	1,063	3,533	1,785	1,454	1,293	2,043	3,066	1,950	2,362	
		Total	936	1,063	3,533	1,785	1,454	1,293	2,043	3,066	1,950	2,362	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	600	457	440	318	136	96	82	114	69	35	
		Total	600	457	440	318	136	96	82	114	69	35	

South Central SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	109	101	90	71	44	16	25	36	17	7
		Total	109	101	90	71	44	16	25	36	17	7
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	10,982	10,049	10,000	9,779	10,209	10,325	9,904	9,596
		-of which MSM	-	-	202	252	262	317	237	276	361	415
		Female	-	-	15,046	13,994	14,086	12,986	13,857	15,361	14,942	14,874
		Total	-	-	26,028	24,043	24,086	22,765	24,066	25,686	24,850	24,470
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	11,551	14,262	18,777	20,724	25,123	27,565	27,870	29,360
		-of which MSM	-	-	434	711	887	1,090	1,046	1,412	1,860	2,313
		Female	-	-	12,030	14,269	19,687	21,691	26,470	30,347	29,735	31,312
		Total	-	-	23,581	28,531	38,464	42,415	51,593	57,912	57,618	60,673
Total new STI diagnoses (for GUM clinics only)	Male (all)	10,635	11,287	13,338	13,690	14,404	13,819	14,258	14,728	13,704	13,063	
	-of which MSM	408	326	463	509	641	721	580	760	827	900	
	Female	7,530	7,893	9,580	10,031	10,305	9,417	10,082	10,135	9,653	9,490	
	Total	18,165	19,180	22,918	23,721	24,709	23,236	24,340	24,863	23,363	22,553	
Total other STI diagnoses	Male (all)	5,713	5,937	5,931	6,286	6,788	6,894	7,763	8,317	9,069	9,639	
	-of which MSM	94	90	89	140	199	217	206	276	285	379	
	Female	5,645	5,610	6,005	6,398	6,706	6,825	7,287	7,599	7,923	7,918	
	Total	11,358	11,547	11,936	12,684	13,494	13,719	15,050	15,916	16,993	17,557	
Total other GUM clinic diagnoses	Male (all)	6,085	5,919	6,039	5,784	5,708	5,487	5,643	7,014	7,301	7,284	
	-of which MSM	2	3	2	3	5	0	6	5	5	8	
	Female	17,148	16,666	18,282	16,499	16,683	14,829	15,440	18,094	17,531	17,668	
	Total	23,233	22,585	24,321	22,283	22,391	20,316	21,083	25,108	24,843	24,952	
Total services provided	Male (all)	20,677	25,369	38,885	42,308	51,370	55,836	61,379	66,883	65,968	69,879	
	-of which MSM	828	818	1,124	1,601	1,875	2,223	1,916	2,433	3,217	3,852	
	Female	22,901	27,869	48,579	49,516	60,187	64,290	73,773	85,639	86,345	92,987	
	Total	43,578	53,238	87,464	91,824	111,557	120,126	135,152	152,522	152,385	162,868	

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 h: South East Coast SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	25	60	88	128	75	93	102	99	94	85
		-of which MSM	12	42	69	100	50	69	49	75	45	39
		Female	2	10	7	5	14	8	8	12	7	5
		Total	27	70	95	133	89	101	110	111	101	90
A3	Early latent syphilis (first 2 years)	Male (all)	14	21	32	43	55	48	67	52	38	36
		-of which MSM	5	16	21	32	40	39	27	41	18	5
		Female	5	15	8	10	10	5	7	5	9	6
		Total	19	36	40	53	65	53	74	57	47	42
A4,A5,A6	Other acquired syphilis	Male (all)	23	26	40	44	43	42	51	64	42	41
		-of which MSM	10	12	18	15	12	18	13	15	14	13
		Female	14	12	14	29	31	32	33	38	32	39
		Total	37	38	54	73	74	74	84	102	74	80
A7	Congenital syphilis, aged under 2	Male (all)	0	0	0	0	0	0	0	1	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	0	0	0	0	0
Total	0	0	0	0	0	0	0	1	0	0		
A8	Congenital syphilis, aged 2 or over	Male (all)	0	0	0	0	0	0	0	1	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	0	0	0	0	0
Total	0	0	0	0	0	0	0	1	0	0		
A9	Epidemiological treatment of suspected syphilis	Male (all)	6	4	7	14	17	26	42	46	39	24
		-of which MSM	0	2	5	13	11	22	20	34	18	11
		Female	2	1	2	4	2	7	6	4	5	4
		Total	8	5	9	18	19	33	48	50	44	28
B1,B2	Uncomplicated gonorrhoea	Male (all)	683	706	635	700	619	616	601	417	486	526
		-of which MSM	342	266	291	318	323	307	196	167	113	179
		Female	153	171	190	193	149	181	154	146	174	141
		Total	836	877	825	893	768	797	755	563	660	667
B3	Gonococcal ophthalmia neonatorum	Male (all)	2	0	0	0	0	1	0	1	0	2
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	1	0	1	1	0	0
Total	2	0	0	0	1	1	1	2	0	2		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	115	169	158	186	218	207	236	247	278	259
		-of which MSM	62	68	85	87	93	86	70	80	68	66
		Female	70	77	122	137	130	145	166	171	193	152
		Total	185	246	280	323	348	352	402	418	471	411
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	4	5	6	4	4	1	5	1	5	5
		-of which MSM	2	5	2	2	2	0	0	1	0	0
		Female	3	2	5	7	3	3	5	10	6	4
		Total	7	7	11	11	7	4	10	11	11	9

South East Coast SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	1	2	1	5	29	15	9	18	14	20
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	1	2	1	0	3	4	5	1	2
		Total	1	3	3	6	29	18	13	23	15	22
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	1,160	1,664	1,781	2,506	2,540	2,889	3,282	3,400	3,125	2,957
		-of which MSM	63	100	138	174	214	174	171	243	147	151
		Female	1,486	2,061	2,349	2,925	2,741	2,864	3,111	3,534	3,198	2,907
		Total	2,646	3,725	4,130	5,431	5,281	5,753	6,393	6,934	6,323	5,864
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	14	27	22	29	45	35	47	50	66	53
		-of which MSM	2	4	5	5	17	5	2	9	4	1
		Female	49	58	48	67	91	83	98	108	127	149
		Total	63	85	70	96	136	118	145	158	193	202
C4D	Chlamydial ophthalmia neonatorum	Male (all)	0	0	0	1	1	3	0	0	2	1
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	2	1	0	1	6	0	2	0	0
		Total	1	2	1	1	2	9	0	2	2	1
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	792	1,107	1,330	1,659	1,667	1,623	1,967	2,058	2,045	1,964
		-of which MSM	35	44	87	91	87	113	136	147	122	105
		Female	648	845	973	1,188	1,350	1,296	1,582	1,900	1,681	1,351
		Total	1,440	1,952	2,303	2,847	3,017	2,919	3,549	3,958	3,726	3,315
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	4,711	5,391	4,913	5,213	5,361	5,655	4,613	4,601	4,196	3,229
		-of which MSM	596	705	681	649	582	575	280	321	276	253
		Female	0	0	738	1,141	1,213	1,099	835	901	800	716
		Total	4,711	5,391	5,651	6,354	6,574	6,754	5,448	5,502	4,998	3,945
C4I	Epidemiological treatment of NSGI	Male (all)	575	710	646	525	538	455	430	500	629	534
		-of which MSM	82	105	92	112	81	79	45	57	47	38
		Female	1,454	1,725	1,449	1,316	1,432	1,471	1,422	1,199	1,134	879
		Total	2,029	2,435	2,095	1,841	1,970	1,926	1,852	1,699	1,763	1,413
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	240	293	234	283	256	235	271	298	368	455
		-of which MSM	33	31	28	31	39	35	29	41	29	20
		Female	440	484	440	515	424	412	388	503	689	650
		Total	680	777	674	798	680	647	659	801	1,057	1,105
C6A	Trichomoniasis	Male (all)	17	13	5	15	8	12	25	10	18	20
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	108	88	84	156	100	122	117	106	134	136
		Total	125	101	89	171	108	134	142	116	152	156
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	101	99	71	104	112	110	99	131	183	262
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3,779	4,385	4,590	4,547	4,622	4,939	5,909	6,602	6,580	7,050
		Total	3,880	4,484	4,661	4,651	4,734	5,049	6,008	6,733	6,764	7,312
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	748	774	602	631	600	459	668	867	652	742
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	650	366	236	197	151	144	185	288	230	215
		Total	1,398	1,140	838	828	751	603	853	1,155	882	957

South East Coast SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C7A	Anogenital candidosis	Male (all)	627	686	612	692	718	773	861	931	906	1,060
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	4,916	5,454	5,090	5,009	4,912	4,747	5,650	5,846	5,723	5,865
		Total	5,543	6,140	5,702	5,701	5,630	5,520	6,511	6,777	6,630	6,925
C7B	Epidemiological treatment of C6 & C7	Male (all)	144	195	309	229	263	208	216	179	479	354
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	338	428	537	502	463	343	366	333	627	479
		Total	482	623	846	731	726	551	582	512	1,106	833
C8,C9	Scabies/pediculosis pubis	Male (all)	272	239	181	142	116	117	83	88	131	107
		-of which MSM	75	78	58	41	38	37	20	24	29	17
		Female	54	26	32	13	19	20	24	12	17	11
		Total	326	265	213	155	135	137	107	100	148	118
C10A	Anogenital herpes simplex – first episode	Male (all)	427	492	464	476	437	506	653	769	821	912
		-of which MSM	26	43	47	45	36	41	37	53	58	53
		Female	683	823	800	831	770	838	1,047	1,209	1,321	1,498
		Total	1,110	1,315	1,264	1,307	1,207	1,344	1,700	1,978	2,142	2,410
C10B	Anogenital herpes simplex – recurrence	Male (all)	483	493	517	490	536	520	476	603	735	830
		-of which MSM	50	54	56	56	47	64	38	55	54	58
		Female	629	690	764	662	688	676	726	906	1,049	1,171
		Total	1,112	1,183	1,281	1,152	1,224	1,196	1,202	1,509	1,784	2,001
C11A	Anogenital warts – first episode	Male (all)	2,473	2,785	2,499	2,599	2,655	2,794	2,901	3,126	3,376	3,330
		-of which MSM	142	193	192	160	190	175	143	185	155	140
		Female	2,127	2,365	2,205	2,504	2,506	2,540	2,863	2,969	2,979	2,880
		Total	4,600	5,150	4,704	5,103	5,161	5,334	5,764	6,095	6,355	6,210
C11B	Anogenital warts – recurrence	Male (all)	1,577	1,647	1,510	1,525	1,767	1,762	1,929	2,075	2,136	2,009
		-of which MSM	101	94	95	87	93	98	75	134	88	99
		Female	908	897	895	986	1,205	1,296	1,612	1,482	1,416	1,380
		Total	2,485	2,544	2,405	2,511	2,972	3,058	3,541	3,557	3,552	3,389
C11C	Anogenital warts – re-registered cases	Male (all)	709	751	673	604	619	625	662	814	754	755
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	604	637	572	666	591	618	638	760	652	562
		Total	1,313	1,388	1,245	1,270	1,210	1,243	1,300	1,574	1,406	1,317
C12	Molluscum contagiosum	Male (all)	412	459	481	564	688	654	711	850	767	733
		-of which MSM	21	16	21	11	27	25	30	23	17	14
		Female	200	248	239	287	378	354	386	351	328	307
		Total	612	707	720	851	1,066	1,008	1,097	1,201	1,095	1,040
C13	Antigen positive viral hepatitis B	Male (all)	40	46	-	-	-	-	-	-	-	-
		-of which MSM	21	13	-	-	-	-	-	-	-	-
		Female	21	18	-	-	-	-	-	-	-	-
		Total	61	64	-	-	-	-	-	-	-	-
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	39	51	41	39	41	22	46	31
		-of which MSM	-	-	15	9	11	10	5	1	8	2
		Female	-	-	16	15	15	20	10	23	15	21
		Total	-	-	55	66	56	59	51	45	61	52

South East Coast SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	5	3	2	2	2	6	3	3	
		-of which MSM	-	-	2	0	0	0	0	1	1	0	
		Female	-	-	2	0	0	1	1	1	1	1	1
		Total	-	-	7	3	2	3	3	7	4	4	
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	4	8	15	14	21	48	44	33	
		-of which MSM	-	-	1	0	2	0	5	9	11	8	
		Female	-	-	10	8	13	12	19	33	19	19	
		Total	-	-	14	16	28	26	40	81	63	52	
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	87	101	105	108	106	75	63	74	75	89	
		-of which MSM	21	20	21	22	22	27	2	6	12	9	
		Female	42	38	43	48	36	16	26	25	30	18	
		Total	129	139	148	156	142	91	89	99	105	107	
E1A	New HIV diagnosis - asymptomatic	Male (all)	111	154	171	206	195	186	202	219	176	175	
		-of which MSM	63	65	96	99	115	124	84	115	83	53	
		Female	55	90	130	144	142	120	86	100	67	56	
		Total	166	244	301	350	337	306	288	319	243	231	
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	1,048	1,381	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	340	559	-	-	-	-	-	-	-	-	
		Total	1,388	1,940	-	-	-	-	-	-	-	-	
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	2,377	3,139	3,967	4,437	3,543	3,684	4,541	4,495	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	-	-	750	1,132	1,484	1,642	1,588	1,819	1,601	1,562	
		Total	-	-	3,127	4,271	5,451	6,079	5,131	5,503	6,143	6,057	
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	48	48	59	70	112	33	23	35	23	20	
		-of which MSM	29	24	32	25	26	12	5	19	10	8	
		Female	20	27	40	42	77	20	19	23	12	8	
		Total	68	75	99	112	189	53	42	58	35	28	
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	1,254	1,252	-	-	-	-	-	-	-	-	
		-of which MSM	0	0	-	-	-	-	-	-	-	-	
		Female	158	197	-	-	-	-	-	-	-	-	
		Total	1,412	1,449	-	-	-	-	-	-	-	-	
E3A*	AIDS – first presentation	Male (all)	38	40	-	-	-	-	-	-	-	-	
		-of which MSM	19	21	-	-	-	-	-	-	-	-	
		Female	17	25	-	-	-	-	-	-	-	-	
		Total	55	65	-	-	-	-	-	-	-	-	
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	7	18	32	11	16	21	19	9	
		-of which MSM	-	-	4	7	9	8	5	9	8	3	
		Female	-	-	7	10	12	4	2	7	9	7	
		Total	-	-	14	28	44	15	18	28	28	16	
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	3	6	14	10	2	11	3	0	
		-of which MSM	-	-	1	4	5	5	1	5	0	0	
		Female	-	-	4	0	6	0	1	1	2	0	
		Total	-	-	7	6	20	10	3	12	5	0	

South East Coast SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	737	841	919	943	1,113	1,209	868	875	1,074	937	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	94	160	189	206	278	313	248	295	286	250	
		Total	831	1,001	1,108	1,149	1,391	1,522	1,116	1,170	1,360	1,187	
D2A	Urinary tract infection	Male (all)	138	109	120	167	97	137	146	196	194	196	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	620	575	477	659	604	628	711	818	889	877	
		Total	758	684	597	826	701	765	857	1,014	1,083	1,073	
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	5,326	5,839	7,280	6,818	7,178	7,004	7,243	9,287	6,146	5,521	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	4,194	4,407	3,858	4,367	4,625	4,369	5,279	6,708	5,703	5,704	
		Total	9,520	10,246	11,138	11,185	11,803	11,373	12,522	15,995	11,850	11,226	
D3	Other episodes not requiring treatment	Male (all)	7,916	9,516	9,023	11,639	12,788	12,643	16,997	20,683	20,372	21,504	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	7,898	10,332	10,130	12,355	14,444	13,920	17,810	23,092	22,604	24,115	
		Total	15,814	19,848	19,153	23,994	27,232	26,563	34,807	43,775	42,979	45,622	
P1A	HIV antibody counselling - with testing	Male (all)	6,687	8,934	4,457	4,357	3,914	2,681	2,664	2,504	2,375	2,531	
		-of which MSM	1,477	1,629	622	610	571	509	377	481	429	380	
		Female	5,133	7,230	3,530	3,538	3,259	2,134	2,066	2,159	1,860	1,755	
		Total	11,820	16,164	7,987	7,895	7,173	4,815	4,730	4,663	4,235	4,286	
P1B	HIV antibody counselling - without testing	Male (all)	970	1,967	6,443	7,944	8,716	9,518	9,281	9,333	9,084	9,327	
		-of which MSM	180	361	674	558	438	473	362	451	419	375	
		Female	807	1,874	7,218	9,208	10,445	11,319	11,737	12,493	12,558	13,098	
		Total	1,777	3,841	13,661	17,152	19,161	20,837	21,018	21,826	21,643	22,425	
P2	Hepatitis B vaccination	Male (all)	790	1,090	1,419	1,174	1,258	1,254	1,392	1,586	1,569	1,710	
		-of which MSM	463	567	902	678	716	725	662	832	582	573	
		Female	201	242	284	339	435	428	505	632	642	579	
		Total	991	1,332	1,703	1,513	1,693	1,682	1,897	2,218	2,212	2,289	
P3	Family Planning	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	3,063	2,019	1,909	1,793	2,729	1,518	1,389	1,545	1,512	2,513	
		Total	3,063	2,019	1,909	1,793	2,729	1,518	1,389	1,545	1,512	2,513	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	375	374	315	428	177	115	117	130	98	94	
		Total	375	374	315	428	177	115	117	130	98	94	

South East Coast SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	193	106	102	91	63	40	41	58	26	12
		Total	193	106	102	91	63	40	41	58	26	12
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	9,431	10,385	10,114	10,198	10,023	9,757	9,180	9,489
		-of which MSM	-	-	853	790	770	961	595	843	612	384
		Female	-	-	12,068	13,367	13,362	12,547	13,185	13,459	12,636	13,534
		Total	-	-	21,499	23,752	23,476	22,745	23,208	23,216	21,817	23,023
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	9,987	13,955	16,717	18,809	22,719	26,639	28,394	30,945
		-of which MSM	-	-	1,540	1,758	1,769	1,901	1,708	2,407	1,945	2,245
		Female	-	-	9,860	13,659	16,387	18,730	23,664	28,595	30,554	32,839
		Total	-	-	19,847	27,614	33,104	37,539	46,383	55,234	58,954	63,788
Total new STI diagnoses (for GUM clinics only)	Male (all)	10,612	12,359	11,579	13,001	13,227	13,900	13,611	14,054	13,723	12,672	
	-of which MSM	1,411	1,588	1,685	1,699	1,708	1,626	1,078	1,326	992	936	
	Female	5,385	6,469	7,324	8,851	8,649	8,676	9,154	10,001	9,878	9,483	
	Total	15,997	18,828	18,903	21,852	21,876	22,576	22,765	24,055	23,603	22,155	
Total other STI diagnoses	Male (all)	7,359	8,421	8,180	9,136	10,500	10,920	10,206	10,980	12,278	11,851	
	-of which MSM	359	400	439	465	429	485	398	527	411	390	
	Female	4,939	5,827	5,735	6,326	7,199	7,502	8,023	8,578	8,051	7,350	
	Total	12,298	14,248	13,915	15,462	17,699	18,422	18,229	19,558	20,330	19,201	
Total other GUM clinic diagnoses	Male (all)	7,211	7,849	9,147	8,811	9,132	8,821	9,360	11,741	8,728	8,291	
	-of which MSM	42	33	39	31	35	37	12	17	32	19	
	Female	15,128	16,151	15,276	15,871	15,681	15,374	18,314	20,865	19,941	20,355	
	Total	22,339	24,000	24,423	24,682	24,813	24,195	27,674	32,606	28,672	28,647	
Total services provided	Male (all)	16,363	21,507	40,760	49,454	53,507	55,103	63,076	70,502	70,974	75,506	
	-of which MSM	2,120	2,557	4,591	4,394	4,264	4,569	3,704	5,014	3,987	3,957	
	Female	17,102	21,697	44,999	54,259	61,061	60,596	70,356	81,975	82,366	88,433	
	Total	33,465	43,204	85,759	103,713	114,568	115,699	133,432	152,477	153,352	163,946	

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 i: South West SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	10	32	53	80	95	131	140	133	93	69
		-of which MSM	6	16	16	44	55	61	91	73	48	42
		Female	3	3	4	17	16	11	12	26	8	12
		Total	13	35	57	97	111	142	152	159	101	81
A3	Early latent syphilis (first 2 years)	Male (all)	19	9	9	28	25	23	36	44	21	19
		-of which MSM	5	3	2	12	11	9	18	24	12	13
		Female	10	6	7	11	11	7	13	6	10	4
		Total	29	15	16	39	36	30	49	50	31	23
A4,A5,A6	Other acquired syphilis	Male (all)	22	19	26	31	50	40	56	65	41	36
		-of which MSM	3	3	4	7	13	10	19	28	15	14
		Female	24	12	11	17	24	21	22	35	23	30
		Total	46	31	37	48	74	61	78	100	64	66
A7	Congenital syphilis, aged under 2	Male (all)	1	0	0	1	0	1	0	1	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	1	0	0	0	0	0
Total	1	0	0	1	1	1	0	1	0	0		
A8	Congenital syphilis, aged 2 or over	Male (all)	0	0	0	0	0	0	0	0	1	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	1	0	1	1	0	0	0	0	0
Total	0	1	0	1	1	0	0	0	0	1	0	
A9	Epidemiological treatment of suspected syphilis	Male (all)	5	7	8	19	26	30	55	65	46	25
		-of which MSM	0	1	3	2	12	14	31	40	16	11
		Female	1	3	2	6	11	7	6	8	10	3
		Total	6	10	10	25	37	37	61	73	56	28
B1,B2	Uncomplicated gonorrhoea	Male (all)	802	1,045	950	765	674	626	497	502	549	475
		-of which MSM	143	154	157	134	111	139	90	144	117	179
		Female	307	382	436	320	287	298	262	243	253	179
		Total	1,109	1,427	1,386	1,085	961	924	759	745	802	654
B3	Gonococcal ophthalmia neonatorum	Male (all)	0	0	0	0	1	1	0	2	1	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	1	3	70	0	0	1	1	1
Total	0	0	1	3	71	1	0	3	2	1		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	175	239	291	249	239	269	301	345	318	310
		-of which MSM	41	46	51	37	68	56	57	83	42	98
		Female	148	179	205	223	224	228	236	274	260	240
		Total	323	418	496	472	463	497	537	619	579	551
B5	Complicated gonococcal infection - including PID and epididymitis	Male (all)	7	13	10	16	4	3	3	6	6	5
		-of which MSM	2	2	1	2	0	1	0	0	0	0
		Female	15	26	13	15	10	17	8	16	16	9
		Total	22	39	23	31	14	20	11	22	22	14

South West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	1	0	3	2	12	2	12	21	6	13
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	3	8	14	6	7	5	13	8	8	10
		Total	4	8	17	8	19	7	25	29	14	23
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	2,298	2,517	2,589	3,085	3,305	4,273	4,590	4,468	3,783	3,732
		-of which MSM	57	61	54	68	79	101	133	210	199	291
		Female	2,775	2,964	3,134	3,330	3,606	3,933	3,962	4,224	3,687	3,384
		Total	5,073	5,481	5,723	6,415	6,911	8,206	8,552	8,692	7,472	7,117
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	51	41	47	35	46	66	64	73	58	60
		-of which MSM	2	0	1	0	2	3	4	2	3	5
		Female	263	266	199	239	203	205	210	228	228	178
		Total	314	307	246	274	249	271	274	301	286	238
C4D	Chlamydial ophthalmia neonatorum	Male (all)	0	0	1	1	1	2	2	4	3	2
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	1	1	0	0	0	1	2	2	3
		Total	0	1	2	1	1	2	3	6	5	5
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	1,413	1,782	1,835	1,842	2,160	2,036	2,152	2,373	2,273	2,077
		-of which MSM	35	55	62	52	80	82	93	152	124	190
		Female	1,293	1,502	1,419	1,386	1,584	1,522	1,634	1,826	1,599	1,379
		Total	2,706	3,284	3,254	3,228	3,744	3,558	3,786	4,199	3,873	3,458
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	4,562	5,091	5,194	5,777	5,771	5,252	5,012	5,211	3,728	3,754
		-of which MSM	206	217	233	238	247	214	243	276	180	265
		Female	0	0	850	1,177	1,211	1,149	1,260	1,327	959	990
		Total	4,562	5,091	6,044	6,954	6,982	6,401	6,272	6,538	4,687	4,745
C4I	Epidemiological treatment of NSGI	Male (all)	373	511	433	464	578	597	713	742	666	671
		-of which MSM	18	21	26	32	43	29	34	53	35	42
		Female	1,404	1,595	1,162	1,173	1,269	1,161	1,095	1,155	1,037	835
		Total	1,777	2,106	1,595	1,637	1,847	1,758	1,808	1,897	1,703	1,506
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	418	443	491	403	514	572	582	660	583	578
		-of which MSM	3	6	10	15	13	16	32	31	30	31
		Female	985	1,061	787	1,044	1,203	1,197	1,252	1,351	1,214	1,233
		Total	1,403	1,504	1,278	1,447	1,717	1,769	1,834	2,011	1,797	1,811
C6A	Trichomoniasis	Male (all)	21	20	12	13	12	26	18	18	16	11
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	275	234	238	210	195	180	226	197	235	232
		Total	296	254	250	223	207	206	244	215	251	243
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	173	204	203	216	230	206	223	297	280	295
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	4,362	4,686	4,239	4,053	4,099	4,181	4,977	5,333	5,221	5,264
		Total	4,535	4,890	4,442	4,269	4,329	4,387	5,200	5,630	5,501	5,559
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	922	987	982	1,057	1,016	960	1,229	1,262	1,323	1,298
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	243	284	208	228	194	226	338	372	416	423
		Total	1,165	1,271	1,190	1,285	1,210	1,186	1,567	1,634	1,741	1,721

South West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C7A	Anogenital candidosis	Male (all)	763	619	560	501	536	586	572	556	576	538
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	5,714	5,607	5,396	4,757	4,282	4,171	4,580	4,729	4,956	4,975
		Total	6,477	6,226	5,956	5,258	4,818	4,757	5,152	5,285	5,532	5,513
C7B	Epidemiological treatment of C6 & C7	Male (all)	298	463	460	450	387	416	381	366	271	294
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	510	547	479	637	765	790	908	981	733	707
		Total	808	1,010	939	1,087	1,152	1,206	1,289	1,347	1,004	1,001
C8,C9	Scabies/pediculosis pubis	Male (all)	313	310	260	171	153	111	120	146	149	141
		-of which MSM	40	50	42	24	23	23	28	28	40	51
		Female	88	62	34	21	30	32	18	33	20	18
		Total	401	372	294	192	183	143	138	179	169	159
C10A	Anogenital herpes simplex – first episode	Male (all)	612	598	550	549	605	540	716	898	913	966
		-of which MSM	27	14	21	22	25	17	27	38	23	48
		Female	965	942	816	773	830	835	1,042	1,326	1,426	1,495
		Total	1,577	1,540	1,366	1,322	1,435	1,375	1,758	2,224	2,340	2,461
C10B	Anogenital herpes simplex – recurrence	Male (all)	687	719	652	595	611	659	647	796	848	894
		-of which MSM	31	27	24	28	35	23	33	42	36	45
		Female	856	876	850	711	768	801	949	1,025	1,203	1,194
		Total	1,543	1,595	1,502	1,306	1,379	1,460	1,596	1,821	2,051	2,088
C11A	Anogenital warts – first episode	Male (all)	3,057	3,121	3,296	3,449	3,471	3,514	3,588	3,808	3,714	3,803
		-of which MSM	113	91	97	108	122	133	114	161	117	163
		Female	2,865	2,999	3,018	3,038	3,221	3,117	3,173	3,481	3,424	3,281
		Total	5,922	6,120	6,314	6,487	6,692	6,631	6,761	7,289	7,139	7,085
C11B	Anogenital warts – recurrence	Male (all)	2,258	2,242	2,126	2,212	2,193	2,292	2,423	2,490	2,575	2,501
		-of which MSM	85	96	93	87	74	80	86	95	112	132
		Female	1,321	1,373	1,296	1,292	1,287	1,364	1,481	1,683	1,596	1,579
		Total	3,579	3,615	3,422	3,504	3,480	3,656	3,904	4,173	4,171	4,080
C11C	Anogenital warts – re-registered cases	Male (all)	658	708	701	756	822	790	752	658	587	528
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	681	615	614	630	756	692	593	512	464	409
		Total	1,339	1,323	1,315	1,386	1,578	1,482	1,345	1,170	1,051	937
C12	Molluscum contagiosum	Male (all)	465	565	614	706	751	827	842	851	804	730
		-of which MSM	9	6	15	21	14	7	13	17	17	14
		Female	292	299	309	356	381	428	419	381	325	307
		Total	757	864	923	1,062	1,132	1,255	1,261	1,232	1,130	1,037
C13	Antigen positive viral hepatitis B	Male (all)	26	26	-	-	-	-	-	-	-	-
		-of which MSM	6	4	-	-	-	-	-	-	-	-
		Female	4	13	-	-	-	-	-	-	-	-
		Total	30	39	-	-	-	-	-	-	-	-
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	25	30	38	19	26	28	30	27
		-of which MSM	-	-	2	3	3	1	0	3	0	4
		Female	-	-	4	6	10	9	13	8	8	10
		Total	-	-	29	36	48	28	39	36	38	37

South West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	3	1	3	2	2	6	4	2
		-of which MSM	-	-	0	1	2	0	1	3	0	0
		Female	-	-	0	0	2	1	3	1	0	2
		Total	-	-	3	1	5	3	5	7	4	4
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	15	2	16	16	13	22	25	9
		-of which MSM	-	-	5	0	1	3	4	2	4	1
		Female	-	-	7	2	4	9	3	4	6	8
		Total	-	-	22	4	20	25	16	26	31	17
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	68	63	93	90	85	55	79	54	54	31
		-of which MSM	2	5	4	2	1	0	3	3	3	1
		Female	26	29	32	32	38	30	11	27	21	16
		Total	94	92	125	122	123	85	90	81	75	47
E1A	New HIV diagnosis - asymptomatic	Male (all)	69	85	102	95	132	127	168	141	103	96
		-of which MSM	24	34	39	41	57	41	62	63	53	50
		Female	28	31	55	61	64	72	60	63	40	32
		Total	97	116	157	156	196	199	228	204	143	128
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	546	597	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	136	151	-	-	-	-	-	-	-	-
		Total	682	748	-	-	-	-	-	-	-	-
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	1,049	1,294	1,494	1,668	1,818	1,836	1,988	1,925
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	333	506	619	691	761	823	827	743
		Total	-	-	1,382	1,800	2,113	2,359	2,579	2,659	2,815	2,668
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	31	30	24	26	36	28	25	14	23	15
		-of which MSM	14	15	4	16	18	10	4	3	12	5
		Female	11	12	4	14	8	16	6	7	7	7
		Total	42	42	28	40	44	44	31	21	30	22
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	413	431	-	-	-	-	-	-	-	-
		-of which MSM	0	0	-	-	-	-	-	-	-	-
		Female	81	124	-	-	-	-	-	-	-	-
		Total	494	555	-	-	-	-	-	-	-	-
E3A*	AIDS – first presentation	Male (all)	27	29	-	-	-	-	-	-	-	-
		-of which MSM	10	13	-	-	-	-	-	-	-	-
		Female	2	4	-	-	-	-	-	-	-	-
		Total	29	33	-	-	-	-	-	-	-	-
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	11	15	12	9	11	12	21	13
		-of which MSM	-	-	4	5	7	1	2	1	8	3
		Female	-	-	7	1	8	3	4	7	2	6
		Total	-	-	18	16	20	12	15	19	23	19
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	11	17	8	9	5	3	2	6
		-of which MSM	-	-	3	5	0	4	2	3	0	3
		Female	-	-	0	1	4	0	0	3	0	1
		Total	-	-	11	18	12	9	5	6	2	7

South West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	269	303	349	463	552	618	612	558	584	594	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	33	31	75	95	100	116	127	140	158	152	
		Total	302	334	424	558	652	734	739	698	742	746	
D2A	Urinary tract infection	Male (all)	129	116	86	87	108	101	109	131	95	106	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	481	421	348	348	396	530	617	953	1,010	999	
		Total	610	537	434	435	504	631	726	1,084	1,105	1,107	
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	3,231	3,415	3,798	3,899	4,247	4,350	4,742	5,607	6,430	6,486	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	4,295	4,475	4,655	5,074	5,278	5,412	5,679	6,513	7,407	7,650	
		Total	7,526	7,890	8,453	8,973	9,525	9,762	10,421	12,120	13,840	14,137	
D3	Other episodes not requiring treatment	Male (all)	8,670	10,181	9,679	13,722	17,297	17,652	19,997	21,191	20,762	20,853	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	8,990	10,389	9,006	13,327	17,424	17,094	20,439	23,126	22,337	21,870	
		Total	17,660	20,570	18,685	27,049	34,721	34,746	40,436	44,317	43,113	42,731	
P1A	HIV antibody counselling - with testing	Male (all)	7,172	10,112	3,485	3,238	2,647	2,133	1,994	1,849	1,954	1,947	
		-of which MSM	741	827	357	364	321	330	270	307	261	376	
		Female	6,292	8,558	2,347	2,180	2,025	1,884	1,765	1,903	1,870	1,809	
		Total	13,464	18,670	5,832	5,418	4,672	4,017	3,759	3,752	3,824	3,756	
P1B	HIV antibody counselling - without testing	Male (all)	1,383	2,720	7,754	9,211	9,631	8,809	9,445	9,698	9,181	9,099	
		-of which MSM	53	91	246	307	284	277	327	366	329	416	
		Female	1,230	2,780	8,070	9,903	10,937	11,235	12,532	13,460	14,307	16,407	
		Total	2,613	5,500	15,824	19,114	20,568	20,044	21,977	23,158	23,494	25,507	
P2	Hepatitis B vaccination	Male (all)	755	859	1,066	972	996	1,184	1,161	1,625	1,520	1,536	
		-of which MSM	264	324	357	339	409	501	471	659	453	727	
		Female	290	291	354	358	433	415	517	666	749	728	
		Total	1,045	1,150	1,420	1,330	1,429	1,599	1,678	2,291	2,270	2,264	
P3	Family Planning	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	1,900	1,905	2,447	2,370	2,415	4,258	5,336	8,118	8,679	9,538	
		Total	1,900	1,905	2,447	2,370	2,415	4,258	5,336	8,118	8,679	9,538	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	280	165	119	87	41	43	41	35	90	44	
		Total	280	165	119	87	41	43	41	35	90	44	

South West SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	48	21	28	20	10	8	8	10	5	4
		Total	48	21	28	20	10	8	8	10	5	4
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	11,553	11,384	11,455	11,268	11,304	11,116	9,609	9,516
		-of which MSM	-	-	433	417	440	447	472	520	410	555
		Female	-	-	13,929	13,086	13,380	14,055	14,641	15,988	15,313	15,406
		Total	-	-	25,482	24,470	24,835	25,323	25,945	27,104	24,930	24,925
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	11,207	15,079	18,856	23,490	28,486	32,766	33,196	33,573
		-of which MSM	-	-	643	814	1,019	1,328	1,733	2,329	2,131	3,150
		Female	-	-	10,066	13,368	17,610	21,572	27,127	31,896	32,865	33,061
		Total	-	-	21,273	28,447	36,466	45,062	55,613	64,662	66,077	66,644
Total new STI diagnoses (for GUM clinics only)		Male (all)	12,736	13,920	14,215	15,215	15,618	16,130	16,424	17,006	14,570	14,480
		-of which MSM	651	669	696	750	784	776	861	1,071	859	1,160
		Female	8,885	9,295	9,925	10,633	11,291	11,505	11,940	12,924	11,862	11,377
		Total	21,621	23,215	24,140	25,848	26,909	27,635	28,364	29,930	26,437	25,860
Total other STI diagnoses		Male (all)	6,847	7,587	7,482	7,944	8,735	9,012	9,536	9,938	9,933	9,569
		-of which MSM	223	262	266	250	325	298	355	496	380	535
		Female	5,980	6,467	5,969	6,044	6,718	6,603	6,905	7,487	7,180	6,569
		Total	12,827	14,054	13,451	13,988	15,453	15,615	16,441	17,425	17,115	16,141
Total other GUM clinic diagnoses		Male (all)	5,610	5,893	6,225	6,333	6,666	6,711	7,376	8,329	9,088	9,086
		-of which MSM	8	9	11	6	7	4	8	11	7	6
		Female	15,963	16,248	15,515	15,244	15,119	15,410	17,178	18,966	19,873	20,102
		Total	21,573	22,141	21,740	21,577	21,785	22,121	24,554	27,295	28,966	29,191
Total services provided		Male (all)	17,980	23,872	44,744	53,606	60,882	64,536	72,387	78,245	76,222	76,524
		-of which MSM	1,058	1,242	2,036	2,241	2,473	2,883	3,273	4,181	3,584	5,224
		Female	18,702	23,923	46,219	54,592	64,224	70,513	82,357	95,157	96,120	98,819
		Total	36,682	47,795	90,963	108,198	125,106	135,049	154,744	173,402	172,387	175,365

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 j: West Midlands SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	40	51	85	168	308	293	249	181	164	132
		-of which MSM	13	21	36	50	108	104	86	80	74	61
		Female	14	5	11	52	69	71	65	47	28	16
		Total	54	56	96	220	377	364	314	228	192	148
A3	Early latent syphilis (first 2 years)	Male (all)	5	14	20	43	51	48	45	50	36	27
		-of which MSM	1	10	6	19	15	16	11	18	10	11
		Female	4	4	9	20	26	30	25	26	22	11
		Total	9	18	29	63	77	78	70	76	58	38
A4,A5,A6	Other acquired syphilis	Male (all)	56	78	85	76	72	70	59	53	96	97
		-of which MSM	3	14	7	5	9	18	2	8	9	28
		Female	38	65	80	53	72	48	73	66	61	62
		Total	94	143	165	129	144	118	132	119	157	159
A7	Congenital syphilis, aged under 2	Male (all)	0	0	0	0	2	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	1	0	1	2	0	0	0
Total	0	0	0	1	2	1	2	0	0	0		
A8	Congenital syphilis, aged 2 or over	Male (all)	0	0	1	1	2	0	1	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	1	1	0	0	0	0	0
Total	0	0	1	2	3	0	1	0	0	0		
A9	Epidemiological treatment of suspected syphilis	Male (all)	6	2	3	12	43	46	38	43	58	67
		-of which MSM	1	0	1	3	11	22	14	20	20	29
		Female	5	1	3	9	14	20	23	19	25	16
		Total	11	3	6	21	57	66	61	62	83	83
B1,B2	Uncomplicated gonorrhoea	Male (all)	1,821	1,689	1,748	1,451	1,139	1,101	1,093	940	972	1,080
		-of which MSM	161	163	216	148	159	151	130	116	114	212
		Female	940	858	856	736	589	597	673	625	673	563
		Total	2,761	2,547	2,604	2,187	1,728	1,698	1,766	1,565	1,645	1,643
B3	Gonococcal ophthalmia neonatorum	Male (all)	0	0	1	1	1	2	0	0	2	2
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	2	0	1	1	0	2	0	0	0	2
Total	2	0	2	2	1	4	0	0	2	4		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	186	200	220	214	214	273	332	312	371	371
		-of which MSM	26	43	49	42	51	57	38	60	38	92
		Female	274	295	285	307	321	290	309	319	326	303
		Total	460	495	505	521	535	563	641	631	697	674
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	6	8	10	22	11	9	8	8	12	18
		-of which MSM	1	1	1	2	2	1	0	0	1	3
		Female	39	18	44	43	37	36	28	28	25	30
		Total	45	26	54	65	48	45	36	36	37	48

West Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	1	2	4	9	5	14	7	11	11	15	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	0	0	6	8	6	11	14	14	7	4	1
		Total	1	2	10	17	11	25	21	18	15	16	
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	3,043	3,432	3,568	3,447	4,032	4,384	4,755	4,781	5,005	4,635	
		-of which MSM	54	64	85	76	78	114	139	195	187	275	
		Female	3,532	4,026	4,344	4,167	4,328	4,298	4,667	5,271	5,190	4,464	
		Total	6,575	7,458	7,912	7,614	8,360	8,682	9,422	10,052	10,197	9,100	
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	24	39	48	68	61	102	108	67	72	55	
		-of which MSM	0	0	3	0	1	6	5	4	2	2	
		Female	108	152	163	199	181	255	253	198	213	159	
		Total	132	191	211	267	242	357	361	265	285	214	
C4D	Chlamydial ophthalmia neonatorum	Male (all)	0	2	5	1	2	2	0	0	2	0	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	2	5	2	3	0	1	1	1	1	1	
		Total	2	7	7	4	2	3	1	1	3	1	
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	1,529	1,794	1,650	1,822	1,856	1,931	2,153	2,333	2,989	2,805	
		-of which MSM	50	60	55	34	43	60	57	111	105	154	
		Female	941	1,085	1,013	1,228	1,426	1,567	1,622	1,886	2,235	2,095	
		Total	2,470	2,879	2,663	3,050	3,282	3,498	3,775	4,219	5,225	4,900	
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	3,633	3,954	4,497	4,701	4,941	4,565	4,393	5,212	4,271	3,580	
		-of which MSM	131	139	194	172	204	187	221	250	238	156	
		Female	0	0	97	303	416	410	494	404	414	405	
		Total	3,633	3,954	4,594	5,004	5,357	4,975	4,887	5,616	4,685	3,985	
C4I	Epidemiological treatment of NSGI	Male (all)	288	203	262	332	299	281	341	324	332	264	
		-of which MSM	20	13	17	21	26	21	23	31	28	22	
		Female	1,365	1,386	1,526	1,465	1,405	1,199	1,083	1,307	1,221	1,134	
		Total	1,653	1,589	1,788	1,797	1,704	1,480	1,424	1,631	1,553	1,398	
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	166	164	177	228	250	270	363	381	481	479	
		-of which MSM	5	6	4	6	11	15	10	19	21	23	
		Female	552	652	651	791	768	954	1,274	1,554	1,748	1,632	
		Total	718	816	828	1,019	1,018	1,224	1,637	1,935	2,229	2,111	
C6A	Trichomoniasis	Male (all)	63	61	31	40	35	50	59	30	28	29	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	781	849	750	567	641	700	621	745	717	702	
		Total	844	910	781	607	676	750	680	775	745	731	
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	66	76	109	107	111	127	121	166	97	115	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	5,158	5,478	5,324	5,425	5,505	5,708	6,792	8,132	8,502	9,088	
		Total	5,224	5,554	5,433	5,532	5,616	5,835	6,913	8,298	8,599	9,203	
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	1,212	1,115	1,084	1,102	994	1,010	1,105	1,167	1,152	965	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	173	171	279	225	184	192	177	273	180	191	
		Total	1,385	1,286	1,363	1,327	1,178	1,202	1,282	1,440	1,332	1,156	

West Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C7A	Anogenital candidosis	Male (all)	433	446	364	380	385	476	493	536	492	475	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	5,679	6,036	5,704	5,356	5,646	5,263	5,798	6,324	6,494	6,550	
		Total	6,112	6,482	6,068	5,736	6,031	5,739	6,291	6,860	6,987	7,025	
C7B	Epidemiological treatment of C6 & C7	Male (all)	339	382	450	402	449	451	601	579	442	525	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	579	826	760	845	788	806	922	1,135	873	833	
		Total	918	1,208	1,210	1,247	1,237	1,257	1,523	1,714	1,315	1,358	
C8,C9	Scabies/pediculosis pubis	Male (all)	352	289	222	158	153	139	160	186	141	138	
		-of which MSM	34	36	19	18	23	22	15	34	17	31	
		Female	89	65	39	30	17	19	23	25	20	29	
		Total	441	354	261	188	170	158	183	211	161	167	
C10A	Anogenital herpes simplex – first episode	Male (all)	611	611	540	579	615	616	827	924	999	1,031	
		-of which MSM	16	16	22	15	31	23	30	47	38	47	
		Female	979	1,100	994	949	985	1,037	1,212	1,502	1,541	1,698	
		Total	1,590	1,711	1,534	1,528	1,600	1,653	2,039	2,426	2,541	2,729	
C10B	Anogenital herpes simplex – recurrence	Male (all)	373	411	382	447	509	524	549	681	730	762	
		-of which MSM	10	8	17	14	24	16	27	23	21	30	
		Female	457	579	608	627	601	667	811	1,008	1,064	1,221	
		Total	830	990	990	1,074	1,110	1,191	1,360	1,689	1,794	1,983	
C11A	Anogenital warts – first episode	Male (all)	2,861	2,841	2,856	3,150	3,099	3,174	3,474	3,812	3,828	3,797	
		-of which MSM	86	101	121	100	130	126	119	144	127	146	
		Female	2,742	2,548	2,646	2,716	2,845	2,878	3,213	3,570	3,385	3,337	
		Total	5,603	5,389	5,502	5,866	5,944	6,052	6,687	7,382	7,214	7,134	
C11B	Anogenital warts – recurrence	Male (all)	1,456	1,384	1,400	1,507	1,523	1,642	1,763	2,113	1,637	1,543	
		-of which MSM	34	39	60	53	55	63	57	82	46	64	
		Female	899	826	824	877	982	1,068	1,121	1,352	1,187	1,076	
		Total	2,355	2,210	2,224	2,384	2,505	2,710	2,884	3,465	2,824	2,619	
C11C	Anogenital warts – re-registered cases	Male (all)	759	612	562	579	532	536	573	613	509	521	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	518	450	449	409	445	497	439	447	397	357	
		Total	1,277	1,062	1,011	988	977	1,033	1,012	1,060	906	878	
C12	Molluscum contagiosum	Male (all)	358	411	488	563	575	672	779	844	775	762	
		-of which MSM	6	7	9	12	12	15	15	10	10	15	
		Female	203	217	235	276	262	318	313	338	295	337	
		Total	561	628	723	839	837	990	1,092	1,182	1,070	1,100	
C13	Antigen positive viral hepatitis B	Male (all)	12	42	-	-	-	-	-	-	-	-	
		-of which MSM	2	6	-	-	-	-	-	-	-	-	
		Female	17	33	-	-	-	-	-	-	-	-	
		Total	29	75	-	-	-	-	-	-	-	-	
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	49	39	41	38	33	26	54	45	
		-of which MSM	-	-	4	2	4	5	3	1	4	4	
		Female	-	-	8	15	25	15	15	14	15	23	
		Total	-	-	57	54	66	53	48	40	69	68	

West Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	2	3	3	2	2	1	2	2
		-of which MSM	-	-	0	1	0	0	0	1	0	0
		Female	-	-	0	0	0	0	2	1	1	1
		Total	-	-	2	3	3	2	4	2	3	3
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	18	18	16	9	14	24	22	16
		-of which MSM	-	-	7	2	4	3	5	8	4	6
		Female	-	-	7	4	7	10	5	8	5	5
		Total	-	-	25	22	23	19	19	32	27	21
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	35	54	45	54	45	37	42	54	51	51
		-of which MSM	1	4	3	3	1	1	1	3	6	7
		Female	19	32	20	29	26	26	14	25	28	24
		Total	54	86	65	83	71	63	56	79	79	75
E1A	New HIV diagnosis - asymptomatic	Male (all)	76	134	160	138	163	151	157	190	166	184
		-of which MSM	24	32	62	37	68	72	55	72	45	67
		Female	52	138	142	174	118	87	74	113	102	86
		Total	128	272	302	312	281	238	231	303	268	270
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	230	381	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	104	207	-	-	-	-	-	-	-	-
		Total	334	588	-	-	-	-	-	-	-	-
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	621	686	1,072	1,603	1,419	1,869	1,444	1,613
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	302	374	653	931	863	1,325	1,012	1,269
		Total	-	-	923	1,060	1,725	2,534	2,282	3,194	2,456	2,882
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	26	50	20	32	30	16	17	30	11	29
		-of which MSM	7	15	5	8	12	8	6	8	5	12
		Female	10	20	15	16	16	9	4	12	17	11
		Total	36	70	35	48	46	25	21	42	28	40
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	334	315	-	-	-	-	-	-	-	-
		-of which MSM	0	0	-	-	-	-	-	-	-	-
		Female	108	118	-	-	-	-	-	-	-	-
		Total	442	433	-	-	-	-	-	-	-	-
E3A*	AIDS – first presentation	Male (all)	36	32	-	-	-	-	-	-	-	-
		-of which MSM	16	8	-	-	-	-	-	-	-	-
		Female	7	7	-	-	-	-	-	-	-	-
		Total	43	39	-	-	-	-	-	-	-	-
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	11	13	13	7	18	17	17	17
		-of which MSM	-	-	1	3	2	0	5	4	2	5
		Female	-	-	9	5	11	3	4	6	7	10
		Total	-	-	20	18	24	10	22	23	24	27
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	1	4	7	8	8	4	1	5
		-of which MSM	-	-	1	2	3	3	4	2	0	1
		Female	-	-	1	1	4	5	0	1	0	1
		Total	-	-	2	5	11	13	8	5	1	6

West Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	139	154	204	269	323	433	316	462	411	507	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	58	53	56	110	184	204	190	234	219	317	
		Total	197	207	260	379	507	637	506	696	630	824	
D2A	Urinary tract infection	Male (all)	124	135	97	118	92	112	137	192	237	143	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	500	539	530	535	529	659	869	1,108	1,052	1,028	
		Total	624	674	627	653	621	771	1,006	1,300	1,289	1,171	
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	2,177	2,333	2,511	2,844	3,145	3,351	3,995	4,969	5,447	5,557	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	2,246	2,081	1,910	2,035	2,136	2,449	3,133	3,881	4,416	4,456	
		Total	4,423	4,414	4,421	4,879	5,281	5,800	7,128	8,850	9,863	10,013	
D3	Other episodes not requiring treatment	Male (all)	8,421	8,727	8,167	9,364	10,904	13,645	17,249	22,189	23,239	23,256	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	8,336	8,617	7,794	9,263	10,602	13,488	17,290	23,024	25,466	25,280	
		Total	16,757	17,344	15,961	18,627	21,506	27,133	34,539	45,213	48,709	48,538	
P1A	HIV antibody counselling - with testing	Male (all)	8,591	11,303	3,819	2,980	2,979	2,825	2,633	2,715	1,918	2,065	
		-of which MSM	813	986	449	365	488	481	358	324	284	291	
		Female	7,400	9,874	2,781	2,018	1,972	1,989	2,312	2,508	2,020	2,000	
		Total	15,991	21,177	6,600	4,998	4,951	4,814	4,945	5,223	3,938	4,066	
P1B	HIV antibody counselling - without testing	Male (all)	578	2,915	7,585	8,947	7,977	8,044	8,753	8,979	9,594	8,627	
		-of which MSM	45	135	293	234	249	250	268	263	229	295	
		Female	561	3,216	8,301	9,921	9,190	8,857	10,272	11,883	12,971	12,769	
		Total	1,139	6,131	15,886	18,868	17,167	16,901	19,025	20,862	22,569	21,397	
P2	Hepatitis B vaccination	Male (all)	717	948	1,077	1,015	978	1,091	1,415	1,765	1,398	1,325	
		-of which MSM	305	392	577	458	496	489	579	793	573	580	
		Female	205	316	287	389	365	436	443	638	524	592	
		Total	922	1,264	1,364	1,404	1,343	1,527	1,858	2,403	1,922	1,918	
P3	Family Planning	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	688	722	1,012	1,404	1,151	993	1,124	1,543	2,015	2,992	
		Total	688	722	1,012	1,404	1,151	993	1,124	1,543	2,015	2,992	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	195	124	200	149	150	133	136	222	216	102	
		Total	195	124	200	149	150	133	136	222	216	102	

West Midlands SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	38	22	34	30	32	15	26	36	41	39
		Total	38	22	34	30	32	15	26	36	41	39
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	12,228	11,411	10,141	9,151	9,481	9,546	9,355	8,424
		-of which MSM	-	-	404	327	354	335	401	457	315	412
		Female	-	-	14,536	13,329	12,304	11,090	11,882	13,834	13,314	13,156
		Total	-	-	26,764	24,740	22,445	20,241	21,363	23,380	22,671	21,582
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	12,795	16,776	19,856	25,487	30,957	37,444	40,267	40,692
		-of which MSM	-	-	945	975	1,109	1,362	1,657	2,408	2,295	2,957
		Female	-	-	12,309	16,405	19,674	26,482	31,998	40,245	43,851	43,623
		Total	-	-	25,104	33,181	39,530	51,969	62,955	77,689	84,124	84,321
Total new STI diagnoses (for GUM clinics only)	Male (all)	13,086	13,750	14,485	14,810	15,481	15,611	16,512	17,664	16,989	16,008	
	-of which MSM	539	611	784	666	856	860	847	1,001	891	1,066	
	Female	10,045	10,652	11,011	11,052	11,315	11,713	12,957	14,471	14,401	13,491	
	Total	23,131	24,402	25,496	25,862	26,796	27,324	29,469	32,135	31,394	29,501	
Total other STI diagnoses	Male (all)	5,392	5,568	5,397	5,951	6,457	7,351	7,552	8,807	8,582	8,557	
	-of which MSM	160	185	207	174	222	260	222	337	267	420	
	Female	4,778	5,077	5,150	5,466	6,108	6,500	6,537	7,965	7,748	7,854	
	Total	10,170	10,645	10,547	11,417	12,565	13,851	14,089	16,772	16,331	16,411	
Total other GUM clinic diagnoses	Male (all)	4,398	4,583	4,729	5,067	5,281	5,613	6,543	7,714	7,996	7,894	
	-of which MSM	3	10	14	8	9	9	9	13	14	17	
	Female	14,604	15,342	14,776	14,648	15,028	15,276	17,889	21,159	21,823	22,340	
	Total	19,002	19,925	19,505	19,715	20,309	20,889	24,432	28,873	29,820	30,234	
Total services provided	Male (all)	18,307	23,893	45,671	50,493	52,835	60,243	70,488	82,638	85,771	84,389	
	-of which MSM	1,163	1,513	2,668	2,359	2,696	2,917	3,263	4,245	3,696	4,535	
	Female	17,190	22,745	47,020	52,729	55,258	63,335	75,321	93,675	100,161	100,412	
	Total	35,497	46,638	92,691	103,222	108,093	123,578	145,809	176,313	185,948	184,814	

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months



3 k: Yorkshire & the Humber SHA

Number of new STI episodes seen at genitourinary medicine clinics by sex: 2001 - 2010

Diagnoses and/or treatment of infection or disease			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
A1,A2	Primary and secondary infectious syphilis	Male (all)	28	53	52	68	103	137	105	150	168	112
		-of which MSM	10	16	21	28	48	69	54	66	111	62
		Female	6	11	8	15	17	10	16	13	19	20
		Total	34	64	60	83	120	147	121	163	187	132
A3	Early latent syphilis (first 2 years)	Male (all)	17	20	16	15	26	38	33	36	37	33
		-of which MSM	4	10	6	5	5	17	12	14	21	21
		Female	10	5	7	12	21	12	8	14	11	12
		Total	27	25	23	27	47	50	41	50	48	45
A4,A5,A6	Other acquired syphilis	Male (all)	30	26	47	59	58	55	52	42	61	49
		-of which MSM	2	3	4	4	5	6	11	4	20	10
		Female	19	22	31	54	46	31	24	40	35	37
		Total	49	48	78	113	104	86	76	82	96	86
A7	Congenital syphilis, aged under 2	Male (all)	1	0	1	0	0	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	0	0	0	0	0
Total	1	0	1	0	0	0	0	0	0	0		
A8	Congenital syphilis, aged 2 or over	Male (all)	0	0	0	0	1	0	1	0	1	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	0	2	0	1	0	0	0	0	0
Total	1	0	2	0	2	0	1	0	1	0		
A9	Epidemiological treatment of suspected syphilis	Male (all)	2	10	6	11	19	14	11	16	19	25
		-of which MSM	0	8	4	5	11	11	9	8	10	15
		Female	1	3	4	1	4	2	8	6	6	10
		Total	3	13	10	12	23	16	19	22	25	35
B1,B2	Uncomplicated gonorrhoea	Male (all)	1,007	1,409	1,510	1,417	1,171	1,073	949	756	812	819
		-of which MSM	117	146	147	138	141	180	149	139	170	201
		Female	495	668	740	784	611	551	560	497	516	518
		Total	1,502	2,077	2,250	2,201	1,782	1,624	1,509	1,253	1,328	1,337
B3	Gonococcal ophthalmia neonatorum	Male (all)	0	0	0	0	1	2	2	3	4	2
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	1	1	3	0	1	5	2	5
Total	0	0	1	1	4	2	3	8	6	7		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	126	154	180	224	244	270	293	282	334	323
		-of which MSM	23	25	19	27	29	30	45	37	63	66
		Female	165	239	279	258	253	235	241	263	258	242
		Total	291	393	459	482	497	505	534	545	592	565
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	10	8	6	10	13	14	9	9	15	10
		-of which MSM	0	2	0	0	0	2	2	4	2	1
		Female	22	32	30	30	20	46	36	27	40	43
		Total	32	40	36	40	33	60	45	36	55	53

Yorkshire & the Humber SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	3	6	4	7	3	5	3	8	10	9
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	0	6	1	1	2	1	8	8	3
		Total	4	6	10	8	4	7	4	16	18	12
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	3,265	4,209	4,556	5,125	5,055	5,248	5,731	6,012	5,230	4,710
		-of which MSM	42	62	79	73	101	131	159	226	291	282
		Female	4,496	5,435	5,763	6,263	5,955	5,548	6,101	6,186	5,319	4,566
		Total	7,761	9,644	10,319	11,388	11,010	10,796	11,832	12,198	10,552	9,277
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	89	87	88	102	84	84	90	126	105	98
		-of which MSM	1	2	2	1	6	2	3	4	5	2
		Female	246	202	226	279	281	297	341	389	312	327
		Total	335	289	314	381	365	381	431	515	417	425
C4D	Chlamydial ophthalmia neonatorum	Male (all)	3	2	2	1	1	0	4	1	2	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1	1	0	0	3	1	1	3	0	0
		Total	4	3	2	1	4	1	5	4	2	0
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	1,895	2,034	2,026	2,298	2,608	2,332	2,607	2,982	2,862	2,381
		-of which MSM	17	10	28	24	47	50	83	82	146	129
		Female	1,053	1,149	1,067	1,333	1,442	1,461	1,725	2,036	1,965	1,492
		Total	2,948	3,183	3,093	3,631	4,050	3,793	4,332	5,018	4,828	3,876
C4H*	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	3,714	3,700	4,292	4,035	3,500	3,288	3,260	3,351	2,943	2,713
		-of which MSM	97	118	115	90	94	104	92	144	157	186
		Female	0	0	287	319	435	442	558	831	832	974
		Total	3,714	3,700	4,579	4,354	3,935	3,730	3,818	4,182	3,776	3,688
C4I	Epidemiological treatment of NSGI	Male (all)	379	370	394	327	360	345	380	411	429	490
		-of which MSM	14	14	18	16	12	18	11	20	22	25
		Female	1,270	1,242	1,148	925	872	824	824	829	618	514
		Total	1,649	1,612	1,542	1,252	1,232	1,169	1,204	1,240	1,047	1,004
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	384	363	386	439	513	431	539	580	603	562
		-of which MSM	13	12	13	14	21	13	18	23	34	41
		Female	719	667	722	833	885	1,045	1,236	1,353	1,148	1,365
		Total	1,103	1,030	1,108	1,272	1,398	1,476	1,775	1,933	1,751	1,927
C6A	Trichomoniasis	Male (all)	13	12	7	19	10	8	18	17	27	24
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	492	499	462	426	362	384	341	408	372	443
		Total	505	511	469	445	372	392	359	425	399	467
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	193	202	194	207	177	266	252	319	289	336
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	5,665	6,280	6,615	5,802	5,386	5,758	6,880	8,251	8,398	8,363
		Total	5,858	6,482	6,809	6,009	5,563	6,024	7,132	8,570	8,688	8,700
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	996	996	836	888	733	648	808	947	834	876
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	327	201	135	585	569	129	91	134	88	184
		Total	1,323	1,197	971	1,473	1,302	777	899	1,081	922	1,060

Yorkshire & the Humber SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
C7A	Anogenital candidosis	Male (all)	452	400	279	274	247	316	444	551	660	627	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	5,385	5,312	4,723	4,060	3,972	3,841	4,046	5,022	5,370	5,928	
		Total	5,837	5,712	5,002	4,334	4,219	4,157	4,490	5,573	6,032	6,556	
C7B	Epidemiological treatment of C6 & C7	Male (all)	330	328	310	341	390	374	392	583	624	464	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	669	682	736	607	655	550	765	862	639	352	
		Total	999	1,010	1,046	948	1,045	924	1,157	1,445	1,264	816	
C8,C9	Scabies/pediculosis pubis	Male (all)	436	322	286	208	163	140	180	143	158	159	
		-of which MSM	56	55	24	32	23	25	29	26	31	33	
		Female	75	57	44	32	25	14	15	13	28	27	
		Total	511	379	330	240	188	154	195	156	187	186	
C10A	Anogenital herpes simplex – first episode	Male (all)	550	569	550	531	554	747	921	1,050	1,061	1,087	
		-of which MSM	9	22	8	12	22	26	25	31	43	55	
		Female	956	1,070	1,029	944	966	1,155	1,467	1,656	1,576	1,700	
		Total	1,506	1,639	1,579	1,475	1,520	1,902	2,388	2,706	2,637	2,787	
C10B	Anogenital herpes simplex – recurrence	Male (all)	444	433	490	407	438	493	577	670	720	733	
		-of which MSM	19	7	16	12	10	18	10	18	29	28	
		Female	563	570	499	531	594	615	773	941	1,056	1,097	
		Total	1,007	1,003	989	938	1,032	1,108	1,350	1,611	1,778	1,831	
C11A	Anogenital warts – first episode	Male (all)	3,170	3,265	3,500	3,671	3,587	3,649	4,171	4,464	4,340	4,272	
		-of which MSM	75	100	88	80	94	87	102	133	160	177	
		Female	3,121	3,145	3,253	3,330	3,079	3,404	3,815	4,161	3,768	3,534	
		Total	6,291	6,410	6,753	7,001	6,666	7,053	7,986	8,625	8,108	7,807	
C11B	Anogenital warts – recurrence	Male (all)	2,034	1,926	1,930	1,896	1,903	1,927	2,386	2,832	2,238	2,179	
		-of which MSM	47	49	65	48	44	50	52	89	78	98	
		Female	1,027	937	1,055	1,092	1,091	1,138	1,422	1,800	1,458	1,242	
		Total	3,061	2,863	2,985	2,988	2,994	3,065	3,808	4,632	3,696	3,421	
C11C	Anogenital warts – re-registered cases	Male (all)	789	725	892	970	918	934	925	1,033	819	736	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	712	511	583	633	607	521	696	660	568	501	
		Total	1,501	1,236	1,475	1,603	1,525	1,455	1,621	1,693	1,388	1,237	
C12	Molluscum contagiosum	Male (all)	494	561	637	707	745	713	878	908	889	768	
		-of which MSM	7	2	8	15	12	8	7	8	24	24	
		Female	270	275	325	355	395	383	420	436	383	364	
		Total	764	836	962	1,062	1,140	1,096	1,298	1,344	1,273	1,132	
C13	Antigen positive viral hepatitis B	Male (all)	39	33	-	-	-	-	-	-	-	-	
		-of which MSM	8	4	-	-	-	-	-	-	-	-	
		Female	14	20	-	-	-	-	-	-	-	-	
		Total	53	53	-	-	-	-	-	-	-	-	
C13A*	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	32	32	40	46	25	51	42	51	
		-of which MSM	-	-	3	1	2	2	2	1	6	3	
		Female	-	-	7	9	13	10	10	26	24	16	
		Total	-	-	39	41	53	56	35	77	66	67	

Yorkshire & the Humber SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
C13B*	**number of which were acute viral hepatitis B	Male (all)	-	-	4	3	3	1	7	5	3	1
		-of which MSM	-	-	0	1	0	0	0	0	1	0
		Female	-	-	2	2	2	1	1	3	2	3
		Total	-	-	6	5	5	2	8	8	5	4
C13C*	Viral hepatitis B: subsequent presentation	Male (all)	-	-	10	10	16	11	13	17	12	22
		-of which MSM	-	-	2	0	0	2	4	2	0	14
		Female	-	-	2	7	10	6	4	6	11	11
		Total	-	-	12	17	26	17	17	23	23	33
C14*	Viral hepatitis C: fist diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	71	76	71	78	63	66	44	46	59	61
		-of which MSM	2	2	3	3	1	3	4	3	7	8
		Female	39	61	38	54	23	29	27	34	31	25
		Total	110	137	109	132	86	95	71	80	90	86
E1A	New HIV diagnosis - asymptomatic	Male (all)	61	91	102	133	137	130	142	138	125	131
		-of which MSM	19	18	19	34	31	40	53	54	55	56
		Female	50	77	136	120	133	101	85	112	79	70
		Total	111	168	238	253	270	231	227	250	204	201
E1B*	Asymptomatic HIV infection – subsequent presentation	Male (all)	259	378	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	127	237	-	-	-	-	-	-	-	-
		Total	386	615	-	-	-	-	-	-	-	-
E1BE2B*	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	658	1,268	1,631	1,801	1,760	1,745	2,233	2,699
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	405	826	1,127	1,164	1,134	1,340	1,787	1,933
		Total	-	-	1,063	2,094	2,758	2,965	2,894	3,085	4,027	4,637
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	33	36	28	45	56	35	28	50	39	24
		-of which MSM	12	8	5	7	9	9	9	17	24	15
		Female	6	22	30	16	22	19	23	28	11	12
		Total	39	58	58	61	78	54	51	78	50	36
E2B*	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	329	427	-	-	-	-	-	-	-	-
		-of which MSM	0	0	-	-	-	-	-	-	-	-
		Female	57	122	-	-	-	-	-	-	-	-
		Total	386	549	-	-	-	-	-	-	-	-
E3A*	AIDS – first presentation	Male (all)	12	15	-	-	-	-	-	-	-	-
		-of which MSM	5	4	-	-	-	-	-	-	-	-
		Female	11	8	-	-	-	-	-	-	-	-
		Total	23	23	-	-	-	-	-	-	-	-
E3A1*	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	2	13	5	8	21	7	13	13
		-of which MSM	-	-	0	5	2	5	12	2	2	3
		Female	-	-	10	8	9	8	6	9	11	10
		Total	-	-	12	21	14	16	27	16	24	23
E3A2*	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	7	13	0	3	4	4	4	4
		-of which MSM	-	-	2	0	0	1	2	0	2	0
		Female	-	-	3	4	2	5	3	3	1	2
		Total	-	-	10	17	2	8	7	7	5	6

Yorkshire & the Humber SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	
E3B	AIDS – subsequent presentation	Male (all)	170	208	218	346	374	409	441	459	576	623	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	28	50	83	178	186	256	256	278	331	396	
		Total	198	258	301	524	560	665	697	737	907	1,019	
D2A	Urinary tract infection	Male (all)	117	155	104	463	92	113	137	150	175	171	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	458	420	394	806	432	512	672	903	871	898	
		Total	575	575	498	1,269	524	625	809	1,053	1,047	1,069	
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	4,422	4,525	4,606	4,383	4,185	3,961	4,529	5,516	5,456	5,009	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	4,457	4,088	4,180	3,936	3,598	3,567	4,158	4,574	4,333	4,180	
		Total	8,879	8,613	8,786	8,319	7,783	7,528	8,687	10,090	9,792	9,191	
D3	Other episodes not requiring treatment	Male (all)	9,581	11,919	11,802	14,073	16,524	17,585	21,579	24,313	22,384	22,432	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	9,673	11,944	13,562	13,456	15,429	16,618	21,521	24,705	22,030	21,411	
		Total	19,254	23,863	25,364	27,529	31,953	34,203	43,100	49,018	44,428	43,853	
P1A	HIV antibody counselling - with testing	Male (all)	8,426	12,088	2,116	2,179	2,189	1,706	1,617	1,753	1,538	1,491	
		-of which MSM	729	817	226	196	194	147	146	205	264	257	
		Female	7,532	11,181	1,755	1,628	1,949	1,466	1,695	1,809	1,606	1,627	
		Total	15,958	23,269	3,871	3,807	4,138	3,172	3,312	3,562	3,146	3,119	
P1B	HIV antibody counselling - without testing	Male (all)	861	2,449	7,236	8,227	8,256	6,824	8,182	11,241	11,930	11,697	
		-of which MSM	69	107	205	182	210	202	172	253	385	440	
		Female	779	2,525	8,792	9,184	9,002	7,882	10,347	14,187	14,528	14,566	
		Total	1,640	4,974	16,028	17,411	17,258	14,706	18,529	25,428	26,461	26,267	
P2	Hepatitis B vaccination	Male (all)	770	878	1,069	1,150	1,290	1,324	1,440	1,635	1,505	1,488	
		-of which MSM	295	320	377	370	383	394	407	577	548	589	
		Female	271	254	340	359	421	471	569	715	650	636	
		Total	1,041	1,132	1,409	1,509	1,711	1,795	2,009	2,350	2,155	2,125	
P3	Family Planning	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	1,093	1,020	1,033	883	1,250	1,204	1,502	1,710	1,389	1,621	
		Total	1,093	1,020	1,033	883	1,250	1,204	1,502	1,710	1,389	1,621	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	472	383	364	576	225	150	122	137	129	66	
		Total	472	383	364	576	225	150	122	137	129	66	

Yorkshire & the Humber SHA: Number of new STI episodes by sex: 2001 - 2010 (continued)

			2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	95	76	64	105	36	33	26	20	23	16
		Total	95	76	64	105	36	33	26	20	23	16
S1*	Sexual health screen (no HIV antibody test)	Male (all)	-	-	12,587	13,006	12,642	11,102	12,077	11,510	11,093	10,562
		-of which MSM	-	-	290	245	299	306	300	355	529	595
		Female	-	-	16,230	15,000	14,158	12,855	15,251	15,306	14,522	14,048
		Total	-	-	28,817	28,006	26,800	23,957	27,328	26,816	25,619	24,614
S2*	HIV antibody test and sexual health screen	Male (all)	-	-	16,257	19,715	21,743	24,975	30,986	38,350	37,642	35,694
		-of which MSM	-	-	862	854	968	1,173	1,327	2,026	2,516	2,774
		Female	-	-	14,837	17,135	18,440	21,288	28,480	37,055	35,833	33,837
		Total	-	-	31,094	36,850	40,183	46,263	59,466	75,405	73,493	69,544
Total new STI diagnoses (for GUM clinics only)	Male (all)	13,274	14,711	16,022	16,545	15,725	15,748	17,078	17,805	16,575	15,544	
	-of which MSM	462	573	535	534	609	718	726	891	1,130	1,159	
	Female	10,965	12,165	13,078	13,767	13,217	13,421	15,029	16,141	14,433	13,988	
	Total	24,239	26,876	29,100	30,312	28,942	29,169	32,107	33,946	31,014	29,535	
Total other STI diagnoses	Male (all)	6,473	6,708	6,851	7,820	8,556	8,585	9,443	10,480	10,302	10,244	
	-of which MSM	127	120	156	136	158	184	223	258	370	371	
	Female	5,035	5,091	5,160	5,836	6,231	6,253	7,108	8,204	8,085	7,471	
	Total	11,508	11,799	12,011	13,656	14,787	14,838	16,551	18,684	18,398	17,724	
Total other GUM clinic diagnoses	Male (all)	6,620	6,715	6,446	6,679	5,946	5,802	6,651	8,185	8,154	7,618	
	-of which MSM	10	6	8	5	3	7	10	6	14	25	
	Female	17,581	17,523	17,260	16,549	14,921	14,586	16,802	19,972	19,919	20,042	
	Total	24,201	24,238	23,706	23,228	20,867	20,388	23,453	28,157	28,081	27,664	
Total services provided		2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	
	Male (all)	19,638	27,334	51,067	58,350	62,644	63,516	75,881	88,802	86,092	83,364	
	-of which MSM	1,093	1,244	1,960	1,847	2,054	2,222	2,352	3,416	4,242	4,655	
	Total	38,986	54,258	107,616	115,995	123,293	125,300	155,246	184,289	176,691	171,143	

Source: HPA Centre for Infections

- data not available/applicable

* KC60 codes were created/revised in quarter 3, 2002

† previous diagnosis of C11A or C11B treated for more than 3 months

Supplementary Tables of Sexually Transmitted Infections in England



Code Groups
New STI diagnoses
Chlamydial infection (uncomplicated and complicated)
Gonorrhoea (uncomplicated and complicated)
Infectious and early latent syphilis
Genital herpes simplex (first episode)
Genital warts (first episode)
New HIV diagnosis
Non-specific genital infection (uncomplicated and complicated)
Chancroid/lymphogranuloma venereum (LGV)/Donovanosis
Molluscum contagiosum
Trichomoniasis
Scabies
Pediculus pubis
Other STI diagnoses
Congenital and other acquired syphilis
Recurrent genital herpes simplex
Recurrent and re-registered genital warts
Subsequent HIV presentations (including AIDS)
Ophthalmia neonatorum (chlamydial or gonococcal)
Epidemiological treatment of suspected STIs (syphilis, chlamydia, gonorrhoea, non-specific genital infection)
Other diagnoses made at GUM clinics
Viral hepatitis B
Viral hepatitis C
Vaginitis and balanitis (including epidemiological treatment)
Anogenital candidiasis (including epidemiological treatment)
Urinary tract infection
Cervical abnormalities
Other conditions requiring treatment at a GUM clinic
Services provided
HIV antibody test
Sexual health screen
Hepatitis B vaccination
Contraception (excluding condom provision)
Other episode not requiring treatment