

---

## Table 1: Total number of STI diagnoses in genitourinary medicine clinics & community settings in England 2008 - 2010



Data are presented by country, SHA & gender

---

### Notes on using these tables:

Please refer to 'notes' page

### Definitions of data totals:

Total new STI diagnoses: }  
Total other STI diagnoses: } Refer to 'Code Groups' for conditions included

### Data sources for chlamydia diagnoses:

England: GUM clinic KC60 returns (2008)  
GUM clinic GUMCAD returns (2009-2010)  
National Chlamydia Screening Programme (NCSP) returns  
"Non-NCSP / Non-GUM" returns from laboratories

### Data sources for all other STI diagnoses:

England: GUM clinic KC60 returns (2008)  
GUM clinic GUMCAD returns (2009-2010)

### Data supplied by:

Health Protection Agency, Health Protection Services Colindale

---

### Data prepared by:

HIV and Sexually Transmitted Infections Department  
Health Protection Agency  
Health Protection Services Colindale  
61 Colindale Avenue  
London  
NW9 5EQ

Tel. 020-8200-6868

Email: [hiv/sti@hpa.org.uk](mailto:hiv/sti@hpa.org.uk)

Website: [www.hpa.org.uk](http://www.hpa.org.uk)

---

**Table 1: Total number of STI diagnoses in genitourinary medicine clinics & community settings in England 2008 - 2010**



**Data are presented by country & gender**

---

**Data tables**

England	1 a
East Midlands	1 b
East of England	1 c
London	1 d
North East	1 e
North West	1 f
South Central	1 g
South East Coast	1 h
South West	1 i
West Midlands	1 j
Yorkshire & the Humber	1 k

Code Groups: Definitions of conditions presented

---

**Further Information**

**England**

Data on selected diagnoses, 1971 to 2010, are available upon request

HPA Health Protection Services Colindale

Contact: Geraldine Leong

Tel: +44 (0)20 8200 6868

Email: [geraldine.leong@hpa.org.uk](mailto:geraldine.leong@hpa.org.uk)

Web: <http://www.hpa.org.uk>

### General

1. Data presented are compiled from a combination of sources and reflect diagnoses made in GUM clinics and include chlamydia diagnoses made in other community healthcare and non-healthcare settings such as general practice. Numbers will therefore be higher than those reported in national GUM clinic statistics.
2. Data presented are the number of diagnoses reported and not the number of people diagnosed.
3. Data on unknown gender are included in the 'Total' row for each diagnosis
4. Data are presented for England only.

### GUM clinic returns

5. Numbers of diagnoses are adjusted for missing GUM clinic data.
6. GUM clinic chlamydia diagnoses include C4A, C4B & C4C (complicated and uncomplicated chlamydia).
7. An improved STI surveillance system (GUMCAD) was introduced in England in 2009. Unlike the KC60 return, duplicate records in GUMCAD can be identified and removed resulting in a slight reduction in the number of diagnoses reported. England data for 2008 have therefore been statistically adjusted to enable fair comparisons of STI diagnoses over time.

### National Chlamydia Screening Programme (NCSP) returns

8. The NCSP offers opportunistic chlamydia screening to those aged 15 to 24 years attending a variety of non-GUM clinic settings. NCSP figures therefore include diagnoses among 15 to 24 year olds only.
9. Data are presented by calendar year (Jan - Dec) and may not match with other data published by the NCSP which are usually reported by financial year (Apr - Mar).
10. Data presented are based on all tests with confirmed positive chlamydia test results including partners/contacts as well as index cases.

### "Non-NCSP / Non-GUM" chlamydia returns

11. The NCSP also collects data from laboratories on numbers of diagnoses made in those aged 15 to 24 years tested outside registered NCSP settings and GUM clinics.
12. Reporting of "Non-NCSP / non-GUM" returns began in April 2008, so there are no data available between January and March 2008.
13. Data are presented by calendar year (Jan - Dec) and may not match with other data published by the NCSP which are usually reported by financial year (Apr - Mar).



## 1 a: England

### Number of STI diagnoses, 2008 - 2010

New STI diagnoses *		2008	2009	2010	% change 2009-2010
Syphilis (primary, secondary and early latent)	Male (all)	2,508	2,497	2,318	-7%
	Female	366	340	291	-14%
	<b>Total</b>	<b>2,874</b>	<b>2,846</b>	<b>2,624</b>	<b>-8%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	9,820	10,695	11,439	7%
	Female	5,165	5,253	5,061	-4%
	<b>Total</b>	<b>14,985</b>	<b>15,978</b>	<b>16,531</b>	<b>3%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	71,061	74,568	76,863	3%
	Female	104,820	112,889	111,767	-1%
	<b>Total</b>	<b>178,001</b>	<b>189,625</b>	<b>189,612</b>	<b>0%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	67,565	63,412	58,694	-7%
	Female	21,276	21,676	21,164	-2%
	<b>Total</b>	<b>88,841</b>	<b>85,188</b>	<b>79,983</b>	<b>-6%</b>
Genital herpes simplex (first episode)	Male (all)	10,104	10,824	11,584	7%
	Female	15,990	16,703	18,090	8%
	<b>Total</b>	<b>26,094</b>	<b>27,564</b>	<b>29,703</b>	<b>8%</b>
Genital warts (first episode)	Male (all)	41,094	41,934	40,953	-2%
	Female	37,062	35,904	34,582	-4%
	<b>Total</b>	<b>78,156</b>	<b>77,900</b>	<b>75,615</b>	<b>-3%</b>
All other new STI diagnoses	Male (all)	15,973	15,300	14,444	-6%
	Female	11,973	10,350	10,055	-3%
	<b>Total</b>	<b>27,946</b>	<b>25,681</b>	<b>24,530</b>	<b>-4%</b>
<b>Total new STI diagnoses</b>	Male (all)	218,125	219,230	216,295	-1%
	Female	196,652	203,115	201,010	-1%
	<b>Total</b>	<b>416,897</b>	<b>424,782</b>	<b>418,598</b>	<b>-1%</b>
<b>Total other STI diagnoses</b>	Male (all)	125,040	126,117	128,701	2%
	Female	96,607	93,486	90,944	-3%
	<b>Total</b>	<b>221,647</b>	<b>219,906</b>	<b>220,049</b>	<b>0%</b>

\* Refer to:

'Code Groups' page for included conditions

'Title' page for data sources

'Notes' page for notes on using data tables



**1 b: East Midlands SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	125	136	145	7%
	Female	31	31	30	-3%
	<b>Total</b>	<b>156</b>	<b>167</b>	<b>175</b>	<b>5%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	548	644	574	-11%
	Female	413	459	438	-5%
	<b>Total</b>	<b>961</b>	<b>1,111</b>	<b>1,014</b>	<b>-9%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	5,349	5,609	6,019	7%
	Female	8,625	9,630	9,512	-1%
	<b>Total</b>	<b>15,002</b>	<b>16,130</b>	<b>15,627</b>	<b>-3%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	3,951	3,799	3,540	-7%
	Female	1,002	932	965	4%
	<b>Total</b>	<b>4,953</b>	<b>4,738</b>	<b>4,506</b>	<b>-5%</b>
Genital herpes simplex (first episode)	Male (all)	682	746	804	8%
	Female	1,042	1,101	1,249	13%
	<b>Total</b>	<b>1,724</b>	<b>1,850</b>	<b>2,055</b>	<b>11%</b>
Genital warts (first episode)	Male (all)	2,714	2,997	2,766	-8%
	Female	2,642	2,501	2,435	-3%
	<b>Total</b>	<b>5,356</b>	<b>5,510</b>	<b>5,212</b>	<b>-5%</b>
All other new STI diagnoses	Male (all)	1,000	910	846	-7%
	Female	847	683	683	0%
	<b>Total</b>	<b>1,847</b>	<b>1,593</b>	<b>1,530</b>	<b>-4%</b>
<b>Total new STI diagnoses</b>	Male (all)	14,369	14,841	14,694	-1%
	Female	14,602	15,337	15,312	0%
	<b>Total</b>	<b>29,999</b>	<b>31,099</b>	<b>30,119</b>	<b>-3%</b>
<b>Total other STI diagnoses</b>	Male (all)	6,661	5,835	5,615	-4%
	Female	5,924	5,343	5,135	-4%
	<b>Total</b>	<b>12,585</b>	<b>11,196</b>	<b>10,756</b>	<b>-4%</b>

\* Refer to:

'Code Groups' page for included conditions

'Title' page for data sources

'Notes' page for notes on using data tables



**1 c: East of England SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	98	76	83	9%
	Female	28	17	18	6%
	<b>Total</b>	<b>126</b>	<b>93</b>	<b>101</b>	<b>9%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	516	608	585	-4%
	Female	309	301	274	-9%
	<b>Total</b>	<b>825</b>	<b>909</b>	<b>859</b>	<b>-6%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	6,005	6,455	6,054	-6%
	Female	9,040	10,243	9,665	-6%
	<b>Total</b>	<b>15,099</b>	<b>16,766</b>	<b>15,753</b>	<b>-6%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	7,926	7,708	6,940	-10%
	Female	1,807	1,861	1,996	7%
	<b>Total</b>	<b>9,733</b>	<b>9,569</b>	<b>8,936</b>	<b>-7%</b>
Genital herpes simplex (first episode)	Male (all)	721	838	984	17%
	Female	1,365	1,433	1,656	16%
	<b>Total</b>	<b>2,086</b>	<b>2,271</b>	<b>2,640</b>	<b>16%</b>
Genital warts (first episode)	Male (all)	3,813	3,913	3,771	-4%
	Female	3,504	3,535	3,355	-5%
	<b>Total</b>	<b>7,317</b>	<b>7,448</b>	<b>7,126</b>	<b>-4%</b>
All other new STI diagnoses	Male (all)	1,347	1,207	1,244	3%
	Female	1,060	908	881	-3%
	<b>Total</b>	<b>2,407</b>	<b>2,115</b>	<b>2,125</b>	<b>0%</b>
<b>Total new STI diagnoses</b>	Male (all)	20,426	20,805	19,661	-5%
	Female	17,113	18,298	17,845	-2%
	<b>Total</b>	<b>37,593</b>	<b>39,171</b>	<b>37,540</b>	<b>-4%</b>
<b>Total other STI diagnoses</b>	Male (all)	9,043	9,126	9,226	1%
	Female	9,947	9,858	9,363	-5%
	<b>Total</b>	<b>18,990</b>	<b>18,984</b>	<b>18,589</b>	<b>-2%</b>

\* Refer to:

'Code Groups' page for included conditions

Title' page for data sources

Notes' page for notes on using data tables



**1 d: London SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	854	1,032	1,032	0%
	Female	99	99	97	-2%
	<b>Total</b>	<b>953</b>	<b>1,140</b>	<b>1,144</b>	<b>0%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	3,939	4,499	5,049	12%
	Female	1,741	1,676	1,703	2%
	<b>Total</b>	<b>5,680</b>	<b>6,197</b>	<b>6,781</b>	<b>9%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	13,630	15,173	16,636	10%
	Female	17,813	19,249	19,457	1%
	<b>Total</b>	<b>31,637</b>	<b>34,651</b>	<b>36,388</b>	<b>5%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	21,611	20,906	20,500	-2%
	Female	6,378	6,881	6,332	-8%
	<b>Total</b>	<b>27,989</b>	<b>27,875</b>	<b>26,954</b>	<b>-3%</b>
Genital herpes simplex (first episode)	Male (all)	2,447	2,726	2,856	5%
	Female	3,669	3,907	4,177	7%
	<b>Total</b>	<b>6,116</b>	<b>6,665</b>	<b>7,060</b>	<b>6%</b>
Genital warts (first episode)	Male (all)	7,567	7,792	7,615	-2%
	Female	6,080	5,991	5,849	-2%
	<b>Total</b>	<b>13,647</b>	<b>13,829</b>	<b>13,531</b>	<b>-2%</b>
All other new STI diagnoses	Male (all)	4,584	4,627	4,495	-3%
	Female	4,184	3,459	3,342	-3%
	<b>Total</b>	<b>8,768</b>	<b>8,114</b>	<b>7,866</b>	<b>-3%</b>
<b>Total new STI diagnoses</b>	Male (all)	54,632	56,755	58,183	3%
	Female	39,964	41,262	40,957	-1%
	<b>Total</b>	<b>94,790</b>	<b>98,471</b>	<b>99,724</b>	<b>1%</b>
<b>Total other STI diagnoses</b>	Male (all)	38,495	40,868	43,676	7%
	Female	26,109	25,889	26,169	1%
	<b>Total</b>	<b>64,604</b>	<b>67,026</b>	<b>70,229</b>	<b>5%</b>

\* Refer to:  
 'Code Groups' page for included conditions  
 'Title' page for data sources  
 'Notes' page for notes on using data tables



**1 e: North East SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	106	113	81	-28%
	Female	10	23	21	-9%
	<b>Total</b>	<b>116</b>	<b>136</b>	<b>102</b>	<b>-25%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	335	394	369	-6%
	Female	157	202	254	26%
	<b>Total</b>	<b>492</b>	<b>596</b>	<b>623</b>	<b>5%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	4,171	4,265	4,683	10%
	Female	5,488	5,935	6,465	9%
	<b>Total</b>	<b>10,283</b>	<b>10,734</b>	<b>11,336</b>	<b>6%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	2,150	2,219	1,975	-11%
	Female	454	495	462	-7%
	<b>Total</b>	<b>2,604</b>	<b>2,714</b>	<b>2,437</b>	<b>-10%</b>
Genital herpes simplex (first episode)	Male (all)	400	484	524	8%
	Female	599	735	783	7%
	<b>Total</b>	<b>999</b>	<b>1,219</b>	<b>1,307</b>	<b>7%</b>
Genital warts (first episode)	Male (all)	2,410	2,408	2,520	5%
	Female	2,255	2,206	2,134	-3%
	<b>Total</b>	<b>4,665</b>	<b>4,614</b>	<b>4,654</b>	<b>1%</b>
All other new STI diagnoses	Male (all)	732	688	642	-7%
	Female	337	284	280	-1%
	<b>Total</b>	<b>1,069</b>	<b>972</b>	<b>922</b>	<b>-5%</b>
<b>Total new STI diagnoses</b>	Male (all)	10,304	10,571	10,794	2%
	Female	9,300	9,880	10,399	5%
	<b>Total</b>	<b>20,228</b>	<b>20,985</b>	<b>21,381</b>	<b>2%</b>
<b>Total other STI diagnoses</b>	Male (all)	4,260	3,914	3,847	-2%
	Female	3,219	3,011	2,918	-3%
	<b>Total</b>	<b>7,479</b>	<b>6,925</b>	<b>6,765</b>	<b>-2%</b>

\* Refer to:  
 'Code Groups' page for included conditions  
 'Title' page for data sources  
 'Notes' page for notes on using data tables



**1 f: North West SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	482	390	374	-4%
	Female	38	39	28	-28%
	<b>Total</b>	<b>520</b>	<b>429</b>	<b>402</b>	<b>-6%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	1,399	1,238	1,441	16%
	Female	764	711	656	-8%
	<b>Total</b>	<b>2,163</b>	<b>1,949</b>	<b>2,097</b>	<b>8%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	11,577	11,798	11,462	-3%
	Female	18,076	18,962	18,409	-3%
	<b>Total</b>	<b>29,799</b>	<b>31,072</b>	<b>30,075</b>	<b>-3%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	6,106	6,459	5,644	-13%
	Female	1,722	1,838	1,664	-9%
	<b>Total</b>	<b>7,828</b>	<b>8,297</b>	<b>7,308</b>	<b>-12%</b>
Genital herpes simplex (first episode)	Male (all)	1,412	1,536	1,572	2%
	Female	2,404	2,471	2,473	0%
	<b>Total</b>	<b>3,816</b>	<b>4,007</b>	<b>4,045</b>	<b>1%</b>
Genital warts (first episode)	Male (all)	6,462	6,600	6,135	-7%
	Female	5,670	5,465	5,246	-4%
	<b>Total</b>	<b>12,132</b>	<b>12,066</b>	<b>11,381</b>	<b>-6%</b>
All other new STI diagnoses	Male (all)	2,175	2,130	1,865	-12%
	Female	1,328	1,125	1,101	-2%
	<b>Total</b>	<b>3,503</b>	<b>3,255</b>	<b>2,966</b>	<b>-9%</b>
<b>Total new STI diagnoses</b>	Male (all)	29,613	30,151	28,493	-5%
	Female	30,002	30,611	29,577	-3%
	<b>Total</b>	<b>59,761</b>	<b>61,075</b>	<b>58,274</b>	<b>-5%</b>
<b>Total other STI diagnoses</b>	Male (all)	18,059	16,210	16,477	2%
	Female	11,575	10,398	10,197	-2%
	<b>Total</b>	<b>29,634</b>	<b>26,608</b>	<b>26,676</b>	<b>0%</b>

\* Refer to:  
 'Code Groups' page for included conditions  
 'Title' page for data sources  
 'Notes' page for notes on using data tables



**1 g: South Central SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	98	99	90	-9%
	Female	11	17	11	-35%
	<b>Total</b>	<b>109</b>	<b>116</b>	<b>101</b>	<b>-13%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	444	455	483	6%
	Female	189	201	249	24%
	<b>Total</b>	<b>633</b>	<b>656</b>	<b>732</b>	<b>12%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	4,793	4,545	4,833	6%
	Female	6,592	6,511	6,722	3%
	<b>Total</b>	<b>11,391</b>	<b>11,077</b>	<b>11,571</b>	<b>4%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	5,527	5,148	4,745	-8%
	Female	1,689	1,865	1,780	-5%
	<b>Total</b>	<b>7,216</b>	<b>7,015</b>	<b>6,525</b>	<b>-7%</b>
Genital herpes simplex (first episode)	Male (all)	801	700	848	21%
	Female	1,218	1,192	1,361	14%
	<b>Total</b>	<b>2,019</b>	<b>1,892</b>	<b>2,209</b>	<b>17%</b>
Genital warts (first episode)	Male (all)	2,918	2,966	2,944	-1%
	Female	2,730	2,650	2,531	-4%
	<b>Total</b>	<b>5,648</b>	<b>5,617</b>	<b>5,475</b>	<b>-3%</b>
All other new STI diagnoses	Male (all)	1,112	1,058	947	-10%
	Female	657	632	524	-17%
	<b>Total</b>	<b>1,769</b>	<b>1,690</b>	<b>1,471</b>	<b>-13%</b>
<b>Total new STI diagnoses</b>	Male (all)	15,693	14,971	14,890	-1%
	Female	13,086	13,068	13,178	1%
	<b>Total</b>	<b>28,785</b>	<b>28,063</b>	<b>28,084</b>	<b>0%</b>
<b>Total other STI diagnoses</b>	Male (all)	8,317	9,069	9,639	6%
	Female	7,599	7,923	7,918	0%
	<b>Total</b>	<b>15,916</b>	<b>16,993</b>	<b>17,557</b>	<b>3%</b>

\* Refer to:  
 'Code Groups' page for included conditions  
 'Title' page for data sources  
 'Notes' page for notes on using data tables



**1 h: South East Coast SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	151	132	121	-8%
	Female	17	16	11	-31%
	<b>Total</b>	<b>168</b>	<b>148</b>	<b>132</b>	<b>-11%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	418	491	531	8%
	Female	156	180	145	-19%
	<b>Total</b>	<b>574</b>	<b>671</b>	<b>676</b>	<b>1%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	4,209	4,469	4,182	-6%
	Female	6,262	6,962	6,586	-5%
	<b>Total</b>	<b>10,483</b>	<b>11,445</b>	<b>10,768</b>	<b>-6%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	4,899	4,564	3,684	-19%
	Female	1,404	1,489	1,366	-8%
	<b>Total</b>	<b>6,303</b>	<b>6,055</b>	<b>5,050</b>	<b>-17%</b>
Genital herpes simplex (first episode)	Male (all)	769	821	912	11%
	Female	1,209	1,321	1,498	13%
	<b>Total</b>	<b>1,978</b>	<b>2,142</b>	<b>2,410</b>	<b>13%</b>
Genital warts (first episode)	Male (all)	3,126	3,376	3,330	-1%
	Female	2,969	2,979	2,880	-3%
	<b>Total</b>	<b>6,095</b>	<b>6,355</b>	<b>6,210</b>	<b>-2%</b>
All other new STI diagnoses	Male (all)	1,241	1,148	1,084	-6%
	Female	604	568	527	-7%
	<b>Total</b>	<b>1,845</b>	<b>1,716</b>	<b>1,611</b>	<b>-6%</b>
<b>Total new STI diagnoses</b>	Male (all)	14,813	15,001	13,844	-8%
	Female	12,621	13,515	13,013	-4%
	<b>Total</b>	<b>27,446</b>	<b>28,532</b>	<b>26,857</b>	<b>-6%</b>
<b>Total other STI diagnoses</b>	Male (all)	10,980	12,278	11,851	-3%
	Female	8,578	8,051	7,350	-9%
	<b>Total</b>	<b>19,558</b>	<b>20,330</b>	<b>19,201</b>	<b>-6%</b>

\* Refer to:  
 'Code Groups' page for included conditions  
 'Title' page for data sources  
 'Notes' page for notes on using data tables



**1 i: South West SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	177	114	88	-23%
	Female	32	18	16	-11%
	<b>Total</b>	<b>209</b>	<b>132</b>	<b>104</b>	<b>-21%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	508	555	480	-14%
	Female	259	269	188	-30%
	<b>Total</b>	<b>767</b>	<b>824</b>	<b>668</b>	<b>-19%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	6,451	6,073	6,274	3%
	Female	9,682	10,216	9,962	-2%
	<b>Total</b>	<b>16,143</b>	<b>16,301</b>	<b>16,249</b>	<b>0%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	5,871	4,311	4,332	0%
	Female	2,678	2,173	2,223	2%
	<b>Total</b>	<b>8,549</b>	<b>6,484</b>	<b>6,556</b>	<b>1%</b>
Genital herpes simplex (first episode)	Male (all)	898	913	966	6%
	Female	1,326	1,426	1,495	5%
	<b>Total</b>	<b>2,224</b>	<b>2,340</b>	<b>2,461</b>	<b>5%</b>
Genital warts (first episode)	Male (all)	3,808	3,714	3,803	2%
	Female	3,481	3,424	3,281	-4%
	<b>Total</b>	<b>7,289</b>	<b>7,139</b>	<b>7,085</b>	<b>-1%</b>
All other new STI diagnoses	Male (all)	1,203	1,122	1,019	-9%
	Female	696	637	612	-4%
	<b>Total</b>	<b>1,899</b>	<b>1,760</b>	<b>1,631</b>	<b>-7%</b>
<b>Total new STI diagnoses</b>	Male (all)	18,916	16,802	16,962	1%
	Female	18,154	18,163	17,777	-2%
	<b>Total</b>	<b>37,080</b>	<b>34,980</b>	<b>34,754</b>	<b>-1%</b>
<b>Total other STI diagnoses</b>	Male (all)	9,938	9,933	9,569	-4%
	Female	7,487	7,180	6,569	-9%
	<b>Total</b>	<b>17,425</b>	<b>17,115</b>	<b>16,141</b>	<b>-6%</b>

\* Refer to:  
 'Code Groups' page for included conditions  
 'Title' page for data sources  
 'Notes' page for notes on using data tables



**1 j: West Midlands SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	231	200	159	-21%
	Female	73	50	27	-46%
	<b>Total</b>	<b>304</b>	<b>250</b>	<b>186</b>	<b>-26%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	948	984	1,098	12%
	Female	653	698	593	-15%
	<b>Total</b>	<b>1,601</b>	<b>1,682</b>	<b>1,691</b>	<b>1%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	6,239	7,541	8,032	7%
	Female	9,974	11,735	11,718	0%
	<b>Total</b>	<b>16,235</b>	<b>19,336</b>	<b>19,817</b>	<b>2%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	5,593	4,752	4,059	-15%
	Female	1,958	2,162	2,037	-6%
	<b>Total</b>	<b>7,551</b>	<b>6,914</b>	<b>6,096</b>	<b>-12%</b>
Genital herpes simplex (first episode)	Male (all)	924	999	1,031	3%
	Female	1,502	1,541	1,698	10%
	<b>Total</b>	<b>2,426</b>	<b>2,541</b>	<b>2,729</b>	<b>7%</b>
Genital warts (first episode)	Male (all)	3,812	3,828	3,797	-1%
	Female	3,570	3,385	3,337	-1%
	<b>Total</b>	<b>7,382</b>	<b>7,214</b>	<b>7,134</b>	<b>-1%</b>
All other new STI diagnoses	Male (all)	1,308	1,149	1,174	2%
	Female	1,246	1,162	1,176	1%
	<b>Total</b>	<b>2,554</b>	<b>2,311</b>	<b>2,351</b>	<b>2%</b>
<b>Total new STI diagnoses</b>	Male (all)	19,055	19,453	19,350	-1%
	Female	18,976	20,733	20,586	-1%
	<b>Total</b>	<b>38,053</b>	<b>40,248</b>	<b>40,004</b>	<b>-1%</b>
<b>Total other STI diagnoses</b>	Male (all)	8,807	8,582	8,557	0%
	Female	7,965	7,748	7,854	1%
	<b>Total</b>	<b>16,772</b>	<b>16,331</b>	<b>16,411</b>	<b>0%</b>

\* Refer to:  
 'Code Groups' page for included conditions  
 'Title' page for data sources  
 'Notes' page for notes on using data tables



**1 k: Yorkshire & the Humber SHA**  
**Number of STI diagnoses, 2008 - 2010**

<b>New STI diagnoses *</b>		<b>2008</b>	<b>2009</b>	<b>2010</b>	<b>% change 2009-2010</b>
Syphilis (primary, secondary and early latent)	Male (all)	186	205	145	-29%
	Female	27	30	32	7%
	<b>Total</b>	<b>213</b>	<b>235</b>	<b>177</b>	<b>-25%</b>
Gonorrhoea (complicated and uncomplicated)	Male (all)	765	827	829	0%
	Female	524	556	561	1%
	<b>Total</b>	<b>1,289</b>	<b>1,383</b>	<b>1,390</b>	<b>1%</b>
Chlamydial infection (complicated and uncomplicated)	Male (all)	8,637	8,640	8,688	1%
	Female	13,268	13,446	13,271	-1%
	<b>Total</b>	<b>21,929</b>	<b>22,113</b>	<b>22,028</b>	<b>0%</b>
Non-specific genital infection (complicated and uncomplicated)	Male (all)	3,931	3,546	3,275	-8%
	Female	2,184	1,980	2,339	18%
	<b>Total</b>	<b>6,115</b>	<b>5,527</b>	<b>5,615</b>	<b>2%</b>
Genital herpes simplex (first episode)	Male (all)	1,050	1,061	1,087	2%
	Female	1,656	1,576	1,700	8%
	<b>Total</b>	<b>2,706</b>	<b>2,637</b>	<b>2,787</b>	<b>6%</b>
Genital warts (first episode)	Male (all)	4,464	4,340	4,272	-2%
	Female	4,161	3,768	3,534	-6%
	<b>Total</b>	<b>8,625</b>	<b>8,108</b>	<b>7,807</b>	<b>-4%</b>
All other new STI diagnoses	Male (all)	1,271	1,261	1,128	-11%
	Female	1,014	892	929	4%
	<b>Total</b>	<b>2,285</b>	<b>2,155</b>	<b>2,057</b>	<b>-5%</b>
<b>Total new STI diagnoses</b>	Male (all)	20,304	19,880	19,424	-2%
	Female	22,834	22,248	22,366	1%
	<b>Total</b>	<b>43,162</b>	<b>42,158</b>	<b>41,861</b>	<b>-1%</b>
<b>Total other STI diagnoses</b>	Male (all)	10,480	10,302	10,244	-1%
	Female	8,204	8,085	7,471	-8%
	<b>Total</b>	<b>18,684</b>	<b>18,398</b>	<b>17,724</b>	<b>-4%</b>

\* Refer to:  
 'Code Groups' page for included conditions  
 'Title' page for data sources  
 'Notes' page for notes on using data tables

## Supplementary Tables of Sexually Transmitted Infections in England



Code Groups
<b>New STI diagnoses</b>
Infectious and early latent syphilis
Gonorrhoea (uncomplicated and complicated)
Chlamydial infection (uncomplicated and complicated)
Non-specific genital infection (uncomplicated and complicated)
Genital herpes simplex (first episode)
Genital warts (first episode)
New HIV diagnosis *
Chancroid/lymphogranuloma venerum (LGV)/Donovanosis *
Molluscum contagiosum *
Trichomoniasis *
Scabies *
Pediculus pubis *
* All other new STI diagnoses
<b>Other STI diagnoses</b>
Congenital and other acquired syphilis
Recurrent genital herpes simplex
Recurrent and re-registered genital warts
Subsequent HIV presentations (including AIDS)
Ophthalmia neonatorum (chlamydial or gonococcal)
Epidemiological treatment of suspected STIs (syphilis, chlamydia, gonorrhoea, non-specific genital infection)